





Guérini in Delessert's Voyage dans l'Inde

Papilio delesserti

Alii subhyalinus, albi, venis maculisque nigris. Anticis
subsinuatis posterioris dentatis, angulo anuli lunula obsoleta flava.
11 cent. exp.

Semitransparent like Idea, 4 cross bands in disc. cell, large 3 ang. marginal
spots at end of veins; + row of smaller submarginal spots between veins
a larger spot near outer angle + another below end of disc. cell of upper.
Penang. 1 spec? in Hope Mus. - Oxford.

Euploea chloe

Alii interrufus fuscus, anticis base violaceo unicoloribus, omnibus utrinque
punctis marginalibus albi serie duplice digestis; subtus punctis discoloribus
albi et subviolaceis. Exp. 10½ cent.

Between Alcathoe + Cocita Godart. Pincang.

Argynnis caerulea.

Alii subrotundati, anticeis apice subconcavi, supra fulvi: anticeis apice et lucis flexuosi, posticeis lineis flexuosis punctisque nigri, maculis costatis duobus albis; subtus griseo fulvi, margaritaceo-virantibus, fascia media conum alba, maculari; extus recta, uteis dentata
Exp. 7½ cent. Malacea.

Vanessa ludoxea. Guer. (1840).

Alii anticeis subfalcatis, posticeis extus subcundatis; omnibus supra ochraceis basi fuscis apice nigris, posticeis lueis duabus marginatibus punctisque subcellaribus nigris; subtus brunnesc. ochracei strijs dentatis albis et fuscis; ocellisque cornu prospicillatis prope marginem.

Exp. 7 cent. Malacea.

very near Polymeria Gr. : a pale variety.

Qh
545
E513
v.1
Ent

F. Moore
Copy
100

A CATALOGUE
OF THE
LEPIDOPTEROUS INSECTS
IN

The Museum

THE HON. EAST-INDIA COMPANY.

BY

THOMAS HORSFIELD, M. & PH. D., F.R.S.,

Keeper of the Company's Museum,

AND

FREDERIC MOORE,

Assistant.

VOL. I.

Printed by Order of the Court of Directors.



LONDON:

WM. H. ALLEN AND CO.

Booksellers to the Hon. East-India Company,

7, LEADENHALL STREET.

1857.

199660

LONDON :

Cox and Wyman, Printers to the Hon. East-India Company,
74-5, GREAT QUEEN STREET, W.C.



P R E F A C E.

IT was my intention, as stated in the Introductory Remarks, to comprise in the first part of the Catalogue of the Lepidopterous Insects contained in the Museum of the East-India Company, the three first Tribes of this Order of Insects; but in arranging this work for publication, it became evident that our materials and the labour of preparing them for publication exceeded the original estimate of their extent: the full illustration of the species, the investigation and enumeration of the multifarious synonyms by which they are indicated by authors, together with the descriptions of a considerable number of new species, rendered a modification of our first plan advisable. I have therefore determined to limit the first part to the PAPILIONES and SPHINGES, reserving the BOMBYCES to the second part.

In submitting this Catalogue to the Entomological public, it is my first duty to acknowledge the very valuable and accommodating assistance which has been afforded to myself and my assistant by the officers of the Zoological Department of the British Museum, in our frequent researches, for the purpose of accurately comparing the subjects in the Company's with those contained in the British Museum. By this means, our species have been verified with a degree of accuracy otherwise unattainable; the synonyms carefully examined, and the character and distinctness of the new species fixed with all possible accuracy.

My best thanks are also due to A. Grote, Esq., of the Company's

Bengal civil service, who has placed at my disposal a very extensive series of drawings of the transformations of Lepidoptera, from continental India; the first portion, containing the *Diurna*, has already reached our Museum, and Plate XII. of illustrations consists almost entirely of Mr. Grote's contributions.

I have also to acknowledge thankfully the contribution of original drawings of transformations of Eastern Lepidoptera: *first*, by Edgar Leopold Layard, Esq., of drawings from Ceylon: *secondly*, by Lady Isabella Rose Gilbert, of drawings from continental India: *thirdly*, by Captain Mortimer Slater, of drawings from northern India: *fourthly*, the Entomological Society has also afforded access to the drawings made by Mrs. Hamilton.

The general plan according to which this work will be conducted is detailed in the Introductory Remarks. The indications afforded by the metamorphoses form the basis, and the subjects will be arranged, as far as possible, according to their affinities or most natural relations. This has been attempted in the first Tribe, the metamorphoses of which are illustrated on the first seven and the twelfth Plate, to which I refer the reader: but in a local collection of limited extent such an attempt must necessarily be imperfect.

The SPHinges, forming the second Tribe, are limited exclusively to those Insects in which the chrysalis is naked, and deposited on the surface or under the earth, the character of which is illustrated by Plates VIII., IX., X., and XI. These form, in the twelfth edition of the "Systema Naturae," the first, second, and part of the third section of the genus *Sphinx*, as defined by Linnaeus. The remainder of this genus, namely, part of the third and fourth sections, consisting chiefly of the genera *Ægeria* and *Zygæna*, belong with more propriety to the BOMBYCES, to which they are allied both in their transformations and in the perfect insect, the chrysalis being enveloped in a silky fabric or cocoon, in the manner of the *Bombyces*.

Thus restricted, the SPHinges are divided into five Stirpes,

which, with the exception of the first Stirps, the metamorphosis of which is as yet unknown, are named from the form of the larvae; viz., the *second* Stirps *Elongatæ*, the *third* Stirps *Acrocephalæ*, the *fourth* Stirps *Amblocephalæ*, and the *fifth* Stirps *Ophthalmicæ*. These are described in detail in pages 258-9. The special character of these divisions, exhibited in the metamorphosis, is strikingly confirmed by the perfect Insect, which, in each of these Stirpes, has a peculiarity obvious to every observer.

In conclusion, it is my duty to state that the descriptive part of this Catalogue has been performed by Mr. F. Moore, the Assistant in the Company's Museum ; and the experienced entomologist will recognize the skill and accuracy of Mr. J. O. Westwood in the preparation of the Illustrations.

THOMAS HORSFIELD.

December 7th, 1857.

INTRODUCTORY REMARKS.

THE ZOOLOGICAL MUSEUM of the East-India Company consists of specimens in all departments of the Science, from the Company's Oriental possessions, contributed by public servants who have been attached as Naturalists to Missions and Deputations on behalf of the Indian Government, or by gentlemen of the civil and military services, as presents to the Honourable Court of Directors.

Collections and Contributions in the particular branch of ENTOMOLOGY, have been received in the Museum in the following chronological order :—

1802. *Eudelin de Joinville*.—A series of Insects from Ceylon, chiefly Lepidoptera, presented to the Indian Government, on the transfer of Ceylon to the British Crown.
- 1813 to 1819. *Thomas Horsfield, M.D.*.—The Entomological Collections made in Java, during the possession of that island by Britain; consisting of a large series of Insects in all Orders, accompanied by Drawings of Lepidopterous Insects in all the stages of their existence.
1823. *George Finlayson*.—Insects collected during the Mission of John Crawford, Esq., to Siam and Hué, the capital of Cochinchina.
1829. *Madras Government*.—Collections made by the Company's Naturalist at Fort St. George.
1831. *Colonel W. H. Sykes*.—The Collections—consisting of a large series of Insects in all Orders, besides Birds, Mammalia, &c.—made during the Statistical Survey of the Dukhun.
1840. *Major R. B. Pemberton*.—Insects collected during his Mission to Bootan in 1837-38.

1841. *The Asiatic Society of Bengal*.—A series of Insects in all Orders.
John McClelland, Esq..—Specimens from Assam.
1842. *J. T. Pearson, Esq.*.—Specimens from Darjeeling, in all Orders.
The Bengal Government.—The Entomological Collection made in Chusan by Theodore Cantor, M.D., acting as Naturalist during the Chinese Expedition.
1843. *The Bombay Government*.—The Entomological Collections made during the mission of Sir W. C. Harris to Abyssinia.
1844. *The Asiatic Society of Bengal*.—Insects in all Orders.
- 1849 to 1856. *Ezra T. Downes, Esq., Deputy Assay-master, Bombay Mint*.—Large Collections of Insects in all Orders (accompanied by Notes), received by several despatches.
1849. *Colonel F. Buckley*.—A large Collection of Insects in all Orders, from the Himalayas, &c.
Brigadier-General J. B. Hearsey.—A small Series of Lepidoptera, received through J. O. Westwood, Esq.
1850. *Colonel W. H. Sykes*.—A Series of Insects preserved in spirit.
Captain Richard Strachey.—A Series of Insects from Ladakh.
1853. *B. H. Hodgson, Esq.*.—A Series of Insects from the central regions of Nepal.
1854. *Theodore Cantor, M.D.*.—A Collection of Insects in all Orders, from Penang.
1856. *J. C. Bowring, Esq.*.—A Series of Insects from North China.
Messrs. H. & R. Schlagintweit.—An extensive Collection of Insects in all Orders, from Darjeeling, &c.
The Bengal Government.—The Collection of Insects transmitted for exhibition at the Exposition Universelle at Paris in 1855.

In the Catalogue which is now offered to the public under the auspices of the Honourable Court of Directors of the East-India Company, it is intended to describe the Lepidopterous Insects contained in the Company's Museum. The Materials for this Work have been enumerated above in the chronological order of their contribution.

When I commenced a Systematic arrangement of the earlier Collections in 1820, soon after my arrival in England, they consisted, with the exception of a small number of specimens from Ceylon, exclusively of the Lepidopterous Insects from Java, which

were collected by me, and deposited in the Museum between the years 1813 and 1819. Since that period, collections of various extent have been contributed from the continental possessions of the Company in the East, and from China.

In the year 1820, the Entomological Cabinet was provisionally arranged according to the plan of Dr. Leach, as given in Samoelle's Entomologists' useful Compendium. My attention having been devoted to other branches of Zoology, chiefly Mammalia and Birds, the Insects remained in the state of their first arrangement until the year 1825, when William Sharp Macleay, Esq., commenced, in the "Annulosa Javanica," a description of the Coleopterous Insects in the Museum.

Some time after this, an attempt was made to give a description of the Lepidopterous Insects in a work entitled, "Descriptive Catalogue of the Lepidopterous Insects in the Company's Museum." This work having been undertaken on a plan which could not insure public support, was discontinued after the publication of the two first numbers, in 1828 and 1829.

In undertaking that work, the systematic arrangement to be adopted was the first consideration; and accordingly, the works of Linnaeus, Latreille, Fabricius, Duméril, and other systematic Entomologists, were carefully studied, in order to direct me to the most natural method of arranging my Collection.

While thus engaged, my attention was almost accidentally called to a comparatively obscure work, rarely consulted by English Entomologists (except as an Index of Species described), which was published in Germany towards the end of the last century, entitled, "Systematisches Verzeichniss der Schmetterlinge der Wienergegend, heraus-gegeben von einigen Lehrern am K. K.—THERESIANUM;" which last word I found designated Messrs. Denis and Schieffermüller, two distinguished officers of the court of Vienna, known in Germany from their charge in the Imperial Academy, by the name of "THERESIANER." In this work I discovered, to my great delight, an attempt to unite Lepidopterous Insects into groups, founded on the various stages of their existence, from the Caterpillar to the perfect Insect, or on their transformations. The maxim of the authors is, "*Ein Aug auf den Schmetterling, das andere Aug die Raupe*" (one eye to the Butterfly, another to the Caterpillar). This maxim gave a new direction to my inquiries; it superseded

any further examination of the systems above mentioned, *so far as relates to system and classification*, these works being founded almost exclusively on a single character, and accordingly artificial, while it greatly confirmed the views which had directed my researches in Java, as will appear in the sequel.

Although the Wiener Verzeichniss has been called a system, it can only be considered as an accumulation of materials of the metamorphoses of European Lepidoptera, to serve for comparison by Entomologists in other countries. The authors have not attempted a new Classification of the whole Order of Lepidoptera on their own principles. In the large divisions they adopt the Genera of Linnaeus, although they have transposed the Genera of that author, commencing with *Sphinx Phalæna*, and terminating with *Papilio*; the cause of this transposition is not explained. Mr. Westwood, in his "Introduction to the Modern Classification of Insects," gives a summary of the subdivisions and families into which the Wiener Verzeichniss is divided, with the remark, that almost every one of the divisions constitutes a modern genus. This work, although little known or regarded in England and France, is held in high estimation in Germany, where, at the period of its publication, it formed an epoch among cultivators of Entomology. Not many years after its first appearance, a new edition was required, which was undertaken on a somewhat reduced scale by the celebrated Illiger, who describes it as one of the most acute and instructive works on Entomology ever published; indeed, in its literary and philosophical character it may be ranked with the most perfect works on Natural Science extant at the time of its publication. It takes a most comprehensive view of Entomology in all its departments; it notices in detail all the authors and systems published at its appearance, near the end of the last century; and the research and learning displayed in its compilation are highly creditable to the authors. After stating the design of their work, the authors describe, in a manner perhaps more comprehensive and complete than is done in any other entomological treatise, the history and peculiarities of Lepidopterous Insects, from the egg to the Butterfly, interspersed with many original remarks, founded on their own observation. The description of the families and of the individual larvæ and chrysalides are given with the greatest minuteness. A series of notes and quotations is carried through the whole work,

furnishing the student with an abstract of all that has been recorded by the most eminent writers at the period above mentioned. The sixth and seventh sections contain many ingenious details and suggestions respecting the relative number of Lepidopterous Insects and Plants found in a given district or country ; and the authors endeavour to show that the number of both Butterflies and Plants will be found nearly equal in the Fauna and Flora of the countries that have been carefully examined. In examining the various systems extant at their time, they discuss the advantages and defects of each, with a view to prove that further researches were required to supply the vacancies remaining, in the attempt to form a natural system of arrangement, of which they had a more correct notion than most other naturalists of their day.

These sections also are invaluable in directing the pursuits of young Entomologists to the importance of collecting Lepidoptera in the various stages of their existence, both with the view to obtain perfect specimens and a complete history of the separate species. The style is colloquial, animated, and classical, and illustrated with numerous appropriate quotations in Latin, French, and English. An acquaintance with this work during my abode in Java would have enabled and directed me to carry on my researches in Entomology to a greater extent than I have accomplished.

In the systematic structure of this Catalogue, I shall be guided, so far as my materials afford the means, by the character in which the subjects manifest themselves in their transformations from the first state of their existence to their perfect development, or, according to the maxim of the authors of the Wiener Verzeichniss, from the larva to the imago ; endeavouring in all cases to discover the most natural arrangement in which they can be disposed.

It is obvious that a complete arrangement of the whole Order of Lepidopterous Insects on this principle cannot be attempted at the present period, although considerable materials have already been contributed towards this object from many parts of the world ; among these, the most important are contained in the following works :—

Hübner (Jacob).—Geschichte Europäische Schmetterlinge,
quammelt van J. H.

Abbott and Smith.—The History of the rarer Lepidopterous Insects of Georgia.

Cramer (Pierre).—Papillons Exotiques, etc. The Supplemental Volume by Stoll.

Merian (Madame).—Métamorphoses Insect. Surinamens.

Harris.—The Aurelian.

Boisduval and Léconte.—Hist. Générale et Iconogr. Lépidopt. et Chenilles de l'Amér. Sept.

Poey.—Centurie de Lépidopt. de l'Isle de Cuba.

Sepp.—Hist. Nat. de Papillons de Surinam.

But many extensive regions are as yet altogether unexplored, and a complete System of Lepidopterous Insects founded on their metamorphoses is reserved for some future Entomologist provided with more ample materials than have as yet been brought together: my only object in the present work is to contribute the result of my labours in the Eastern Islands.

Deeply impressed with the importance and necessity of an accurate knowledge of Lepidoptera in all stages of their existence, I devoted, during several successive seasons, all the means at my command to the investigation of Javanese Lepidoptera; and it may be not out of place here to give a brief extract from the Descriptive Catalogue above mentioned, of the method pursued by me. "I
"lived at this time at Surakarta, a province in the interior, be-
"longing to the native princes. I was amply provided with every
"convenience and facility for preserving what I had collected.
"Several draughtsmen had likewise been trained, under my super-
"intendence, for botanical delineations; and the skill they acquired
"in those, soon fitted them for the annulose department. I was,
"therefore, enabled to enter upon a history of the Metamorphoses
"of Javanese Lepidoptera, a design which had long engaged my
"anxious solicitude. Although I did not, at this period, so fully
"conceive the paramount necessity of an acquaintance with the
"Metamorphoses of Lepidoptera, towards the establishment of a
"natural arrangement, as I have been led to do in later periods,
"yet I was so strongly impressed with its essential importance in
"attempting a complete history of Insects, that I commenced with
"a fixed determination to prosecute the inquiry with unremitting
"industry and zeal, to collect all the larvæ of Lepidopterous Insects
"which I might possibly obtain, and to trace them through the

" various periods of their existence. With this view, I fitted up a
" large apartment adjoining my residence with breeding-cages and
" receptacles for chrysalides. At the commencement of the rainy
" season, the period when, in tropical climates, the foliage of
" vegetables is renewed, I daily went out in search of caterpillars,
" accompanied by the most intelligent of my native assistants. The
" caterpillars thus collected were placed in separate breeding-cages,
" and several of the assistants were instructed to provide daily, at
" regular periods, the food the individuals required, and to secure
" the cleanliness of the cages. As soon as the caterpillars were
" approaching to perfection, a drawing was made of them. The
" same individual which had been submitted to the draughtsman
" was then separately confined, watched with the most diligent
" care, and as soon as it had passed into the state of a chrysalis,
" again made the object of the pencil. A determinate number was
" carefully attached to the drawing and to the cage of the chrysalis.
" As soon as the perfect insect had appeared and expanded its
" wings, it was secured, set, and numbered, in accordance with the
" larva and chrysalis. During this period, every possible solicitude
" was employed to prevent mistakes. The original series, consisting
" of the perfect insects and the chrysalides obtained by this mode
" of proceeding, and numbered in accordance with the collection of
" drawings made at the same time, is now deposited in the Museum
" of the Honourable East-India Company, and affords an authentic
" document of the accuracy of the details regarding the metamor-
" phoses of Javanese Lepidoptera, which will be offered in the
" course of this work."

By the method thus described, I obtained the complete history of the transformations of nearly 200 species of Javanese Lepidoptera. The collection varies considerably in the principal divisions of this Order. In the first great division or Tribe, comprising the Diurnal Butterflies, or Papiliones, I have been most successful, and in this tribe my materials are sufficiently extensive to illustrate the minor Groups or Stirpes into which it has been subdivided; and in these are included also a large proportion of the Genera found in the tropical countries of the East.

Having above detailed the advantages I derived from an acquaintance with the Wiener Verzeichniss, so far as relates to the smaller groups of Lepidoptera, and the association of them into

Natural Genera on the principle employed by Messrs. Denis and Schieffermüller, I have now to refer to another work which opened a new view to my study of Zoology generally, and especially of Entomology,—the “*Horæ Entomologicæ*” of William Sharp Macleay, Esq.; and I consider it a fortunate coincidence that this work made its appearance about the period of my arrival in England, and that my attention was early directed to it,—the first part having been published in 1819, the second in 1821,—while I was engaged on the provisional arrangement of the Lepidopterous Insects in the Company’s Museum.

In this work, Mr. Macleay has sketched a plan, according to which the whole animal kingdom forms one natural group. In explanation of his system, he brings into one view almost all that is recorded by Zoological authors, both ancient and modern, in order to illustrate his theory. It is, in fact, an epitome of Zoology, and the study of it will greatly benefit the Zoologist and the Entomologist, whether he adopts or rejects his views. It is the production of an original mind, and may be justly considered as one of the most important contributions of late years to the Philosophy of Natural History.

Without attempting an analysis of the work, I shall briefly state those points which have a reference to this Catalogue.

First. The sentiment which pervades this work, impressing in the strongest manner on the student and the cultivator of Natural History, the paramount importance of the maxim of Linnæus: *METHODUS NATURALIS EST ULTIMUS FINIS BOTANICÆ.*

Secondly. The clearness with which relations of *Affinity* and *Analogy* observable in different groups of subjects of Natural History, both animal and vegetable, are propounded and explained. This law was discovered by Mr. Macleay in 1819, in studying Lamellicorn Insects; and was subsequently more fully discussed and elucidated in an Essay contained in the fourteenth volume of the Transactions of the Linnaean Society,* chiefly with reference to a similar discovery made almost simultaneously by Elias Fries; *Griphiswaldiae*, 1821, but without any communication or inter-

* Entitled, “Remarks on the Identity of certain General Laws which have been lately observed to regulate the Natural Distribution of Insects and Fungi.” By William Sharp Macleay, Esq.

change of opinions of the respective authors, in a group of the lower vegetables, *Fungi*, and which is made public in a work entitled "Systema Mycologicum, sistens Fungorum Ordines, Genera, et Species."*

Thirdly. The exposition of the law, that the series of affinities in groups of the animal kingdom is progressive, and that it returns into itself; and thus the groups form circles.

Fourthly. That the primary groups of those departments of the animal kingdom which have hitherto been investigated have been ascertained to be limited to five.

The laws enumerated in the third and fourth points have been discovered to prevail in *Mammalia* by Dr. J. E. Gray, detailed in the Annals of Philosophy, New Series, No. LIX., for November, 1823; in *Birds*, by N. A. Vigors, Esq., and have been detailed with great clearness in an admirable Essay contained in the fourteenth volume of the Transactions of the Linnaean Society, entitled, "Observations on the Natural Affinities that connect the Orders and Families of Birds;" and by Dr. J. J. Kaup, of Darmstadt, in an excellent monograph of *Falconidae*, the subdivisions of which agree with those of N. A. Vigors, Esq., independent of any mutual communication of the respective authors; in *Crustacea*, by Dr. De Haan, of Leyden, who, in the Introduction of his work has the following passage:—"Secuti sumus methodum circulorum quinariam ab viro ornato Macleay in Horis Entomologicis expositum, qua quippe via totum naturæ schema aptius exponi et affinitates facilius indicari nobis visæ sunt."†—*Pramissa IX.*

* Referring to Mr. Macleay's paper, contained in the Transactions of the Linnaean Society, here cited, it may be proper in this place to give the substance of the proposition of Fries in his own words:—"Affinia igitur sunt quæ in eadem serie sequuntur, et in se invicem transire videntur. Hæc in ulterioribus congruunt, sed in ceterioribus rationibus differunt. Analogia autem dicimus quæ in diversis seriebus locis parallelis posita sunt, et sibi invicem correspondent."

It is remarkable, that a similar law was noticed by Agardh, which in his *Aphorismi Botanici* is described in the following words:—"Analogia quedam et similitudo in diversis seriebus vegetabilium interdum cernatur, quasi progressa esset natura ad perfectionem per eosdem gradus sed diversâ viâ."

† In the second chapter, De ratione, qua quinque sectiones Crustaceorum inter se ligantur, he has the following remarks:—"Plures Botanici et Zoologi in eo conveniunt et vegetabilia et animalia non tantum plures series distinctas

Fifthly. That each group in a circle is analogically represented by the corresponding group in other circles.

Sixthly. The arrangement which Mr. Macleay has made of a class of annulose animals, designated Ametabola in his System ; and its application in the present work.*

Ametabola.—This Class, when first defined and introduced into the system by Dr. Leach, consisted of two divisions only,—Thysanura and Anoplura ; to these Mr. Macleay has added the two divisions of Myriapoda, namely Chilopoda and Chilognatha, and also Vermes, as defined by him, *Horæ Entom.* p. 286. The Class, therefore, consists of the following Orders :—Vermes, Anoplura, Thysanura, Chilopoda, and Chilognatha.

The researches and inquiries which led Mr. Macleay to this arrangement, are detailed in the same work, in pp. 286-7, &c. They are founded on a very careful study of the character of these animals, with reference especially to recent accurate analyses and descriptions of Messrs. Savigny, Latreille, and Lamarck, and they afford a favourable example of the method pursued by Mr. Macleay in his inquiries.

I have now, very briefly, to state the application which Mr. Macleay has made of the several divisions of the class Ametabola, in an analogical point of view, to the larvæ of Coleopterous Insects.

In the seventh chapter, treating of the tribes of Mandibulata, he gives an explanation, of which the following is the substance.—“ When reviewing the Coleoptera with a view to a natural division, I first threw the whole into those great natural groups which occurred to my sight, and, prosecuting my inquiries, at length, when

formare, quæ analogis jungantur, sed invicem connexa esse, quasi annulum annulo. Naturam non facere saltum, est adagium Linneanum ; multum autem abest, huncce nexus in omnibus partibus esse indicatum. Illud vero vinculum extricare videtur scopus, in quem omnes perquisitiones in Historia Naturali debeat attendere, ut inde pateat in creatis et consensus et ordo et sapientia, si talis præ humanis excellentia humanis verbis exprimi possit. In hoc propositum Cl. Macleay *Horæ Entomologicas* conserpsit ; eximio hocce in opere invenimus filum Ariadneum, quo ulterius persequendo, viam illo in labyrintho invenire possumus. Formas nempe in circulis in se redeuntibus disponere conatus est ; circularis illa dispositio nobis videtur naturæ maxime conveniens.”

* Mr. Macleay's four other primary divisions of the Annulosa being the Crustacea, Arachnida, Mandibulated, and Hanstellated insects.

thinking of their *larvæ*, I discovered that each of my groups, as far as my knowledge of them went, had a peculiar character,"—which he then briefly describes on p. 422.

Referring to the detailed description of the larvæ of Coleopterous Insects which he has detected, and arranged in five groups, it is sufficient for my present purpose to state, that the *first* group, examples of which are *Carabus* and *Dytiscus*, are represented analogically by the Chilopodiform Ametabola; the *second* type, of which *Scarabaeus* is an example, is represented analogically by the Chilognathiform Ametabola. The *third* type, of which *Curculio* and *Cerambyx* are examples, is represented analogically by the Vermiform Ametabola. The *fourth* type, of which *Coccinella* and *Chrysomela* are examples, is represented analogically by the Anopluriform Ametabola; and the *fifth* type, an example of which is *Meloe*, is represented by the Thysanuriform Ametabola.*

To guard against any misapprehension of the characters of the relations here pointed out, Mr. Macleay gives the following caution (see H. E. p. 423): "The distinction, however, between affinity and analogy, is perhaps nowhere in Entomology more necessary to be attended to than here; since, in terming larvae *Chilognathiformes* or *Chilopodiformes*, it is not meant that they are *Scolopendræ* or *Juli*, or even near to them in affinity, but only that they are so constructed that certain analogical circumstances attending them strongly remind us of these *Ametabola*."

Considering, therefore, the determination to which Mr. Macleay has arrived regarding the primary groups of Coleoptera, and their analogical relations with certain divisions of Ametabola, according to which the whole Order resolves itself into five natural divisions, I submitted the entire series of illustrations of the Metamorphoses of Indian Lepidoptera contained in the Company's Museum to a rigid examination; reserving the indications afforded by this examination, so far as regards the whole Order, to a future stage of this Catalogue; it is sufficient for my present purpose to state, that in the first Tribe of this Order, the Diurnal Lepidoptera, or Papiliones, I discovered five decided types of form, each of which

* To illustrate the five types of form observed in the first Tribe of Lepidoptera, the Papiliones, and the corresponding analogies with the Ametabola, I have prepared the accompanying plate exhibiting the larva and the analogous Ametabola.

manifested a clear analogy to a corresponding division of Ametabola; namely,—

First. A *larva* of a linear oblong form, attenuated at both ends, depressed or cylindrico-convex, of a sluggish appearance, with short and scarcely perceptible feet, distinctly marked above with transverse striae. A *pupa* nearly smooth, or with comparatively few protuberances, very obtuse at the anterior extremity, attached by the abdomen, braced, vertically suspended with the head upwards or tending forwards in a horizontal direction. Representing the *Vermiform* Ametabola.

Secondly. A *larva* of a cylindrical form, generally swelled or thickened at the fourth or fifth section of the body, attenuated towards the head and posterior extremity; in the typical genera naked, or covered with short, obtuse, fleshy protuberances; in the extreme genera, at the boundary of the neighbouring groups, covered with a close silky down, or with short scattered hair, most remarkably distinguished by a furcula or forked tentaculum, situated between the head and neck, which may be drawn back or thrust forward at pleasure. *Pupa* angulated and mostly tuberculated; in the typical genera, and in those at the confines of the first group, attached by the posterior extremity, braced, and vertically suspended with the head upwards; in the genera approaching the third stirps, perpendicularly suspended, according to the habits of that stirps. Representing the *Chilognathiform* or *Juliform* Ametabola.

Thirdly. A cylindrical *larva*, strikingly characterized by its terrific or threatening aspect, being covered with sharp, rigid, erect processes, often of great length, but diversified in the different subdivisions, arranged in regular longitudinal lines along the body of the larva, and beset with numerous diverging spines disposed in a verticillate manner. The attitude of the *pupa* is the reverse of that of the typical forms in the two former divisions: it is attached by the abdomen, while the head is directed perpendicularly downwards; in form it is greatly diversified. Representing the *Chilopodiform* or *Scolopendriform* Ametabola.

Fourthly. A lengthened cylindrical striated *larva*, somewhat smaller at each end, apparently of an agile habit, naked, or covered with a slight down, having a head of moderate size, armed with two erect spines, or provided with a movable shield, often of great

size, and beset with erect hornlike processes : but the chief characteristic of this division consists in two very strongly marked lengthened filiform or spinous appendages at the extremity of the abdomen. The *pupa* is attached by the tail and suspended perpendicularly, as in the last division ; its head is in general terminated by two points of various form and length. Representing the *Thysanuriform* Ametabola.

Fifthly. A cylindrical *larva*, nearly naked, with a very large head, often globular, and attached to the body by a long neck ; characteristically distinguished from the other subdivisions by its bluntness and abrupt termination behind ; the *pupa* being covered by a convoluted leaf. Representing the *Anopluriform* Ametabola.

These separate groups, agreeably to the plan of Mr. Macleay, in the “*Annulosa Javanica*,” will be denominated stirpes.

Tabular Review of the Larvae and their Analogies above enumerated.

1st Division, named according to the form of the larva	}\ Vermiform.
2nd Ditto	Juliform or Chilognathiform.
3rd Ditto	Scolopendriform or Chilopodiform.
4th Ditto	Thysanuriform.
5th Ditto	Anopluriform.

To answer the question which may arise, whether, in applying to one *Tribe* of Lepidoptera—the Papiliones—the same principle which Mr. Macleay has applied to the whole *Order* of Coleoptera, I have not departed from his scheme ? I offer the following considerations :—Mr. Macleay has divided the whole Animal Kingdom into quinary groups, which are again subdivided into groups of consecutively descending value : thus, the Animal Kingdom is divided in descending order into Sub-kingdom, Class, Order, Tribe, and Stirps. This arrangement is fully exemplified in the Diagram of the whole Animal Kingdom, on page 318 of the “*Horæ Entomologicæ*,” and in that of the sub-kingdom the “*Annulosa*,” on page 390 of the same work ; to which I refer the reader. The same law of groups of consecutively minor value has been observed to prevail in Birds by N. A. Vigors, Esq., and in Crustacea by Dr. De Haan, of Leyden, who thus expresses himself :—“*Methodum quinariam, cuius in hac invertebratorum classi specimen Entomologorum judicio submittimus, in aliis usque ad familias, in aliis*

usque ad genera persecuti sumus. Ita, secundum opus laudatum, Animalia in quinque subregna ; quinque classes dividuntur. Porro, secundum nostras observationes, Crustacea continent quinque Ordines, quinque Tribus, quinque Familias, quinque Genera."

In the foregoing introductory remarks, I have endeavoured to detail the principles which will direct me in the compilation of this Catalogue. The indications afforded by the several individuals described in all stages of their existence, will be carefully studied and applied. I have acknowledged the advantage derived from the examination of the "Horae Entomologicae" of Mr. Macleay, and especially from his remark, that "*considerations founded on the metamorphoses must ultimately produce the most natural plan of entomological arrangement.*" But I desire that it may be distinctly understood, that it is not my purpose to give an illustration of his particular system ; my object being to contribute a fragment towards a future enterprise of a more qualified entomologist. In a collection from a territory of limited geographical extent, large chasms remain to be supplied from more extensive localities. In the construction of my groups, I shall endeavour to discover the prominent forms, and to arrange them according to their affinities ; and where their analogies to the Ametabola are evident, I shall point them out ; but I shall not attempt to form them strictly on the model given by Mr. Macleay, or to pursue their division into typical and aberrant forms.

My reference to the works of Mr. Vigors on Birds, and to Dr. De Haan on Crustacea, is to afford a confirmation of Mr. Macleay's views, but not as an indication of my intention to pursue the subject in this Catalogue, to the same extent to which they have pursued it. The most natural division of the order of Lepidoptera, which may result from my examination, is reserved for a future stage of this work ; in the first part I propose to describe the Papiliones, Sphinges, and Bombyces.

THOMAS HORSFIELD.

LIBRARY, EAST-INDIA HOUSE,

February 11th, 1857.

CATALOGUE
OF
LEPIDOPTEROUS INSECTS.

SYSTEMATIC LIST

OF THE

GENERA AND SPECIES

OF

LEPIDOPTEROUS INSECTS

CONTAINED IN THE

MUSEUM OF THE EAST-INDIA COMPANY.

Class INSECTA.

Order LEPIDOPTERA.

Tribe I. PAPILIONES.

Stirps I.,

with VERMIFORM Larvæ.

Genus MILETUS, Hübner.

M. Symethus, Cramer Sp.	p. 18
— Boisduvali, Moore	19
— Horsfieldi, Moore	19

Genus PITHECOPS, Horsfield.

P. Hylax, Fabr. Sp.	19
---------------------	----

Genus POLYOMMATUS, Latr.

P. Akasa, Horsf.	20
— Puspa, Horsf.	20
— Laius, Cramer Sp.	21

Genus LYCENA, Fabricius.

L. Pluto, Fabr. Sp.	21
— Amyntas, Fabr. Sp.	22
— Aratus, Cramer Sp.	22
— Parrhasius, Fabr. Sp.	22
— Pandava, Horsf.	22

L. Cneus, Fabr. Sp.	p. 22
— Damoetes, Fabr. Sp.	23
— Kandarpa, Horsf.	23
— Boëtica, Linn. Sp.	23
— Pavana, Horsf.	24
— Elpis, Godart Sp.	24
— Celeno, Cramér Sp.	24
— Alexis, Stoll Sp.	25
— Plinius, Fabr. Sp.	25
— Theophrastus, Fabr. Sp.	25
— Rosimon, Fabr. Sp.	26
— Malaya, Horsf.	26
— Roxus, Godart Sp.	26
— Pseudoroxus, Doubleday	27
— Nyseus, Guérin Sp.	27

Genus CHRYSOPHANUS, Hüb.

C. Phlæas, Linn. Sp.	27
— Pavana, Kollar Sp.	28

Genus ILERDA, Doubleday.

I. Sena, Kollar, Sp.	28
— Tamu, Kollar Sp.	28
— Androcles, Boisd. Sp.	29
— Brahma, Moore	29
— Epicles, Godart Sp.	29

Genus THECLA, Fabricius.

T. Onyx, Boisd.	30
-----------------	----

SYSTEMATIC LIST

Genus DIPSAS, <i>Doubleday.</i>		M. Acte, <i>Doubleday</i> p. 47
D. Syla, <i>Kollar Sp.</i>	p. 30	— Tharis, <i>Hübner Sp.</i> 47
— Ataxus, <i>Boisd. Sp.</i>	31	— Lapithis, <i>Boisd.</i> 48
— Xenophon, <i>Fabr. Sp.</i> ...	31	— Lisias, <i>Fabr. Sp.</i> 48
— Melampus, <i>Cramer Sp.</i> ...	32	— Triopas, <i>Cramer Sp.</i> 49
— Epijarbas, <i>Boisd. Sp.</i> ...	32	— Etolus, <i>Fabr. Sp.</i> 49
— Chrysomallus, <i>Hüb. Sp.</i>	33	— Amyntor, <i>Herbst Sp.</i> ... 49
— Isocrates, <i>Fabr. Sp.</i>	33	— Erylus, <i>Godart</i> 50
— Malika, <i>Horsf. Sp.</i>	37	— Jalindra, <i>Horsf. Sp.</i> 50
— Kessuma, <i>Horsf. Sp.</i> ...	37	— Nedymond, <i>Cramer Sp.</i> ... 50
— Nasaka, <i>Horsf. Sp.</i>	37	— Chitra, <i>Horsf. Sp.</i> 51
— Varuna, <i>Horsf. Sp.</i>	37	— Ravindra, <i>Horsf.</i> 51
Genus APHNÆUS, <i>Hübner.</i>		— Sugriva, <i>Horsf.</i> 51
A. Pindarus, <i>Fabr. Sp.</i>	37	Genus LOXURA, <i>Horsfield.</i>
— Etolus, <i>Cramer Sp.</i>	38	L. Atymnus, <i>Cramer Sp.</i> ... 51
— Lohita, <i>Horsf. Sp.</i>	38	— Surya, <i>Moore</i> 52
— Syama, <i>Horsf. Sp.</i>	38	— Pita, <i>Horsf.</i> 52
Genus AMBLYPODIA, <i>Horsf.</i>		Genus ANOPS, <i>Boisduval.</i>
A. Narada, <i>Horsf.</i>	39	A. Thetys, <i>Drury Sp.</i> 52
— Vivarna, <i>Horsf.</i>	39	— insularis, <i>Horsf. Sp.</i> 53
— Apidanus, <i>Cramer Sp.</i> ...	39	— Cinyra, <i>Cramer Sp.</i> 53
— Centaurus, <i>Fabr. Sp.</i>	40	— Bulis, <i>Boisd.</i> 53
— Pseudo-Centaurus, <i>Doubl.</i>	40	— Santana, <i>Moore</i> 54
— Silhetensis, <i>Boisd. Sp.</i> ...	40	Stirps II.,
— Helus, <i>Cramer Sp.</i>	40	with CHILOGNATHIFORM
— Camdeo, <i>Doubleday</i>	41	or JULIFORM Larvæ.
— Eumolphus, <i>Cramer Sp.</i>	41	A.
— Rama, <i>Kollar Sp.</i>	42	Genus CALLIDRYAS, <i>Boisd.</i>
— Perimuta, <i>Boisd.</i>	42	C. Pyranthe, <i>Linn. Sp.</i> 56
— Epimuta, <i>Boisd.</i>	42	— Philippina, <i>Cramer Sp.</i> ... 56
— Quercetorum, <i>Boisd.</i> ...	42	— Alcmeone, <i>Cramer Sp.</i> ... 56
— Querceti, <i>Boisd.</i>	43	— Hilaria, <i>Cramer Sp.</i> 57
— Dodonea, <i>Boisd.</i>	43	— Scylla, <i>Linn. Sp.</i> 58
— Ganesa, <i>Moore</i>	44	Genus GONEPTERYX, <i>Leach.</i>
— Timoleon, <i>Stoll Sp.</i>	44	G. Nipalensis, <i>Doubleday</i> ... 59
— Rochana, <i>Horsf.</i>	44	Genus DERCAS, <i>Boisduval.</i>
— Vidura, <i>Horsf.</i>	45	D. Verhuelli, <i>Van der Hoeven</i> Sp. 59
— Longinus, <i>Fabr. Sp.</i>	45	Genus HEBOMOIA, <i>Hübner.</i>
— Pseudolonginus, <i>Doubl.</i>	45	H. Glaucippe, <i>Linn. Sp.</i> ... 60
— Hypatada, <i>Boisd.</i>	45	
— Nissa, <i>Kollar Sp.</i>	46	
— Deva, <i>Moore</i>	46	
— Jangala, <i>Horsf.</i>	46	
Genus MYRINA, <i>Godart.</i>		
M. Jafra, <i>Godart</i>	47	

OF THE GENERA AND SPECIES.

Genus ERONIA, <i>Hübner.</i>	
E. Valeria, <i>Cramer Sp.</i> ...	p. 61
— Avatar, <i>Moore</i>	61
Genus COLIAS, <i>Fabricius.</i>	
C. Edusa, <i>Fabr. Sp.</i>	62
— Hyale, <i>Linn. Sp.</i>	62
— Neriene, <i>Fischer</i>	62
Genus TERIAS, <i>Swainson.</i>	
T. Hecabe; <i>Linn. Sp.</i>	63
— Sari, <i>Horsf.</i>	64
— Tilaha, <i>Horsf.</i>	64
— Blanda, <i>Boisd.</i>	64
— Harina, <i>Horsf.</i>	64
— Læta, <i>Boisd.</i>	65
— Venata, <i>Moore</i>	65
— Drona, <i>Horsf.</i>	65
Genus THESTIAS, <i>Boisduval.</i>	
T. Ænippe, <i>Cramer Sp.</i>	66
— Marianne, <i>Cramer Sp.</i> ...	66
— Pyrene, <i>Linn. Sp.</i>	66
— Venilia, <i>Godart Sp.</i>	67
Genus IDMAIS, <i>Boisduval.</i>	
I. Calais, <i>Cramer Sp.</i>	67
— Phisadia, <i>Godart Sp.</i> ...	68
— Dynamene, <i>Klug Sp.</i> ...	68
— Fausta, <i>Olivier Sp.</i>	68
Genus CALLOSUNE, <i>Doubleday.</i>	
C. Eucharis, <i>Fabr. Sp.</i>	68
— Danaë, <i>Fabr. Sp.</i>	69
— Etrida, <i>Boisd. Sp.</i>	69
Genus PONTIA, <i>Fabricius.</i>	
P. Nina, <i>Fabr.</i>	69
Genus PIERIS, <i>Schrink.</i>	
P. Nero, <i>Fabr. Sp.</i>	70
— Rouxi, <i>Boisd.</i>	71
— Panda, <i>Godart</i>	71
— Hippo, <i>Cramer Sp.</i>	71
— Nerissa, <i>Fabr. Sp.</i>	72
— Ega, <i>Boisd.</i>	72
— Neombo, <i>Boisd.</i>	72
P. Mesentina, <i>Cramer Sp.</i> p. 72	
— Libythea, <i>Fabr. Sp.</i>	73
— Pandione, <i>Hübner Sp.</i> ...	73
— Paulina, <i>Cramer Sp.</i>	73
— Durvasa, <i>Moore</i>	73
— Indra, <i>Moore</i>	74
— Nipalensis, <i>G. R. Gray</i>	75
— Gliciria, <i>Cramer Sp.</i>	75
— Daplidice, <i>Linn. Sp.</i>	75
— Remba, <i>Moore</i>	75
— Nama, <i>Doubleday</i>	76
— Coronis, <i>Cramer Sp.</i>	76
— Judith, <i>Fabr. Sp.</i>	77
— Lea, <i>Doubleday</i>	77
— Thestyliis, <i>Doubleday</i> ...	78
— Seta, <i>Moore</i>	78
— Belladonna, <i>Fabr. Sp.</i> ...	78
— Sanaca, <i>Moore</i>	79
— Crithoë, <i>Boisd.</i>	79
— Egialea, <i>Cramer Sp.</i>	79
— Pasithoë, <i>Linn. Sp.</i>	79
— Thisbe, <i>Cramer Sp.</i>	80
— Eucharis, <i>Drury Sp.</i>	80
— Hyparete, <i>Linn. Sp.</i>	80
— Hierte, <i>Hübner Sp.</i>	81
— Agostina, <i>Hewitson</i>	81
— Descombesi, <i>Boisd.</i>	81
— Belisama, <i>Cramer Sp.</i> ...	82
— Autothisbe, <i>Hübner Sp.</i>	82
— Coronea, <i>Cramer Sp.</i> ...	82
— Vishnu, <i>Moore</i>	83
Genus APORIA, <i>Hübner.</i>	
A. Agathon, <i>G. R. Gray</i> ...	83
— Soracta, <i>Boisd.</i>	83
B.	
Genus PARNASSIUS, <i>Latr.</i>	
P. Hardwicki, <i>G. R. Gray</i>	84
Genus LEPTOCIRCUS, <i>Swains.</i>	
L. Curius, <i>Fabr. Sp.</i>	85
— Meges, <i>Zinken-Sommer</i>	85
Genus TEINOPALPUS, <i>Hope.</i>	
T. imperialis, <i>Hope</i>	86

SYSTEMATIC LIST

Genus ORNITHOPTERA, Boisd.	P. Gyas, <i>Westwood</i> p. 110
O. Richmondia, <i>G. R. Gray</i> , p. 86	— Payeni, <i>Boisd.</i> 110
— Darsius, <i>G. R. Gray</i> ... 87	— Xuthus, <i>Linn.</i> 111
— Pompeus, <i>Cramer Sp.</i> ... 87	— Machaon, <i>Linn.</i> 111
— Rhadamaanthus, <i>Boisd.</i> ... 88	— Cloanthus, <i>Westwood</i> ... 112
— Amphrisius, <i>Cramer Sp.</i> 88	— Sarpedon, <i>Linn.</i> 113
Genus PAPILIO, Linnæus.	— Eurypylus, <i>Linn.</i> 113
P. Nox, <i>Swainson</i> 89	— Bathycles, <i>Zinken-Som.</i> 114
— Astina, <i>Horsf.</i> 89	— Agamemnon, <i>Linn.</i> 114
— Macareus, <i>Godart</i> 90	— Nomius, <i>Esper</i> 115
— Megarus, <i>Westwood</i> 90	— Anticrates, <i>Doubleday</i> ... 115
— Xenocles, <i>Doubleday</i> 90	— Antiphates, <i>Cramer</i> 116
— Delesserti, <i>Guérin</i> 91	— Agetes, <i>Westwood</i> 116
— Agestor, <i>G. R. Gray</i> ... 91	— Glycerion, <i>G. R. Gray</i> ... 117
— dissimilis, <i>Linn.</i> 91	
— Castor, <i>Westwood</i> 93	
— Hector, <i>Linn.</i> 93	
— Diphilus, <i>Esper</i> 94	
— Antiphus, <i>Fabr.</i> 94	
— Coon, <i>Fabr.</i> 95	
— Alcinous, <i>Klug</i> 95	
— Philoxenus, <i>G. R. Gray</i> 96	
— Dasarada, <i>Moore</i> 96	
— Ravana, <i>Moore</i> 96	
— Minereus, <i>G. R. Gray</i> ... 97	
— Janaka, <i>Moore</i> 97	
— Varuna, <i>White</i> 98	
— Rhetenor, <i>Westwood</i> 98	
— Protenor, <i>Cramer</i> 98	
— Memnon, <i>Linn.</i> 99	
— Polymnestor, <i>Cramer</i> ... 101	
— Iswara, <i>White</i> 101	
— Helenus, <i>Linn.</i> 101	
— Chaon, <i>Westwood</i> 102	
— Romulus, <i>Cramer</i> 102	
— Polytes, <i>Linn.</i> 103	
— Pammon, <i>Linn.</i> 104	
— Demolion, <i>Cramer</i> 105	
— Erithonius, <i>Cramer</i> 105	
— Arjuna, <i>Horsf.</i> 107	
— Paris, <i>Linn.</i> 107	
— Ganesa, <i>Doubleday</i> 107	
— Arcturus, <i>Westwood</i> 108	
— Krishna, <i>Moore</i> 108	
— Polycitor, <i>Boisd.</i> 109	
— Crino, <i>Fabr.</i> 109	
— Peranthus, <i>Fabr.</i> 109	
— Bianor, <i>Cramer</i> 110	
	Stirps III., with CHILOPODIFORM or SCOLOPENDRIFORM Larvæ.
	Genus DANAI, Latreille.
D. Limniae, <i>Cramer Sp.</i> 121	
— Melissa, <i>Cramer Sp.</i> 121	
— similis, <i>Linn. Sp.</i> 122	
— Grammica, <i>Boisd.</i> 122	
— Aglea, <i>Cramer Sp.</i> 122	
— Juventa, <i>Cramer Sp.</i> 122	
— Cleona, <i>Cramer Sp.</i> 123	
— Melaneus, <i>Cramer Sp.</i> ... 123	
— Tytia, <i>G. R. Gray</i> 123	
— Ablata, <i>Zinken-Som. Sp.</i> 124	
— Philene, <i>Cramer Sp.</i> 124	
— Plexippus, <i>Linn. Sp.</i> ... 124	
— Melanippus, <i>Cramer Sp.</i> 125	
— Lotis, <i>Cramer Sp.</i> 125	
— Chrysippus, <i>Linn. Sp.</i> ... 126	
	Genus EUPLÆA, Fabricius.
E. Rhadamanthus, <i>Fabr.</i> ... 126	
— Mitra, <i>Moore</i> 127	
— Elusine, <i>Cramer Sp.</i> 127	
— Mazares, <i>Doubleday</i> 127	
— Melina, <i>Godart Sp.</i> 128	
— Hübneri (<i>Boisd.</i>), <i>Moore</i> 128	
— Crameri (<i>Boisd.</i>), <i>Moore</i> 129	
— Eunice, <i>Godart Sp.</i> 129	
— Klugi (<i>Boisd.</i>), <i>Moore</i> ... 130	
— Prothoë, <i>Godart Sp.</i> 130	

OF THE GENERA AND SPECIES.

E. superba, <i>Herbst Sp.</i> ... p. 131	Genus PRECIS, <i>Hübner.</i>
— Alcaethoë, <i>Godart Sp.</i> ... 131	P. Ida, <i>Cramer Sp.</i> p. 142
— Core, <i>Cramer Sp.</i> 131	— Iphita, <i>Cramer Sp.</i> 143
— Deione, <i>Westwood</i> 132	— Hara, <i>Moore</i> 143
— Ochsenheimeri (<i>Boisd.</i>), Moore 132	— Polynice, <i>Cramer Sp.</i> ... 144
— Midamus, <i>Linn. Sp.</i> 133	Genus ERGOLIS, <i>Boisduval.</i>
Genus IDEOPSIS, <i>Horsfield.</i>	E. Ariadne, <i>Linn. Sp.</i> 144
I. Gaura, <i>Horsf.</i> 134	— Coryta, <i>Cramer Sp.</i> 144
— Daos, <i>Boisd. Sp.</i> 134	Genus CYNTHIA, <i>Fabricius.</i>
Genus HESTIA, <i>Hübner.</i>	C. Arsinoë, <i>Cramer Sp.</i> 145
H. Lynceus, <i>Drury Sp.</i> 134	Genus AMNOSIA, <i>Boisduval.</i>
— Belia, <i>Westwood</i> 135	A. decora, <i>Boisd.</i> 146
— Jasonia, <i>Westwood</i> 135	Genus CYRESTIS, <i>Boisduval.</i>
— Leuconoë, <i>Erickson</i> 135	C. lutea, <i>Zinken-Sommer</i> ... 146
Genus TELCHINIA, <i>Hübner.</i>	— Thyodamas, <i>Boisd.</i> 146
T. Violæ, <i>Fabr. Sp.</i> 135	— Periander, <i>Fabr. Sp.</i> 147
Genus PAREBA, <i>Doubleday.</i>	— Risa, <i>Doubleday</i> 147
P. Vesta, <i>Fabr. Sp.</i> 136	— Rahria, <i>Westwood</i> 147
Genus VANESSA, <i>Fabricius.</i>	Genus PARTHENOS, <i>Hübner.</i>
V. Antiopa, <i>Linn. Sp.</i> 136	P. Sylvia, <i>Cramer Sp.</i> 147
— Xanthomelas, <i>Denis et</i> <i>Schiff.</i> 137	— Gambrisius, <i>Fabr. Sp.</i> ... 148
— Cashmirensis, <i>Kollar</i> ... 137	Genus PROTHOË, <i>Hübner.</i>
— Charonia, <i>Drury Sp.</i> 137	P. Francki, <i>Godart</i> 148
Genus GRAPTA, <i>Kirby.</i>	Genus TERINOS, <i>Boisduval.</i>
G. C. aureum, <i>Linn. Sp.</i> ... 138	T. Clarissa, <i>Boisd.</i> 149
Genus PYRAMEIS, <i>Hübner.</i>	Genus CIRROCHROA, <i>Doubleday.</i>
P. Callirhoe, <i>Hübner</i> 138	C. Thais, <i>Fabr. Sp.</i> 149
— Cardui, <i>Linn. Sp.</i> 138	— Aoris, <i>Doubleday</i> 149
Genus JUNONIA, <i>Hübner.</i>	— Clagia, <i>Godart Sp.</i> 149
J. Lemonias, <i>Linn. Sp.</i> 139	— Bajadeta, <i>Moore</i> 150
— Erigone, <i>Cramer Sp.</i> 139	— Ravana, <i>Moore</i> 150
— Laomedia, <i>Linn. Sp.</i> 140	Genus MESSARAS, <i>Doubleday.</i>
— Oenone, <i>Linn. Sp.</i> 140	M. Erymanthis, <i>Drury Sp.</i> 150
— Orithyia, <i>Linn. Sp.</i> 141	— Alcipe, <i>Cramer Sp.</i> ... 151
— Vellida, <i>Linn. Sp.</i> 141	Genus ATELLA, <i>Doubleday.</i>
— Asterie, <i>Linn. Sp.</i> 142	A. Phalanta, <i>Drury Sp.</i> 151
— Almana, <i>Linn. Sp.</i> 142	— Egista, <i>Cramer Sp.</i> 152

SYSTEMATIC LIST

Genus LAOGONA, Boisduval.	N. Nandina, Moore p. 168
L. Hypoocla, Cramer Sp. p. 153	— Aceris, Esper Sp. 168
— Hypselis, Godart 153	
Genus CETHOSIA, Fabricius.	Genus ATHYMA, Westwood.
C. Biblis, Drury Sp. 153	A. Leucothoë, Linn. Sp. ... 170
— Penthesilea, Cramer Sp. 154	— Asura, Moore 171
— Æole, De Haan 154	— Opalina, Kollar 171
— Cyane, Fabr. Sp. 155	— Larymna, Doubleday ... 172
Genus ARGYNNIS, Fabricius.	— Jina, Moore 172
A. Niphe, Linn. Sp. 155	— Pravara, Moore 173
— Aruna, Moore 156	— Nefte, Cramer Sp. 173
— Issaea, G. R. Gray 156	— Inara, Doubleday Sp. ... 173
— Kamala, Moore 156	— Cama, Moore 174
— Rudra, Moore 157	— Selenophora, Kollar 175
— Childreni, G. R. Gray... 157	— Ranga, Moore 175
Genus DIADEMA, Boisduval.	— Mahesa, Moore 176
D. Bolina, Linn. Sp. 158	Genus ABROTA, Moore.
— Auge, Cramer Sp. 158	A. Ganga, Moore 178
— Antilope, Cramer Sp. ... 159	Genus LIMENITIS, Fabricius.
Genus PENTHEMA, Doubleday.	L. Procris, Cramer Sp. 179
P. Lisarda, Doubleday 160	— Zulema, Doubleday 179
Genus HESTINA, Westwood.	— Alankara, Horsf. Sp. ... 179
H. assimilis, Linn. Sp. 160	— Paduka, Moore 179
— consimilis, Westwood ... 161	— Ismene, Doubleday 180
— Isa, Moore..... 161	— Daraxa, Doubleday 180
— persimilis, Westwood.... 161	— Dudu, Westwood 180
— Nama, Doubleday 162	— Danava, Moore 180
Genus CALINAGA, Moore.	— Zayla, Doubleday 181
C. Buddha, Moore 163	Genus PANDITA, Moore.
Genus NEPTIS, Fabricius.	P. Sinope (Boisd.), Moore 182
N. Hordonia, Stoll Sp. 164	Stirps IV., with
— Heliodore, Fabr. Sp..... 164	THYSANURIFORM Larvæ.
— Miah, Moore..... 164	
— Vikasi, Horsf. 165	Genus ADOLIAS, Boisduval.
— Manasa, Moore..... 165	A. Aconthea, Cramer Sp. ... 185
— Ananta, Moore 166	— Parta, Moore..... 185
— Radha, Moore 166	— Garuda, Moore 186
— Columella, Cramer Sp.... 166	— Phemius, Doubleday 187
— Jumba, Moore 167	— Anosia (Boisd.), Moore 187
— Nata, Moore 168	— Alpheda, Godart Sp. ... 188

OF THE GENERA AND SPECIES.

- | | |
|---|---|
| A. Pulasara, <i>Moore</i> <i>p.</i> 190 | Genus NYMPHALIS, <i>Latreille</i> . |
| — Puseda, <i>Moore</i> 191 | N. Fabius, <i>Fabr. Sp.</i> <i>p.</i> 204 |
| — Trigerta, <i>Moore</i> 191 | — Athamas, <i>Drury Sp.</i> 205 |
| — Iapis, <i>Godart Sp.</i> 192 | — Schreiberi, <i>Godart</i> 205 |
| — Ambalika, <i>Moore</i> 192 | — Delphis, <i>Doubleday Sp.</i> ... 206 |
| — Jahnii, <i>Moore</i> 192 | — Eudamippus, <i>Doubl. Sp.</i> ... 206 |
| — Sikandi, <i>Moore</i> 193 | — Dolon, <i>Westwood</i> 206 |
| — Evelina, <i>Stoll Sp.</i> 193 | — Marmax, <i>Westwood</i> 206 |
| — Coctyna, <i>Horsf. Sp.</i> 193 | — Bernardus, <i>Fabr. Sp.</i> ... 206 |
| — Cocytus, <i>Fabr. Sp.</i> 194 | — Baya, <i>Moore</i> 207 |
| — Adima, <i>Moore</i> 194 | |
| — Apiades, <i>Menetries</i> 195 | Genus KALLIMA, <i>Doubleday</i> . |
| — Sancara, <i>Moore</i> 195 | K. Paralekta, <i>Horsf. Sp.</i> ... 208 |
| — Iva, <i>Moore</i> 195 | — Inachis, <i>Boisd.</i> 208 |
| — Epiona, <i>G. R. Gray</i> 196 | — Bisaltide, <i>Cramer Sp.</i> ... 209 |
| — Durga, <i>Moore</i> 196 | |
| — Teuta, <i>Doubleday</i> 196 | Genus AMATHUSIA, <i>Fabricius</i> . |
| — Franciae, <i>G. R. Gray</i> ... 197 | A. Phidippus, <i>Linn. Sp.</i> ... 209 |
| — Coresia, <i>Hübner Sp.</i> 197 | — Amythaon, <i>Doubleday</i> ... 210 |
| — Nicea, <i>G. R. Gray</i> 197 | |
| — Nesiomachus, <i>Boisd.</i> 197 | Genus ZEUXIDIA, <i>Hübner</i> . |
| — Dirtea, <i>Fabr. Sp.</i> 198 | Z. Luxeri, <i>Hübner</i> 210 |
| — Siva, <i>Westwood</i> 198 | |
| Genus EURIPUS, <i>Doubleday</i> . | Genus DISCOPHORA, <i>Boisd.</i> |
| E. Halitherses, <i>Doubleday</i> 199 | D. Tullia, <i>Cramer Sp.</i> 211 |
| — Hallirothius, <i>Westwood</i> 199 | — Celinde, <i>Stoll Sp.</i> 211 |
| Genus HERONA, <i>Doubleday</i> . | |
| H. Marathus, <i>Doubleday</i> ... 199 | Genus ENISPE, <i>Doubleday</i> . |
| Genus CASTALIA, <i>Boisduval</i> . | E. Euthymius, <i>Doubleday</i> ... 212 |
| C. Dichroa, <i>Kollar Sp.</i> 199 | — Cyenus, <i>Westwood</i> 212 |
| — Chandra, <i>Moore</i> 200 | — Lepida, <i>Moore</i> 213 |
| Genus DILIPA, <i>Moore</i> . | |
| D. Morgiana, <i>Westw. Sp.</i> ... 201 | Genus CLEROME, <i>Boisduval</i> . |
| Genus APATURA, <i>Fabricius</i> . | C. Arcesilaus, <i>Fabr. Sp.</i> ... 213 |
| A. Ambica, <i>Kollar</i> 201 | — Phaon, <i>Erichson</i> 214 |
| — Parvata, <i>Moore</i> 202 | |
| — Parisatis, <i>Westwood</i> 202 | Genus DRUSILLA, <i>Swainson</i> . |
| — Nakula, <i>Moore</i> 203 | D. Horsfieldi, <i>Swainson</i> ... 214 |
| Genus SYMPHEDEA, <i>Hübner</i> . | |
| S. Thyelia, <i>Fabr. Sp.</i> 204 | Genus THAUMANTIS, <i>Hübner</i> . |
| | T. Odana, <i>Godart Sp.</i> 214 |
| | — Diores, <i>Doubleday</i> 215 |
| | — Ramdeo, <i>Moore</i> 215 |
| | — Nourmahal, <i>Westwood</i> ... 215 |
| | — Lucipor, <i>Westwood</i> 215 |
| | — Noureddin, <i>Westwood</i> ... 215 |
| | — Camadeva, <i>Westwood</i> ... 216 |

SYSTEMATIC LIST

- | | |
|---|---|
| <p>Genus DEBIS, Boisduval.</p> <p>D. Europa, <i>Fabricius Sp. p.</i> 216
 — Neilgherriensis, <i>Guérin Sp.</i> 217
 — Rohria, <i>Fabricius Sp.</i> ... 217
 — Verma, <i>Kollar Sp.</i> 217
 — Bhairava, <i>Moore</i> 217
 — Scanda, <i>Moore</i> 218
 — Nada, <i>Moore</i> 218
 — Arcadia, <i>Cramer Sp.</i>..... 219
 — Mekara, <i>Moore</i> 219
 — Chandica, <i>Moore</i> 219
 — Kansa, <i>Moore</i> 220</p> <p>Genus ZOPHOESSA, Doubleday.</p> <p>Z. Sura, <i>Doubleday</i> 220
 — Yama, <i>Moore</i> 221</p> <p>Genus CYLLO, Boisduval.</p> <p>C. Constantia, <i>Cramer Sp.</i> 221
 — Crishna, <i>Westwood</i> 221</p> <p>Genus MELANITIS, Fabricius.</p> <p>M. Leda, <i>Linnæus Sp.</i> 222
 — Banksia, <i>Fabricius</i> 222
 — Vamana, <i>Moore</i>..... 223
 — Ambasara, <i>Moore</i> 223
 — Bela, <i>Moore</i> 223
 — Suyudana, <i>Moore</i> 224
 — Varaha, <i>Moore</i> 224
 — Gokala, <i>Moore</i> 224
 — Suradeva, <i>Moore</i> 225</p> <p>Genus CÆLITES, Boisduval.</p> <p>C. Epiminthia, <i>Boisduval</i> ... 225</p> <p>Genus ORINOMA, Doubleday.</p> <p>O. Damaris, <i>G. R. Gray Sp.</i> 225</p> <p>Genus NEORINA, Westwood.</p> <p>N. Hilda, <i>Westwood</i> 226</p> <p>Genus RAGADIA, Westwood.</p> <p>R. Makuta, <i>Horsfield Sp.</i>.... 226</p> <p>Genus EREBIA, Dalman.</p> <p>E. Scanda, <i>Kollar</i> 226
 — Annada, <i>Moore</i> 226</p> | <p>Genus LASIOMMATA, Westw.</p> <p>L. Schakra, <i>Kollar</i> p. 227</p> <p>Genus ENOPE, Moore.</p> <p>E. Pulaha, <i>Moore</i> 227
 — Bhadra, <i>Moore</i> 227</p> <p>Genus ZENICA, Westwood.</p> <p>Z. Achanta, <i>Donovan Sp.</i> ... 228</p> <p>Genus SATYRUS, Latreille.</p> <p>S. Padma, <i>Kollar</i> 228
 — Avatarā, <i>Moore</i>..... 229
 — Swaha, <i>Kollar</i> 229
 — Saraswati, <i>Kollar</i> 229</p> <p>Genus ERITES, Boisduval.</p> <p>E. Madura, <i>Horsfield Sp.</i>.... 229</p> <p>Genus MYCALESIS, Hübner.</p> <p>M. Polydecta, <i>Fabricius Sp.</i> 230
 — Otreā, <i>Cramer Sp.</i> 230
 — Francisca, <i>Cramer Sp.</i> ... 230
 — Visala, <i>Moore</i> 230
 — Sanatana, <i>Moore</i> 231
 — Mamerta, <i>Cramer Sp.</i> ... 231
 — Drusia, <i>Cramer Sp.</i> 231
 — Malsara, <i>Moore</i> 231
 — Gotama, <i>Moore</i> 232
 — Patnia, <i>Moore</i> 232
 — Anapita, <i>Moore</i> 232
 — Heri, <i>Moore</i> 233
 — Safitza, <i>Hewitson</i> 233
 — Samba, <i>Moore</i> 233
 — Mandata, <i>Moore</i> 233
 — Hesione, <i>Cramer Sp.</i>.... 234
 — Runeka, <i>Moore</i> 234
 — Janardana, <i>Moore</i> 234</p> <p>Genus THEOPE, Moore.</p> <p>T. Himachala, <i>Moore</i> 234</p> <p>Genus YPHTHIMA, Hübner.</p> <p>Y. Lysandra, <i>Cramer Sp.</i> ... 235
 — Philomela, <i>Hübner</i> 235
 — Pandocus (<i>Boisd.</i>), <i>Moore</i> 235
 — Hyagriva, <i>Moore</i> 236</p> |
|---|---|

OF THE GENERA AND SPECIES.

- Y. Sakra, *Moore* p. 236
 — Narasingha, *Moore* 236

Genus ELYMNIAS, *Hübner*.

- E. Lais, *Fabr. Sp.* 237
 — undularis, *Fabr. Sp.* 237
 — Dusara, *Horsf. Sp.* 238
 — Vasudeva, *Moore* 238
 — Patna, *Westwood Sp.* 238
 — leucocyma, *Godart Sp.* 238
 — Kamara, *Moore* 239

Genus HYPANIS, *Boisduval*.

- H. Ilithyia, *Drury Sp.* 239

Genus LIBYTHEA, *Fabricius*.

- L. Myrrha, *Godart* 240
 — Lepita, *Moore* 240

Stirps V., with

ANOPLURIFORM Larvæ.

Genus ZEMEROS, *Boisduval*.

- Z. Flegyas, *Cramer Sp.* 242

Genus TAXILA, *Doubleday*.

- T. Durga, *Kollar Sp.* 242
 — Egeon, *Boisd. Sp.* 242
 — Fatna, *Boisd. Sp.* 243
 — Erato, *Boisd. Sp.* 243
 — Echerius, *Stoll Sp.* 243
 — Neophron, *Boisd. Sp.* 243
 — Drupadi, *Horsf. Sp.* 244
 — Fylla, *Boisd. Sp.* 244

Genus GONILOBA, *Westwood*.

- G. Chromus, *Cramer Sp.* ... 244
 — Sena, *Moore* 245
 — Derma, *Moore* 245
 — Badra, *Moore* 245
 — Japetus, *Cramer Sp.* 245
 — Gana, *Moore* 245
 — Ravi, *Moore* 246
 — Tapana, *Moore* 246
 — Menaka, *Moore* 246
 — Gopala, *Moore* 246
 — Madhava, *Moore* 246

- G. Pralaya, *Moore* p. 246
 — Bhagava, *Moore* 246
 — Sambara, *Moore* 246

Genus ISMENE, *Swainson*.

- I. Cœdipoda, *Swainson* 247
 — Jaina, *Moore* 247
 — Harisa, *Moore* 247
 — Amara, *Moore* 247
 — Vasutana, *Moore* 247
 — Benjami, *Guérin* 248
 — Jayadeva, *Moore* 248
 — Sinta, *Moore* 248
 — Dasa, *Moore* 248
 — Ladon, *Cramer Sp.* 248
 — Ambasa, *Moore* 248
 — Chaon, *Boisd.* 248
 — Batara, *Moore* 249

Genus PYRGUS, *Hübner*.

- P. Agama, *Moore* 249
 — Superna, *Moore* 249
 — Danna, *Moore* 249
 — Purendra, *Moore* 250

Genus NISONIADES, *Hübner*.

- N. Salsala, *Moore* 250
 — Daha, *Moore* 250
 — Diocles, *Boisd.* 250
 — Amerta, *Moore* 250

Genus PAMPHILA, *Fabricius*.

- P. Vedanga, *Moore* 250
 — Naranata, *Moore* 251
 — Augias, *Linn. Sp.* 251
 — Sumbadra, *Moore* 251

Genus ACHLYODES, *Hübner*.

- A. Chandrasa, *Moore* 251
 — Sura, *Moore* 251
 — Vasava, *Moore* 252

Genus HESPERIA, *Fabricius*.

- H. Folus, *Cramer Sp.* 252
 — Fatih, *Kollar* 252
 — Pulomaya, *Moore* 252
 — leucocerca, *Kollar* 252
 — Putra, *Moore* 253

SYSTEMATIC LIST

- H. Asmara, *Moore* *p.* 253
- Alyssos, *Boisd.* 253
- Praba, *Moore* 253
- Indrani, *Moore* 253
- Dan, *Fabricius* 253
- Sema, *Moore* 253
- Thrax, *Linn. Sp.* 254
- Irava, *Moore* 254
- Pandia, *Moore* 254
- Esa, *Moore* 254
- Druna, *Moore* 254
- Aria, *Moore* 254
- Chaya, *Moore* 255
- Agna, *Moore* 255
- Mangala, *Moore* 255
- Cinnara, *Moore* 255
- Kumara, *Moore* 255
- Sutapa, *Moore* 255
- Divodasa, *Moore* 255

- Genus NYCTALEMON, *Dalman*.
N. Patroclus, *Linn. Sp.* ... 256

Tribe II. SPHINGES.

Stirps II.

Larvæ ELONGATÆ.

- Genus SESIA, *Fabricius*.
S. Hylas, *Linn. Sp.* 260

Genus SATASPES, *Moore*.

- S. infernalis, *Westwood Sp.* 261

Genus MACROGLOSSA, *Ochs.*

- M. Stellatarum, *Linn. Sp.* ... 261
- Corythus, *Boisd.* 262
- Gilia, *Boisd.* 262
- gyrans, *Boisd.* 262
- Sitiene, *Boisd.* 262
- Passalus, *Drury Sp.* 262
- divergens, *Walker* 263
- Nycteris, *Kollar* 263
- Faro, *Cramer Sp.* 263

Genus LOPHURA, *Boisduval*.

- L. Hyas, *Boisduval* 263

Stirps III.

Larvæ ACROCEPHALÆ.

- Genus SMERINTHUS, *Latreille*.
S. Dryas, *Boisduval* *p.* 264

- dentatus, *Cramer Sp.* ... 264

Stirps IV.

Larvae AMBLOCEPHALÆ.

- Genus LEUCOPHLÆBIA, *Westw.*
L. lineata, *Westwood* 265

Genus BASIANA, *Walker*.

- B. cervina, *Walker* 265

Genus AMBULYX, *Boisduval*.

- A. substrigilis, *Westwood* ... 266

Genus CALYMNIA, *Boisduval*.

- C. Panopus, *Cramer Sp.* ... 266

Genus ACHERONTIA, *Ochs.*

- A. Styx, *Westwood* 266
- Satanás, *Boisd.* 267

Genus SPHINX, *Linneaus*.

- S. Convolvuli, *Linn.* 267

Genus MACROSILA, *Boisduval*.

- M. nyctiphanes, *Boisd.* 268
- discistriga, *Walker* 268

Genus ZONILIA, *Boisduval*.

- Z. Morpheus, *Cramer Sp.* ... 269

Stirps V.

Larvæ OPHTHALMICÆ.

Genus PANACRA, *Walker*.

- P. Automedon, *Boisd. Sp.* ... 269
- scapularis, *Horsf. Sp.* ... 269
- Vigil, *Guérin Sp.* 270
- Busiris, *Boisd. Sp.* 270

OF THE GENERA AND SPECIES.

Genus PHILAMPELUS, *Harris.*

- P. Anceus, *Cramer Sp.* ... p. 270
- sericeus, *Walker* 271
- Naga, *Moore* 271

Genus DARAPSA, *Walker.*

- D. Hypothoüs, *Cramer Sp.* 271

Genus DAPHNIS, *Hübner.*

- D. Nerii, *Linn. Sp.* 272

Genus PERGESA, *Walker.*

- P. Acteus, *Cramer Sp.* 272
- Castor, *Boisd. Sp.* 273

Genus ELIBIA, *Walker.*

- E. Dolichus, *Westwood Sp.* 273

Genus DEILEPHILA, *Ochs.*

- D. Lathyrus, *Boisd.* p. 274
- Livornica, *Esper Sp.* 274

Genus CHÆROCAMPA, *Dup.*

- C. Celerio, *Linn. Sp.* 274
- Alecto, *Linn. Sp.* 275
- suffusa, *Walker* 275
- pallicosta, *Boisd.* 276
- Thyelia, *Linn. Sp.* 276
- lineosa, *Walker* 276
- Nessus, *Drury Sp.* 276
- Clotho, *Drury Sp.* 277
- Lucasi, *Boisd.* 277
- Lyceutes, *Cramer Sp.* 277
- Oldenlandiæ, *Fabr. Sp.* 278
- bisecta, *Horsf.* 278

CATALOGUE
OF
THE LEPIDOPTEROUS INSECTS
IN THE
Museum of the East-India Company.

Class INSECTA.

Div. I. INSECTA HAUSTELLATA.

INSECTA HAUSTELLATA (*Clairville*). *Stephens, Ill. Brit. Ent. vol. I. p. 1 (1828).*

Order LEPIDOPTERA.

LEPIDOPTERA, *Linnæus, Syst. Nat. (1748).*

GLOSSATA, *Fabricius, Syst. Entom. (1775).*

LEPIDOPTERA, *Clairville.*

LÉPIDOPTÈRES, *Duméril, Consid. Gén. des Ins. p. 134 (1823).
Boisduval, Spéc. Gén. Lép. I. p. 162 (1836).*

Tribe I. PAPILIONES.

LEPIDOPTERA DIURNA, *Latreille, Gen. IV. p. 186 (1809). Stephens.
G. R. Gray.*

RHOPALOCÈRES, *Duméril, Consid. Gén. des Ins. p. 222 (1823). Bois-
duval, Spéc. Gén. Lép. I. p. 162 (1836).*

PAPILIONIDÆ, *Horsfield, Cat. Lep. Mus. E.I.C. pp. 20, 57 (1828).*

PAPILIONIDES, *Boisduval, Hist. Gén. des Lép. Amér. Sep. p. 2
(1829).*

RHOPALOCERA, *Stainton, Ins. Brit. Lepid. Tineina, p. 2 (1854); et
Manual Brit. Butt. p. 10 (1856). Walker, Cat. Lep. Brit. Mus.
pt. I. Lep. Het. p. 1 (1854).*

SUCCINCTI, *Boisduval, Spéc. Gén. Lép. I. pp. 162, 171 (1836).*

SUCCINCTÆ, *Boisduval, Index Méth. p. 1 (1840).*

Stirps I., with VERMIFORM Larvæ.

VERMIFORM, *Horsfield*, *Cat. Lep. Mus. E.I.C.* pp. 21, 38, 64 (1828).

LYCÆNIDÆ, *Leach*, *Edinb. Encyclop.* IX. p. 129 (1815). *Stephens*.

Doubleday. *Westwood*. *Stainton (Manual)*, pp. 10, 40.

POLYOMMATIDÆ, *Swainson*, *Phil. Mag.* I. p. 187 (1827).

POLYOMMATES, *Boisduval*, *Hist. Gén. des Lép. Amér. Sep.* p. 2 (1829).

LYCÆNIDES, *Boisduval*, *Icon.* I. (1832); id. *Sp. Gén. Lép.* I. p. 164 (1836); id. *Index Méth.* p. 8 (1840).

PLEBEI RURALES, *Linnæus*.

POLYOPHTHALMI, *Aldrovandus*.

Genus MILETUS, *Hübner*.

MILETUS, *Hübner*, *Verz. bek. Schmett.* (1816). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 502.

SYMETHA, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 59, t. 2, f. 2 (1828).

GERYDUS, *Boisduval*, *Sp. Gén. Lép.* p. 6 (1836).

1. *MILETUS SYMETHUS*, *Cramer Sp.*

♀ Papilio Symethus, *Cramer*, *Pap. Exot.* II. t. 149, f. B. C. (1779). *Stoll*, *Suppl. à Cramer's Pap.* I. t. 37, f. 3, 3c. *Fabricius*, *Mant. Ins.* II. p. 69.

Miletus Symethus, *Hübner*, *Verz. bek. Schmett.* *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 503, No. 1.

Polyommatus Symethus, *Godart*, *Enc. Méth.* IX. p. 675.

♀ Gerydus Symethus, *Boisduval*, *Sp. Gén. Lép.* p. 6, t. 23, f. 2. *Doubleday*, *List Lep. Brit. Mus. pt. II.* p. 57.

♀ Symetha Pandu, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 2, f. 2, 2a (1828).

? Symæthus Leos, *Guérin*, *Voy. Coquille Ins.* t. 18, f. 8.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The characters of this genus are peculiar, and its situation in a Natural Series remains for future investigation. Its chief peculiarity consists in the character of its legs, of which Mr. Westwood gives the following description: "Legs rather short, slender, scaly, compressed. The tarsi in all the feet, with the basal joint, remarkably

elongated, widened, and quite compressed ; the tarsus in the fore-legs of the male being exarticulate, and as long as the femur and tibia united ; second, third, and fourth joints in the four hind-legs very short, terminal joint small, subovate, notched at its obliquely-truncated extremity, with extremely minute unguis."

2. *MILETUS BOISDUVALI*, Moore (Plate 1a, fig. 1).

M. Boisduvali, n. sp.—UPPER-SIDE brown ; fore-wing with a whitish fascia running from near the middle of the anterior margin to near the anal angle ; hind-wings slightly angulated. UNDER-SIDE pale brown ; fore-wing, with fascia, as on upper side, intersected with veinlets, and having three spots in the discoidal cell, one on each of the three anterior subcostal veinlets ; also an undulated spot near the anterior angle, and a small blackish dot between each veinlet near the exterior margin ; hind-wing also marked with undulated spots and stripes, and with a dot between each veinlet. Form of wings as in ♀ *Symætha* ; expanse of wings $1\frac{4}{5}$ in.

a. b. ♀. Java. From Dr. Horsfield's Collection.

3. *MILETUS HORSFIELDI*, Moore (Plate 1a, fig. 2).

M. Horsfieldi, n. sp.—UPPER-SIDE brown, fore-wing of the male with an oval longitudinal whitish spot in the centre ; female wholly brown, with but a faint indication of lighter colour in the centre of the fore-wing. UNDER-SIDE of both sexes creamy-white, and densely covered with undulating brown strigæ, both wings also having near the exterior margin a darker brown dot between each veinlet. Hind-wings rounded, and, in the male, slightly angulated ; in the female, these are acute. Expanse of wings in the male $1\frac{7}{8}$ in., female $1\frac{5}{8}$ in.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

Genus PITHECOPS, Horsfield.

PITHECOPS, Horsfield, Cat. Lep. Mus. E.I.C. p. 66 (1828).

LYCENA, p. Westwood, Diurnal Lep.

4. *PITHECOPS HYLAX*, Fabricius Sp.

Hesperia Hylax, Fabricius, Mant. Ins. II. p. 77 (1787) ; Ent. Syst. III. pt. I. p. 304.

Polyommatus Hylax, Godart, Enc. Méth. H. N. IX. p. 701. Donovan, Ins. of India, t. 46, f. 2.

δ ♀ Pithecopus Hylax, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 66, t. 1, f. 2, 2a, 2b (1828). *Doubleday, List Lep. Brit. Mus. pt. II.* p. 57.

Lycæna Hylax, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 496, No. 198, t. 76, f. 8.

a. b. δ . Java. From Dr. Horsfield's Collection.

c. d. ♀. India. From the Asiatic Society of Bengal.

"In many essential points, the form of the palpi and antennæ, the peculiarities of the feet, &c., this insect agrees with the character of the genus *Polyommatus*, but its habit and aspect are very peculiar: this is owing to the great length and lateral expansion of the wings, to their comparative narrowness, and to their being regularly elliptical and rounded in the anal region."—(Horsfield, Catal.)

"This insect has also naked eyes, scaly palpi, and spurless fore tibiae, but the wings have the veins arranged in the ordinary manner, and the unguis and their appendages are as small and inconspicuous as in the genus *Lycæna*."—(Westwood, Diurn. Lep.)

Genus POLYOMMATUS, *Latreille, Pr. S. D.*

POLYOMMATUS, Latreille, Règ. An. III. p. 553 (1817).

POLYOMMATUS, Stephens. Curtis. Swainson.

LYCÆNA, p. Doubleday. Westwood (Diurn. Lep.).

5. *POLYOMMATUS AKASA*, Horsfield.

♀ Polyommatus Akasa, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 67, t. 1, f. 1, 1a (1828).

Lycæna Akasa, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 491, No. 55.

a. b. ♀. Java. From Dr. Horsfield's Collection.

"This species is not abundant in the island of Java, two specimens only being obtained."—(Horsfield.)

6. *POLYOMMATUS PUSPA*, Horsfield.

δ ♀ Polyommatus Puspa, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 67 (1828).

Lycæna Puspa, Doubleday, List Lep. Brit. Mus. pt. II. p. 44. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 491, No. 56.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. N. India. Presented by Colonel Buckley.

"Common in Java."—(Horsfield.)

7. *POLYOMMATUS LAIUS*, Cramer Sp.

♀ Papilio Laius, Cramer, *Pap. Exot.* IV. t. 319, f. D. E. (1782).

Lycæna Laius, *Doubleday, List Lep. Brit. Mus.* pt. II. p. 44. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 491, No. 63.

Hesperia Caius, *Fabricius, Ent. Syst.* III. pt. I. p. 296 (1793).

Polyommatus Caius, *Godart, Enc. Méth.* IX. p. 701.

a. b. ♂ ♀. India. From the Asiatic Society of Bengal.

Genus LYCÆNA, Fabricius.

LYCÆNA, *Fabricius (Illiger's Mag.* VI. 1807). *Westwood, Diurn. Lep.*

POLYOMMATUS, *p. Godart.*

8. *LYCÆNA PLUTO*, Fabricius Sp.

Hesperia Pluto, *Fabricius, Ent. Syst.* III. pt. I. p. 288 (1793).

Lycæna Pluto, *Donovan, Ins. of India*, t. 45, f. 2. *Hombron et Jacquinot, Voy. Pôle Sud, Lép.* t. 3, f. 9, 10. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 42.

Lycæna Nila, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 78 (1828). *E. Doubleday, List Lep. Brit. Mus.* pt. II. p. 43 (nec. T. Nila, *Kollar*).

a. b. ♂. N. India. From the Asiatic Society of Bengal.

c. ♂. Ceylon. Presented by the Trustees of the British Museum.

d. ♂. Java. (*L. Nila, Horsf.*) From Dr. Horsfield's Collection.

9. *LYCÆNA AMYNTAS*, Fabricius Sp.

Papilio Amyntas, *Fabricius*, *Mant. Ins. II.* p. 70 (1787).
Wien. Verz. p. 185.

Lycæna Amyntas, *Ochsenheimer, Schmett. von Europa*, IV.
 p. 26. *E. Doubleday, List Lep. Brit. Mus.* pt. II.
 p. 43. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 43.

Polyommatus Amyntas, *Godart, Enc. Méth.* IX. p. 659.
 a. ♂. China (?). Donor unknown.

10. *LYCÆNA ARATUS** Cramer Sp.

♀ Papilio aratus, *Cramer, Pap.* IV. t. 365, f. A.B. ? (1782).
 Lycæna atratus, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 78.
Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 490, No. 29.

a. ♀. Java. From Dr. Horsfield's Collection.

If new, *L. Kurava*, Moore.

11. *LYCÆNA PARRHASIUS*, Fabricius Sp. (*Plate 1a, fig. 3*).

Hesperia Parrhasius, *Fabricius, Ent. Syst.* III. pt. I.
 p. 289 (1793).

Lycæna Parrhasius, *Horsfield, Cat. Lep. Mus. E.I.C.*
 p. 86. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 489, No. 13.

Papilio Parrhasius, *Donovan, Ins. India*, t. 45, f. 5.

Polyommatus Parrhasius, *Godart, Enc. Méth.* IX. p. 657.

a. b. ♂. Java. From Dr. Horsfield's Collection.

12. *LYCÆNA PANDAVA*, Horsfield.

♀ Lycæna Pandava, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 84
 (1829). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 489, No. 14.

a. ♀. Java. From Dr. Horsfield's Collection.

13. *LYCÆNA CNEJUS*, Fabricius Sp.

Hesperia Cnejus, *Fabricius, Ent. Syst.* V. *Suppl.* p. 430
 (1798).

* This species was named *aratus* by Cramer, not *atratus* as quoted by authors.

Lycæna Cnejus, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 83.

Doubleday, *List Lep. Brit. Mus. pt. II.* p. 43.

Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 40.

Polyommatus Cnejus, *Godart*, *Enc. Méth. IX.* p. 657.

a. ♀. Java. From Dr. Horsfield's Collection.

b. c. ♂ ♀. India. From the Asiatic Society of Bengal.

14. *LYCÆNA DAMOETES*, *Fabricius* Sp.

Papilio Damoetes, *Fabricius*, *Mant. Ins. II.* p. 77 (1787).

Donovan, *Ins. New Holl.* t. 31, f. 2.

Lycæna Damoetes, *Horsfield*, *Cat. Lep. Mus. E. I. C.*

p. 81. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 496, No. 193.

Polyommatus Damoetes, *Godart*, *Enc. Méth. IX.* p. 680.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

15. *LYCÆNA KANDARPA*, *Horsfield*.

Lycæna Kandarpa, *Horsfield*, *Cat. Lep. Mus. E. I. C.*

p. 82 (1829). *Doubleday*, *List. Lep. Brit. Mus. pt. II.* p. 43. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 39.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. e. ♂. N. India. From the Asiatic Society of Bengal.

f. ♀. N. India. Presented by Colonel Hearsey.

16. *LYCÆNA BOETICA*, *Linnæus* Sp.

Papilio Boeticus, *Linnæus*, *Syst. Nat. II.* p. 789 (1766).

Hübner, *Eur. Schmett. Pap. t. 74*, f. 373, 375.

Ochsenh. Pap. Eur. t. 1, 2, p. 99; id. *Schmett. v. Eur.* p. 27.

Lycæna Boetica, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 80.

Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 418.

Doubleday, *List Lep. Brit. Mus. pt. II.* p. 42.

Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 25. *Gerhard*, *Lycænidæ*, t. 11, f. 1.

Hesperia Boetica, *Fabricius*, *Ent. Syst.* III. pt. I. p. 280.
Polyommatus Boeticus, *Godart*, *Enc. Méth.* IX. p. 653.
Papilio Coluteæ, *Rossi*, *Faun. Etr.* II. p. 155 (1795).

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. f. g. h. ♂ ♀. N. India. Presented by Colonel Hearsey.

17. *LYCÆNA PAVANA*, *Horsfield*.

Lycæna Pavana, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 77 (1828). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 41 (nec. *P. Pavana*, *Kollar*).

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

18. *LYCÆNA ELPIS*, *Godart Sp.*

Polyommatus Elpis, *Godart*, *Enc. Méth.* IX. p. 654 (1819).

Lycæna Elpis, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 76, t. 1, f. 4. *Doubleday, List Lep. Brit. Mus.* pt. II. p. 42. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 28.

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. N. India. From the Asiatic Society of Bengal.

19. *LYCÆNA CELENO*, *Cramer Sp.*

♂ *Papilio Celeno*, *Cramer*, *Pap. Exot.* I. t. 31, f. C. D. (1779). *Donovan's Drawings in Bibl. Hope, Oxford.*

Lycæna Celeno, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 490, No. 27.

Papilio Celerio, *Fabricius*, *Mant. Ins.* II. p. 66 (1787).

Polyommatus Celerio, *Godart*, *Enc. Méth.* IX. p. 654.

Lycæna Celerio, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 75. *Doubleday, List Lep. Brit. Mus.* pt. II. p. 42.

- a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.
d. ♂. Borneo. Presented by A. Lowe, Esq.

20. LYCÆNA ALEXIS, Stoll Sp.

Lycæna Alexis, *Doubleday, List Lep. Brit. Mus. pt. II.*
p. 42.

♂ *Papilio Alexis*, *Stoll, Suppl. Cramer's Pap.* *p. 167, t. 38,*
f. 3, 3a (1791).

Hesperia Ælianuſ, *Fabricius, Ent. Syst. III. pt. I.* *p. 280*
(1793).

Lycæna Ælianuſ, *Horsfield, Cat. Lep. Mus. E.I.C.* *p. 73,*
t. IV. f. 1, 1a. Westwood, in Doubleday and Hewitson's Diurnal Lep. *p. 490, sp. 38.*

Polyommatus Ælianuſ, *Godart, Enc. Méth. IX.* *p. 654.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. N. India. From the Asiatic Society of Bengal.

g. h. ♂ ♀. Canara. Presented by S. N. Ward, Esq.

"The larva of *L. Alexis* feeds on the *Butea frondosa*, and was observed in the month of February in Java."—(Horsfield.)

The larva and pupa of this species are figured on Plate 1, fig. 1, 1a.

21. LYCÆNA PLINIUS, Fabricius Sp.

Hesperia Plinius, *Fabricius, Ent. Syst. III. pt. I.* *p. 284*
(1793).

Lycæna Plinius, *Horsfield, Cat. Lep. Mus. E.I.C.* *p. 72.*
Westwood, in Doubleday und Hewitson's Diurnal Lep. *p. 489, No. 17.*

Papilio Plinius, *Donovan, Ins. of India*, *t. 45, f. 1.*

Polyommatus Plinius, *Godart, Enc. Méth. IX.* *p. 658.*

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. e. f. ♂ ♀. N. India. Presented by Colonel Hearsey.

22. LYCÆNA THEOPHRASTUS, Fabricius Sp.

Hesperia Theophrastus, *Fabricius, Ent. Syst. III. pt. I.*
p. 281 (1793).

Lycæna Theophrastus, *Horsfield, Cat. Lep. Mus. E.I.C.*
p. 73. Doubleday, List Lep. Brit. Mus. pt. II. *p. 41.*

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 489, No. 10.

Polyommatus Theophrastus, Godart, Enc. Méth. IX. p. 658.

a. b. India. From the Asiatic Society of Bengal.

c. Canara. Presented by S. N. Ward, Esq.

d. e. f. g. N. India. Presented by Colonel Hearsey.

23. *LYCÆNA ROSIMON, Fabricius Sp.*

Papilio Rosimon, Fabricius, Mant. Ins. II. p. 71 (1787).

Lycæna Rosimon, Horsfield, Cat. Lep. Mus. E.I.C. p. 71. *Doubleday, List Lep. Brit. Mus. pt. II.* p. 41.

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 489, No. 8.

Polyommatus Rosimon, Godart, Enc. Méth. IX. p. 658.

Papilio Corydon, Cramer, Pap. t. 340, f. C.D.E. 1782 (*nec. Fabr.*).

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

f. g. Silhet. Presented by Colonel F. Buckley.

24. *LYCÆNA MALAYA, Horsfield.*

Lycæna Malaya, Horsfield, Cat. Lep. Mus. E.I.C. p. 70 (1828). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 489, No. 15.

a. b. Java. From Dr. Horsfield's Collection.

"This species is comparatively scarce in Java, having obtained only two specimens. In its habit it greatly resembles the individuals of the genus *Polyommatus*."—(Horsfield.)

25. *LYCÆNA ROXUS, Godart Sp.*

Polyommatus Roxus, Godart, Enc. Méth. IX. p. 659 (1819).

Lycæna Roxus, Horsfield, Cat. Lep. Mus. E.I.C. p. 70, t. 2, f. 4, 4a. *Doubleday, List Lep. Brit. Mus. pt. II.* p. 41. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 489, No. 7.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

"Hitherto this species has only been found in Java; it occurs in considerable numbers in the skirts of large forests; but from the great delicacy of the wings, it is not easily obtained in a perfect state."—(Horsfield.)

26. LYCAENA PSEUDOROXUS, Doubleday.

Lycaena pseudoroxus, Doubleday, *List Lep. Brit. Mus.* pt. II. p. 41 (1847).

a. Silhet. Presented by the Trustees of the British Museum.

27. LYCAENA NYSEUS, Guérin Sp.

Polyommatus Nyseus, Guérin-Méneville, in Delessert's *Souv. Voy. Inde*, pt. II. p. 78, t. 22, f. 1, 1a (1843).

Lycaena Nyseus, Doubleday, *List Lep. Brit. Mus.* pt. II. p. 40. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 489, No. 6.

a. b. c. Canara. Presented by S. N. Ward, Esq., E.I.C.C.S.

d. e. Madras. From Captain J. M. Jones's Collection.

Genus CHRYSOPHANUS, Hübner.

CHRYSOPHANUS, Hübner, *Verz. bek. Schmett.* p. 72 (1816). Westwood. Stephens.

POLYOMMATUS, Boisduval. H. Schaeffer. Doubleday.

LYCAENA, Stephens (Ill.). Curtis (Guide).

28. CHRYSOPHANUS PHLÆAS, Linnaeus Sp.

Papilio Phlæas, Linnaeus, *Syst. Nat.* II. p. 793 (1766).

Chrysophanus Phlæas, Hübner, *Eur. Schmett. Pap.* f. 362, 363. Stephens, *Cat. Brit. Lep. Brit. Mus.* p. 17. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 498, No. 4.

Polyommatus Phlæas, Godart, *Enc. Méth.* IX. p. 670. Doubleday, *List Lep. Brit. Mus.* pt. II. p. 53. Kollar, in Hugel's *Kaschmir*, IV. pt. II. p. 417.

Lycaena Phlæas, Stephens, Ill. *Brit. Lep. H. I.* p. 79.

♀ *Papilio Virgaureæ*, Scopoli, *Ent. Carn.* p. 181.

Var. ♀ *Papilio Timæus*, Cramer, *Pap.* t. 146, f. E.F.

Var. *Papilio turcicus*, Gerhard, *Lycæn.* t. 5, f. 5.

- a. b.* ♂ ♀. Bootan. From Pemberton's Collection.
c. d. ♂ ♀. N. India. Presented by Colonel Hearsey.
e. ♀. Darjeeling. Collected by the Messrs. Schlag-intweit.

29. *CHRYSORHANUS PAVANA*, Kollar Sp.

- ♂ Polyommatus Pavana, Kollar, in Hügel's Kaschmir, IV.
 pt. II. p. 416, t. 5, f. 56 (1844), (nec. L. Pavana, Horsf.).
 Thecla Panava, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 487, No. 143 (1852).
a. b. ♂. N. India. Presented by Colonel Buckley.
c. d. e. f. ♂ ♀. N. India. Presented by Colonel Hearsey.

Genus ILERDA, E. Doubleday.

- ILERDA, E. Doubleday, List Lep. Brit. Mus. pt. II. p. 25 (1847).
 THECLA, p. Boisduval. Westwood, in Doubleday and Hewitson's Diurnal Lep.

30. *ILERDA SENA*, Kollar Sp.

- ♂ Polyommatus Sena, Kollar, in Hügel's Kaschmir, p. 415, t. 5, f. 3, 4 (1844).
 Thecla Sena, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 487, No. 142.
 ♀ Thecla Cadma, Boisduval, MS. (1847).
 Ilerda Cadma, Doubleday, List Lep. Brit. Mus. pt. II. p. 25.
a. b. ♂. Darjeeling. Reserved from the Indian Collection, Exposition Universelle at Paris, 1855.
b. c. ♀. N. India. Presented by Colonel Buckley.

31. *ILERDA TAMU*, Kollar Sp.

- Polyommatus Tamu, Kollar, in Hügel's Kaschmir, IV.
 pt. II. p. 417, t. 5, f. 7, 8 (1844).
 Thecla Tamu, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 487, No. 144.
a. b. ♂. Bootan. From Pemberton's Collection.

Remark.—Upper-side of wings with glittering blue patches.

32. *ILERDA ANDROCLES*, Boisduval Sp.

Thecla Androcles, *Boisduval*, *MS.* (1847). *Westwood*,
in Doubleday and Hewitson's Diurnal Lep. p. 487,
No. 141, tab. 75, f. 2.

Ilerda Androcles, *E. Doubleday, List Lep. Brit. Mus.*
pt. II. p. 25.

- a. ♂. Silhet. Presented by the Trustees of the British Museum.
- b. c. ♂. Darjeeling. Presented by Lieutenant Hugo James.
- d. e. ♂. Darjeeling. Collected by Messrs. Schlagintweit.

Remark.—Upper-side of wings with glittering green patches.

33. *ILERDA BRAHMA*, Moore (*Plate 1a, Fig. 4*).

Ilerda Brahma, *n. sp.*.—Differs from the two preceding species in having the patches of the wings glittering coppery-gold colour, and a broader and longer exterior red band to the hind-wings.

- a. ♂. Darjeeling. Collected by the Messrs. Schlagintweit.

34. *ILERDA EPICLES*, Godart Sp.

Polyommatus Epicles, *Godart*, *Enc. Méth.* IX. p. 646 (1819).

Thecla Epicles, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 92, t. 1, f. 3. *Westwood*, *in Doubleday and Hewitson's Diurnal Lep.* p. 487, No. 140.

Ilerda Epicles, *Doubleday, List Lep. Brit. Mus.* pt. II. p. 25.

♀ *Heliophorus Belenus*, *Hübner*, *Zutr.* f. 785, 786 (1832).

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
- e. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

“This species is not uncommon in the acclivities near the confines of the extensive medial plain of Java.”—(Horsfield.)

"The *Ilerdæ* are beautiful species, with naked eyes, setose palpi, antennæ with long joints ringed with white, and a very well-marked club, and with three branches to the postcostal vein of the fore-wings."—(Westwood, *Diurnal Lep.* p. 482.)

Genus THECLA, *Fabricius*.

THECLA, *Fabricius* (*in Illiger's Magaz.* VI.) (1807).

POLYOMMATUS, *p.* *Godart*.

35. THECLA ONYX, *Boisduval*.

Thecla Onyx, *Boisduval, MS.* (1847).

Myrina Onyx, *E. Doubleday, List Lep. Brit. Mus.* pt. II. p. 22.

a. Moulmein. Presented by the Trustees of the British Museum.

Thecla Onyx.—UPPER-SIDE has the *fore-wings* dark brown, with a patch of white in the centre; the posterior base, from the subcostal veinlet to the middle of the disc, cyaneous. *Hind-wings* light brown; centre, from base to near exterior margin, cyaneous, paling to light bluish-brown along the abdominal margin. Tails three, two inner longish. UNDER-SIDE pale ferruginous brown; a white fascia crossing the anterior and posterior wings towards anal angle, and ending upwards to abdominal margin in a greenish-metallic line, the latter being bounded inwardly by a narrow blackish line; a blackish spot and line at the anal angle, and another spot some distance off on the exterior margin.

Genus DIPSAS, *Doubleday*.

DIPSAS, *Doubleday, List Lep. Brit. Mus.* pt. II. p. 25 (1847).

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 479 (1852).

APHNEUS, *p. Doubleday, List Lep. Brit. Mus.* pt. II. p. 25 (1847).

SITHON, *Hübner, Verz. bek. Schmett.* p. 77 (1816).

36. DIPSAS SYLA, *Kollar Sp.*

♂ *Thecla Syla, Kollar, in Hügel's Kaschmir*, IV. pt. II. p. 414, t. 4, f. 7, 8 (1844).

Dipsas Syla, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 1.

♂ *Thecla Pholus, Boisduval, MS.* (1847).

- Dipsas Pholus, *Doubleday, List Lep. Brit. Mus.* pt. II.
p. 25.
- ♀ Amblypodia Euphranor, *Doubleday, List Lep. Brit. Mus.*
pt. II. p. 25 (1847).
- a. ♂. Darjeeling. From Pearson's Collection.
- b. c. d. ♂. N. India. Presented by Colonel Buckley.
- d. e. ♀. Darjeeling. Reserved from the Indian Col-
lection, Exposition Universelle at Paris.

"This species, which is the type of the genus, has a general resemblance to the types of *Amblypodia*, the male being golden-green on the upper surface of the wings, with a black margin (thus resembling *Amblyp. Eumolphus*), whilst the female has the disc of the fore-wings blue, with a white spot beyond the middle, and a broad dark brown margin. From the typical *Amblypodiae* they are at once distinguished by the hairy eyes and palpi, and the want of an upper disco-cellular vein in the fore-wings."—(Westwood.)

37. DIPSAS ATAXUS, Boisduval Sp.

- Thecla Ataxus, *Boisduval, MS.* (1847).
- ♂ Dipsas Ataxus, *Doubleday, List Lep. Brit. Mus.* pt. II.
p. 25. *Westwood, in Doubleday and Hewitson's
Diurnal Lep.* p. 480, t. 74, f. 7.
- a. ♂. N. India. Presented by Colonel Buckley.
- b. ♂. Darjeeling. From Messrs. Schlagintweit's
Collection.

38. DIPSAS XENOPHON, Fabricius Sp.

- Hesperia Xenophon, *Fabricius, Ent. Syst.* III. pt. I.
p. 272 (1793).
- Dipsas Xenophon, *Westwood, in Doubleday and Hewit-
son's Diurnal Lep.* p. 480, No. 3.
- Polyommatus Xenophon, *Godart, Enc. Méth.* IX. p. 640.
- Papilio Xenophon, *Donovan, Ins. of India,* t. 41, f. 3.
- Thecla Xenophon, *Horsfield, Cat. Lep. Mus. E. I. C.*
p. 94, t. IV. f. 2, 2a.

- a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Col-
lection.

"The larva of this species feeds on *Schmiedelia racemosa*, and was found in Java in considerable numbers."—(Horsfield.)

The larva and pupa of *D. Xenophon* are figured on Plate 1, fig. 3, 3a.

39. DIPSAS MELAMPUS, Cramer Sp.

♂ *Papilio Melampus*, *Cramer, Pap. Exot. IV. t. 362, f. G.H.* (1782).

Aphnaeus Melampus, *Doubleday, List Lep. Brit. Mus. pt. II. p. 25.*

Hesperia Jarbas, *Fabricius, Mant. Ins. II. p. 68* (1787).

Dipsas Jarbas, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 5.*

Polyommatus Jarbas, *Godart, Enc. Méth. IX. p. 646.*

Papilio Jarbas, *Donovan, Ins. of India, t. 40, f. 3.*

Thecla Jarbas, *Horsfield, Cat. Lep. Mus. E.I.C. p. 93.*

♂ *Thecla Sorya, Kollar, in Hügel's Kaschmir, IV. pt. II. p. 414, t. 5, f. 1, 2* (1844).

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

f. g. h. ♂. N. India. Presented by Colonel Buckley.

The larva and pupa of *D. Melampus* are figured on Plate 1, fig. 2, 2a.

40. DIPSAS EPIJARBAS, Boisduval Sp.

Thecla Epijarbas, Boisduval, MS.

Aphnaeus Epijarbas, *Doubleday, List Lep. Brit. Mus. pt. II. p. 26.*

a. ♂. Canara. Presented by S. N. Ward, Esq., E.I.C.S.

b. ♀. N. India. Presented by the Trustees of the British Museum.

c. ♂. Darjeeling. From the Indian Collection, Exposition Universelle at Paris, 1855.

This is larger than *D. Jarbas*, the male having on the **UPPER-SIDE** of the fore-wings the dark brown border occupying the whole space between the anterior margin and the median vein (in the latter of which it is nearly black), and then branching off to exterior margin, and ending in a point at the middle of the hind margin. The **UNDER-SIDE** is of a duller tint, and has two whitish undulating lines crossing

the fore and hind-wings, as also two short lines from the median to subcostal veinlet. The outer black spot is entirely encircled with yellow, and above the anal spot is a well-defined lunar-shaped streak of metallic green. The female is somewhat larger, with rounder wings, and is of a fulvous-brown, and may be distinguished by the similarity of the markings of the under-side. Expanse of wings in the male $1\frac{7}{12}$ in., in the female $1\frac{9}{10}$ in.

41. *DIPSAS CHRYSOMALLUS*, Hübner Sp.

♂ *Zesius chrysomallus*, Hübner, *Zutrage*, f. 301, 302 (1823).

Chrysophanus chrysomallus, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 499, No. 29.

? ♀ Myrina Setho, *E. Doubleday, List Lep. Brit. Mus. pt. II.* p. 22.

a. ♂. S. India. Donor not known.

b. ♀. Canara. Presented by S. N. Ward, Esq.

c. d. ♀. Locality and donor unknown.

D. chrysomallus.—UPPER-SIDE of male pale cupreous, outer-margin brownish; at anal angle three blackish spots. Tails three. UNDER-SIDE pinkish creamy-white; a band composed of roundish pinkish spots disposed irregularly across both wings; fore-wing with five, and hind-wing with eight pinkish spots towards the base; anal angle with three spots, the two outer black, banded above with red, and the centre one bluish; near outer-margin is a line of indistinct marks; extreme outer-margin and tails pinkish. Female larger, wings more square, brown, tinged with bluish at base; spots on under-side, as in the male, but of a brighter colour.

42. *DIPSAS ISOCRATES*, Fabricius Sp.

Hesperia Isocrates, Fabricius, *Ent. Syst. III. pt. I.* p. 266 (1793).

Dipsas Isocrates, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 480, No. 4.

Polyommatus Isocrates, Godart, *Enc. Méth. IX.* p. 633.

Thecla Isocrates, Westwood, *Trans. Linn. Soc. II.* p. 4, t. 1, and *Donovan's Ins. India*, 2nd edit. *Downes, Calcutta Journ. N. H. II.* p. 408.

Aphnaeus Isocrates, *Doubleday, List Lep. Brit. Mus. pt. II.* p. 26.

Hesperia Pann, *Fabricius, Ent. Syst. III. pt. I. p. 276.*
 Papilio Pann, *Donovan, Ins. India, t. 38, f. 1 (nec. Drury).*

- a. b. c. d. ♂ ♀. Silhet. Presented by Colonel Buckley.
 e. f. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

The following notice of the transformations of this interesting species was read before the Entomological Society of London, February 2, 1835, by J. O. Westwood, Esq., having been traced by Mr. Charles King, of Madras.

"The larvæ of this butterfly reside in the interior of the pomegranate; seven or eight, at least, having been reared in the interior of a small specimen of this fruit. Of the mode in which the eggs are deposited by the female in the interior of the pomegranate, no information has been received; it is, however, probable that this is effected whilst the fruit is in its very young state. The caterpillars feed upon the seeds and inner part of the fruit, which is thus weakened, and rendered unable to support its own weight, and consequently liable to have its stem broken, and to fall to the ground with the first wind. This, however, would be destruction to the inclosed insects; since, in all probability, they would find it impossible to make their escape were the fruit to be suffered to lie rotting upon the ground. To obviate this evil, the caterpillars, when full fed, have the instinct to eat a hole, about a quarter of an inch in diameter, through the hard shell of the fruit, whilst it remains upon the tree; through this hole they then creep to the stem of the fruit, and spin a white web, which they attach to the basal part of the fruit as well as to the stem, for about the distance of an inch along the latter. This web is sufficiently strong to support the pomegranate from falling after the wind has broken the stem near to the fruit, as is the case in the specimen of the fruit in hand."

From the circumstance of this specimen having as many holes in it as there were caterpillars inhabiting it, it is most probable that the web thus spun is a joint production of the whole.*

* It is curious, as evidencing the instinctive impulses under which each of the inclosed larvæ must have acted, that, instead of availing themselves of the first aperture made in the fruit, each caterpillar should be at the trouble of making a hole for itself, a circumstance which renders it the more probable that all joined in spinning the web.

But it will be at once asked, What necessity could there be for the caterpillars to secure the fruit from falling after each has bored a hole and thus made its escape? This question is answered by the curious circumstance that, after so securing the fruit, the caterpillars return again into the pomegranate, in the hollow interior of which they undergo their transformation to the chrysalis state.

Here, too, we may notice another interesting fact, namely, that the insect has the precautionary instinct, which acts as a second inducement, to make the aperture in the fruit in that stage of its existence in which it is furnished with organs best adapted for the purpose; for, had the larva omitted taking this step, the consequence would have been, that the poor insect, when come to its butterfly state, would have been a prisoner, totally unable to make its escape, being unprovided with any instrument sufficiently powerful to make a hole in the shell; therefore, in this butterfly, it is absolutely necessary that a complete aperture should be left open; and as this is a circumstance which necessarily leaves the pupa exposed, it is not surprising that nature should seldom resort to such a proceeding in the case of internal-feeding insects. Perhaps even in this almost solitary instance, we may fairly imagine that the situation is sufficiently retired to insure them protection from many of their enemies. These chrysalides are attached horizontally upon the inner walls of the pomegranate, by means, first, of a patch of silk laid upon its surface, to the centre of which the tail of the chrysalis is affixed, and, second, of a slender silken thread, passing from side to side over the base of the abdominal segments. As to the manner in which the girthed suspension is effected after the larva has attached itself, I may be allowed to make the following short extract from Messrs. Kirby and Spence's third volume, p. 212. In this order of insects, which have several modus operandi, some of the larvæ which have a short and more rigid body (as *Lycæna Argus*, and many more of the *Papiliones rurales* and *urbicolæ*), 'after having bent the head on one side so as to fix one end of the thread, bring themselves into a straight position, and, by a manœuvre not easily described, contrive to introduce the head under the thread, which they then bend themselves to fasten on the other side, pushing it to its proper situation by the successive tension and contraction of their segments.' A short time after this is effected, the skin of the caterpillar bursts, and its skin is gradually sloughed off beneath the girth, until the pupa is entirely naked; the exuviaæ being collected at the extremity of the abdomen.

Another curious instance of instinct yet remains to be noticed. The butterfly, so soon as ever it has escaped from the puparium, must make its escape out of the hole formed by the larva; delay would be death, as the wings would soon acquire their full expansion of nearly a couple of inches; in which state it would, of course, be unable to creep out.

In the chrysalis state, the belly of the insect is placed in contact with the inner surface of the fruit; consequently, as the slit by which the butterfly escapes out of the puparium extends along the back, the under surface of the latter remains entire, the anterior lateral portions on each side the slit (extending as far as the whole coverings of the wings) curling up and lying close upon those parts which had covered the breast and limbs, leaving the abdominal portion in the same form as when it inclosed the insect."

E. T. Downes, Esq., in the Calcutta Journal of Natural History, observes: "I am induced to call the attention of Entomologists to the caterpillar of the butterfly which is so destructive to the fruit of the pomegranate (*Punicea granatum*), as my observations regarding its habits do not correspond with the account given above. At the time the pomegranate is in flower, and at a very early period, the Hair Streak may be seen very busily occupied about the flowers, and I have little doubt that the eggs are deposited at the bottom of the calyx, from the position in which I have seen the abdomen of the butterfly placed; as the fruit enlarges, the eggs are inclosed, and in this situation matured. Recollecting the history of this insect, I determined to enjoy the pleasure of seeing it go through its various stages (the pomegranate-tree being close to my door). In order that I might obtain a perfect insect, I surrounded several of the fruit with fine gauze, but in such a manner as not in the least to interfere with the caterpillar in its labour of connecting the fruit and stalk by means of the web; but, to my astonishment and disappointment, this never took place; the caterpillars issued from the fruit, and, finding their escape impeded, underwent their change on the external part; and so soon as this was effected, I removed the fruit from the tree for the purpose of placing it in a mosquito gauze house in my room. Subsequently, I examined several of the fruit, but I never found any chrysalidæ, or the remains of any, inside the fruit. I was very careful in my observations, and I came to the conclusion that the caterpillars, in this instance, deserted the fruit when ready to undergo their change; and I have no doubt of the butterfly being the same, viz., '*Thecla Isocrates.*' "

43. *DIPSAS MALIKA*, Horsfield Sp. (Plate 1a, fig. 5).

Thecla Malika, Horsfield, Cat. Lep. Mus. E.I.C. p. 90 (1829).

Dipsas Malika, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 11.

a. ♂. Java. From Dr. Horsfield's Collection.

44. *DIPSAS KESSUMA*, Horsfield Sp.

Thecla Kessuma, Horsfield, Cat. Lep. Mus. E.I.C. p. 89 (1829).

Dipsas Kessuma, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 10.

a. ♀. Java. From Dr. Horsfield's Collection.

45. *DIPSAS NASAKA*, Horsfield Sp.

Thecla Nasaka, Horsfield, Cat. Lep. Mus. E.I.C. p. 91 (1829).

Dipsas Nasaka, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 8.

a. ♂. Java. From Dr. Horsfield's Collection.

46. *DIPSAS VARUNA*, Horsfield Sp.

Thecla Varuna, Horsfield, Cat. Lep. Mus. E.I.C. p. 91 (1829).

Dipsas Varuna, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 480, No. 9.

a. b. ♂. Java. From Dr. Horsfield's Collection.

Genus APHNAEUS, Hübner.

Aphnaeus, Hübner, Verz. bek. Schmett. p. 81 (1816).

Zeritis, Boisduval, MS.

47. *APHNAEUS PINDARUS*, Fabricius Sp.

Hesperia Pindarus, Fabricius, Ent. Syst. III. pt. I. p. 262 (1793). Donovan, Ins. Ind. t. 38, f. 2.

Aphnaeus Pindarus, E. Doubleday, List Lep. Brit. Mus. pt. II. p. 26.

Amblypodia Pindarus, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 479, No. 26.

Zeritis Brahmina, Boisduval, MS.

- a.* ♂. N. India. Presented by the Trustees of the British Museum.
- b.* ♂. N. India. From Mr. Argent's Collection.
- c.* ♂. N. India. Presented by Lieutenant Hugo James, 44th Beng. N. I.
- d.* ♀. Canara. Presented by S. N. Ward, Esq.

48. *APHNÆUS ETOLUS*, Cramer Sp.

- ♀ Papilio Etolus, Cramer, *Pap. Exot.* t. 208, f. E.F. (1782) (*nec. Fabr.*).
- Papilio Vulcanus, Fabricius, *Mant. Ins.* II. p. 66 (1787). *Donovan, Ins. Ind.* t. 38, f. 3.
- Aphnæus Vulcanus, Hübner, *Verz. bek. Schmett.* p. 81. *E. Doubleday, List Lep. Brit. Mus. pt. II.* p. 26.
- Polyommatus Vulcanus, Godart, *Ene. Méth.* IX. p. 644.
- Amblypodia Vulcanus, Horsfield, *Cat. Lep. Mus. E.I.C.* pp. 106, 107. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 479, No. 27.
- a. b.* ♂. India. From the Asiatic Society of Bengal.
- c.* ♂. Canara. Presented by S. N. Ward, Esq.
- d.* ♀. Calcutta. Presented by Colonel Buckley.
- e.* ♀. Java. From Dr. Horsfield's Collection.

49. *APHNÆUS LOHITA*, Horsfield Sp.

- Amblypodia Lohita, Horsfield, *Cat. Lep. Mus. E.I.C.* pp. 106, 107 (1829). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 479, No. 19.

- a. b.* ♂ ♀. Java. From Dr. Horsfield's Collection.

50. *APHNÆUS SYAMA*, Horsfield Sp.

- Amblypodia Syama, Horsfield, *Cat. Lep. Mus. E.I.C.* pp. 107, 108 (1829). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 479, No. 20.

- a. b.* ♂ ♀. Java. From Dr. Horsfield's Collection.

Genus AMBLYPODIA, Horsfield.

AMBLYPODIA, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 98 (1828). *Westwood, Diurn. Lep.* p. 478.

ARHOPALA, Boisduval.

DIPSAS, p. E. Doubleday.

51. AMBLYPODIA NARADA, Horsfield.

♀ Amblypodia Narada, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 98, t. 1, f. 8; t. 4, f. 4, 4a (1829). *Doubleday, List Lep. Brit. Mus. pt. II.* p. 23. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 9.

Theela Talazala, Boisduval, *MS. Mus. E.I.C.*

a. ♀. Java. From Dr. Horsfield's Collection.

b. ♂. Java. From Mr. Argent's Collection.

The larva and pupa of *A. Narada* are figured on Plate 1, fig. 4, 4a.

52. AMBLYPODIA VIVARNA, Horsfield.

Amblypodia Vivarna, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 99 (1829). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 479, No. 10.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♂. Java. From Mr. Argent's Collection.

53. AMBLYPODIA APIDANUS, Cramer Sp.

Papilio Apidanus, Cramer, *Pap. t. 137, f. F.G.* ♂ (1779). *Fabricius, Mant. Ins. II.* p. 69 (1787).

♂ ♀ Amblypodia Apidanus, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 100. *Doubleday, List Lep. Brit. Mus. pt. II.* p. 24. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 3.

Hesperia Apidanus, Fabricius, *Ent. Syst. III. pt. I.* p. 280.

Polyommatus Apidanus, Godart, *Enc. Méth. IX.* p. 652. Papilio Dorimond, Stoll, *Suppl. to Cramer's Pap. t. 37, f. 4, 4D.* ♀ (1791).

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. h. ♂ ♀. Bornco. Presented by Hugh Lowe, Esq.

"In Java, the larva of *Ambl. Apidanus* feeds on the leaves of several species of *Eugenia* and *Calyptranthes*. It occurred not unfrequently, and was from time to time undergoing its change in my breeding-cages."—(Horsfield.)

The larva and pupa are figured on Plate 1, fig. 5, 5a.

54. AMBLYPODIA CENTAURUS, Fabricius Sp.

Papilio Centaurus, Fabricius, Mant. Ins. II. p. 68 (1787).

Amblypodia Centaurus, Doubleday, List Lep. Brit. Mus. pt. II. p. 24. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 478, No. 1.

Polyommatus Centaurus, Godart, Enc. Méth. H. N. IX. p. 658.

a. b. ♂ ♀. Cherra Poonjee. c. ♀. Silhet. Presented by Colonel Buckley.

c. d. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

55. AMBLYPODIA PSEUDO-CENTAURUS, Doubleday.

Amblypodia Pseudo-Centaurus, Doubleday, List. Lep. Brit. Mus. pt. II. p. 24 (1847). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 478, No. 2.

♂ ♀ Amblypodia Centaurus, Horsfield, Cat. Lep. Mus. E.I.C. p. 102 (nec. Fabr.).

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

56. AMBLYPODIA SILHETENSIS, Boisduval Sp.

Thecla silhetensis, Boisduval, Sp. Gén. II. (ined.).

Amblypodia silhethensis, Doubleday, List Lep. Brit. Mus. pt. II. Suppl. p. 36 (1847).

a. b. ♂ ♀. Silhet. Presented by Colonel Buckley.

57. AMBLYPODIA HELUS, Cramer Sp.

♂ Papilio Helus, Cramer, Pap. t. 201, f. F. G. (1782).

Amblypodia Helus, Horsfield, Cat. Lep. Mus. E.I.C. p. 103. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 478, No. 4.

Polyommatus Helus, Godart, Enc. Méth. H. N. IX. p. 652. Lucas, Hist. Nat. Lep. Exot. t. 44, f. 4.

a. ♂. Java. From Dr. Horsfield's Collection.

58. *AMBLYPODIA CAMDEO*, Doubleday (Plate 1a, fig. 6).

Amblypodia Camdeo, Doubleday, *List Lep. Brit. Mus.* pt. II. p. 24 (1847).

- a. b. ♂ ♀. N. India. Presented by Colonel Sykes.
- c. d. ♂. Silhet. Presented by Colonel Buckley.
- e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

Amblypodia Camdeo.—*Male*, UPPER-SIDE pale violet-blue, with a whitish tint in the centre, and a dark mark at the end of the discoidal cell of the fore-wing; extreme edge of anterior margin of fore-wing, ciliæ, tail, and a short tuft at anal angle, dark brown. UNDER-SIDE buff-brown; fore-wing having near the base of discoidal cell a small blackish spot, then a longer square one, and parallelly beyond this another squarish undulated spot; then a series of seven oval and lunate spots across the wing, one between each veinlet, also some irregular-shaped spots disposed at the posterior side of the median vein, all these being encircled by a narrow white band; near and parallel with the exterior margin are two series of pale white marks, the space between these being of a darkish brown. Hind-wing with five black spots near the base (these spots on both wings forming a circle); beyond these are three, then two, and afterwards a series of somewhat smaller spots, this series ending upwards to abdominal margin in a long narrow mark, all being encircled as those in fore-wing; near and parallel with the exterior margin are markings as in fore-wing; near anal angle are disposed some patches of black and metallic blue.

Female, UPPER-SIDE bright cyaneous, with a broad anterior and exterior margin to both wings; anal angle with two or three cyaneous lunar-shaped marks; abdominal margin in both sexes of a creamy colour; body dark brown. UNDER-SIDE as in the male. Expanse of wings $2\frac{3}{8}$ in.

59. *AMBLYPODIA EUMOLPHUS*, Cramer Sp.

♂ Papilio Eumolphus, Cramer, *Pap. Exot.* III. t. 299, f. G.H. (1782).

♂ ♀ Amblypodia Eumolphus, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 103. Doubleday, *List Lep. Brit. Mus.* pt. II. p. 24. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 478, No. 8.

Polyommatus Eumolphus, *Godart, Enc. Méth. H.N.* IX.
p. 652.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

60. *AMBLYPODIA RAMA*, *Kollar Sp.*

♀ Thecla Rama, *Kollar, in Hügel's Kaschmir*, IV. pt. II.
p. 412, t. 4, f. 1, 2 (1844).

a. ♂. Canara. Presented by S. N. Ward, Esq.

b. ♀. N. India. Presented by Colonel Sykes.

61. *AMBLYPODIA PERIMUTA*, *Boisduval.*

Amblypodia Perimuta, *Boisduval, MS.* E. Doubleday,
List of Lep. Brit. Mus. pt. II. p. 24.

a. ♂. Silhet. Presented by the Trustees of the
British Museum.

A. Perimuta.—UPPER-SIDE of the male glossy purple, with a narrow black border on exterior margins. UNDER-SIDE brown, glossed over with purple; fore-wing with two oval marks and an undulating line of a paler colour; hind-wing with a broad, uneven, and indistinctly-defined fascia of a creamy colour crossing from the anterior to abdominal margin, these wings being also marked with several oval and undulating lines; fore-wing pointed at the anterior angle; hind-wings rounded, without tails. Expanse of wings $1\frac{5}{12}$ in.

62. *AMBLYPODIA EPIMUTA*, *Boisduval.*

Amblypodia Epimuta, *Boisduval, MS.* E. Doubleday,
List Lep. Brit. Mus. pt. II. p. 24.

a. ♂. Borneo. Presented by Hugh Lowe, Esq.

A. Epimuta.—UPPER-SIDE of the male wholly glossy purple. UNDER-SIDE brown, and covered with a series of undulating lines; anal angle of hind-wing with a patch of metallic blue. Wings somewhat rounded. Without tails. Expanse of wings $1\frac{7}{10}$ in.

63. *AMBLYPODIA QUERCETORUM*, *Boisduval* (*Plate 1a,*
fig. 7.).

Amblypodia Quercetorum, *Boisduval, MS.* Doubleday,
List Lep. Brit. Mus. pt. II. p. 25 (1847).

a. ♂. Silhet. Presented by the Trustees of the British Museum.

b. c. ♂. N. India. Donor unknown.

A. Quercetorum.—UPPER-SIDE dark brown; centre of wings deep purple. UNDER-SIDE brown; *fore-wing* with a series of small dark spots near the exterior margin, then an undulating line, and towards the base some indistinct spots; *hind-wing* with a dark brown fascia running from the anterior angle across to middle of abdominal margin; an undulating dark line near exterior margin, and two small whitish elongated spots near anterior margin; anterior and posterior angle of *fore-wing* pointed; *hind-wing* with the anal angle elongated. Expanse of wings $1\frac{5}{8}$ in.

64. *AMBLYPODIA QUERCETI*, Boisduval.

Amblypodia Querceti, *Boisduval*, MS. *Doubleday, List Lep. Brit. Mus. pt. II. p. 25 (1847)*.

a. b. c. d. e. ♂ ♀. N. India. Presented by Colonel Buckley.

A. Querceti.—Male, UPPER-SIDE purple; *fore-wing* with broadish brown exterior margin commencing from middle of anterior margin; *hind-wing* with broad brown margins. Female, UPPER-SIDE with discoidal cell and posterior base of *fore-wing*, and middle of *hind-wing*, purple, the disc of *fore-wing* being somewhat lighter and pinky; the rest of the wings brown. UNDER-SIDE greyish-brown, slightly rufescent on *fore-wing*; an undulating broad band of rufescent brown crosses the fore and *hind-wings*, another inwardly on the latter half across the wing, then two or three spots; near exterior margin of both wings is an indistinct angulated line; *fore-wing* angulated exteriorly; *hind-wing* rounded; tail, one. Expanse of wings $1\frac{1}{2}$ in.

65. *AMBLYPODIA DODONÆA*, Boisduval (*Plate 1a, fig. 8*).

Amblypodia Dodonæa, *Boisduval*, MS. *Doubleday, List Lep. Brit. Mus. pt. II. p. 25 (1847)*.

a. b. c. d. ♀. N. India. Presented by Colonel Buckley.

A. Dodonæa.—Female, UPPER-SIDE, *fore-wing* with discoidal cell and posterior base blue, palest in the centre of disc, and intersected by dark veinlets, the rest of wing dusky-brown; *hind-wing* with

centre, blue; outer margin brown; anterior and abdominal margins paler. UNDER-SIDE dark cream-colour; *fore-wing* paler posteriorly, with an ill-defined band near outer margin, then a broad undulating band, and some spots of a darker shining tint; *hind-wing* also with an ill-defined but more angulated band near outer margin, then two series of undulating lines, and also spots at base of wing. Shape of wings as in *A. Querceti*, but exterior margin of fore-wing more angulated. Expanse of wings $1\frac{5}{8}$ in.

66. *AMBLYPODIA GANESA*, Moore (*Plate 1a, fig. 9*).

A. Ganesa, n. sp.—UPPER-SIDE, *fore-wings*, with discoidal cell and posterior base, sky-blue; the end of discoidal cell and between discoidal veinlets white; rest of wing dark brown; *hind-wing* sky-blue to near exterior margin, rest brown; abdominal margin and ciliae paler. UNDER-SIDE cream-white; *fore-wings* nearly covered with broad undulating brown bands; *hind-wings* with ill-defined undulating bands; near outer margin of both wings appear a very faint undulated line and a series of dots. Wings shaped as in *A. Querceti* and *A. Dodonæa*. Without tails. Expanse of wings $1\frac{1}{4}$ in.

a. b. c. d. N. India. Presented by Colonel Buckley.

67. *AMBLYPODIA TIMOLEON*, Stoll Sp.

Papilio Timoleon, Stoll, *Suppl. to Cramer, Pap. t. 32, f. 4, 4D* (1791).

Amblypodia Timoleon, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 478, No. 23.

Thecla Nila, Kollar, in Hügel's *Kaschmir*, p. 413, t. IV. f. 5, 6 (1844), (nec. Th. Nila, Horsf.).

Amblypodia Lyco, Doubleday, *List Lep. Brit. Mus. pt. II.* p. 23 (1847).

a. ♂. Bootan. (*A. Lyco*, Doubleday.) From Pemberton's Collection.

68. *AMBLYPODIA ROCHANA*, Horsfield (*Plate 1a, f. 10*).

♂ Amblypodia Rochana, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 108 (1829).

Thecla Timoleon, Boisduval, *Sp. Gén. Lép. t. 22, f. 4?*

a. ♂. Java. From Dr. Horsfield's Collection.

69. *AMBLYPODIA VIDURA*, Horsfield.

♂ Amblypodia Vidura, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 113, t. 1, f. 6, 6a (1829). *E. Doubleday, List Lep. Brit. Mus. pt. II. p. 23. Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 18.

a. ♂. Java. From Dr. Horsfield's Collection.

70. *AMBLYPODIA LONGINUS*, Fabricius Sp.

Hesperia Longinus, *Fabricius*, *Ent. Syst. V.* p. 430 (1798).

Amblypodia Longinus, *Doubleday, List Lep. Brit. Mus. pt. II. p. 23. Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 16.

Polyommatus Longinus, *Godart, Enc. Meth. IX.* p. 634. *Lucas, Lep. Exot. t. 44, f. 3.*

Bithys Longinus, *Hübner, Zuträge*, f. 933, 934 (1837).

a. b. c. ♂ ♀. N. India. Presented by the Asiatic Society of Bengal.

71. *AMBLYPODIA PSEUDOLONGINUS*, Doubleday.

Amblypodia Pseudolonginus, *Doubleday, List Lep. Brit. Mus. pt. II. p. 23 (1847. Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 17.

♂ ♀ Amblypodia Longinus, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 110, t. 1, f. 7 (nec. *Fabr.*).

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

"The larva of *Amblyp. Pseudolonginus* feeds on the leaves of a species of *Loranthus*, and was observed in the month of February."
—(Horsfield, MS.)

The larva and pupa of *A. Pseudolonginus* are figured on Plate 1, fig. 6, 6a.

72. *AMBLYPODIA HYPATADA*, Boisduval Sp.

Thecla Hypatada, *Boisduval, MS.*

Myrina Hypatada, *Doubleday, List Lep. Brit. Mus. pt. II. p. 23.*

a. b. ♂. N. India. Presented by Colonel Buckley.

A. Hypatada.—Male, UPPER-SIDE, *fore-wing*, with discoidal cell, posterior base to posterior angle, brilliant light sky-blue; rest of wing dark brown; *hind-wing* brilliant light sky-blue, with brown ciliae; abdominal margin cream-colour. UNDER-SIDE light creamy-brown, with a well-defined narrow undulating band of reddish-brown crossing the *fore* and *hind-wings*, terminating abdominally in a zig-zag manner on the latter; near exterior margins an indistinct line; near anal angle two black spots, bordered anteriorly with red, and posteriorly with powdered white. Tails broken off. Expanse of wings $1\frac{5}{12}$ in.

73. *AMBLYPODIA NISSA*, Kollar Sp.

Thecla Nissa, Kollar, in Hügel's Kaschmir, IV. pt. II.
p. 412, t. 4, f. 3, 4 (1844).

a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.

74. *AMBLYPODIA DEVA*, Moore.

Ambl. Deva, n. s.—Male, UPPER-SIDE, *fore-wing*, with discoidal cell and posterior base, brilliant deep blue, intersected by the dark median and submedian veinlets; costal margin and anterior half of wing to beyond posterior angle dark brown; *hind-wing*, with the centre from the base, brilliant deep blue; a broad anterior and narrow exterior margin of dark brown; along the exterior margin are disposed some blackish marks; a black spot at the anal angle, surrounded with whitish; abdominal margin pale brown; tails two, brown; ciliae whitish. UNDER-SIDE cream-colour, with a series of interrupted marks disposed in an undulating line across the *fore* and *hind-wings*, and terminating in a zig-zag manner abdominally; anal angle with a black spot, bordered anteriorly with red, and another of the same a short distance off on the exterior margin. Female, UPPER-SIDE paler blue, and with lighter brown margins. Expanse of wings in male $1\frac{2}{3}$ in.; female $1\frac{5}{8}$ in.

a. b. ♂ ♀. Canara. Presented by S. N. Ward, Esq.

75. *AMBLYPODIA JANGALA*, Horsfield (Plate 1a, f. 11).

♀ Amblypodia Jangala, Horsfield, Cat. Lep. Mus. E.I.C.
p. 113 (1829). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 478, No. 22.

a. ♀. Java. From Dr. Horsfield's Collection.

b. c. ♂. Darjeeling. From the Indian Collection, Exposition Universelle at Paris, 1855.

A. Jangala.—Male, UPPER-SIDE brownish-black, with the discoidal cell and posterior base of the fore-wing and centre of hind-wing shining indigo-blue.

Genus MYRINA, Godart.

MYRINA, Godart, Enc. Méth. IX. p. 592 (1819).

76. *MYRINA JAFRA*, Godart.

Myrina Jafra, Godart, Enc. Méth. IX. p. 592 (1819).

Horsfield, Cat. Lep. Mus. E.I.C. p. 118, t. 11, f. 5, 5a. Lucas, Hist. Nat. Lep. Exot. t. 43, f. 4. Guérin-Méneville, Icon. Règ. An. Ins. p. 489, t. 81, f. 3. Doubleday, List Lep. Brit. Mus. pt. II. p. 21. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 476, No. 1.

a. b. c. ♂ ♀. Java: From Dr. Horsfield's Collection.

77. *MYRINA ACTE*, Doubleday.

Myrina Acte, Doubleday, List Lep. Brit. Mus. pt. II. p. 21 (1847).

a. N. India. Presented by the Trustees of the British Museum.

b. c. N. India. Presented by Lieutenant Hugo James, 44th Bengal N. I.

M. Acte.—UPPER-SIDE dark smoky-brown; hind-wing with three spots of white near anal angle; tail and anterior margin whitish. UNDER-SIDE reddish chrome-yellow, with an indistinct undulating line crossing the fore and hind-wings, terminating abdominally on the latter in separate lighter spots; near anal angle brownish, with two black spots bounded anteriorly with metallic blue. Expanse of wings $1\frac{1}{2}$ in.

78. *MYRINA THARIS*, Hübner Sp.

Oxylides Tharis, Hübner, Zutr. f. 883, 884 (1837).

Myrina Tharis, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 476, No. 12.

Myrina Pharis, *E. Doubleday, List Lep. Brit. Mus. pt. II.*
p. 22 (1847); Diurnal Lep. t. 74, f. 3.

a. India. Presented by the Trustees of the British Museum.

M. Tharis.—UPPER-SIDE brown; fore-wing with a broad band from angle of abdominal margin across the wing to middle of exterior margin; a spot near to, and the margin of anal angle, tails and broad ciliae pure white. UNDER-SIDE, with fore-wing chrome-red; basal half of hind-wing slightly paler; anal half white, with patches of black disposed at anal angle and between the veinlets along exterior margin; across the wing, from abdominal margin to near anterior angle (separating the red and white colours), is a zig-zag line. Expanse of wing $1\frac{1}{2}$ in.; middle tail $\frac{1}{2}$ in.

79. *MYRINA LAPITHIS*, Boisduval Sp.

Theela Lapithis, *Boisduval, MS. (1847)*.

Amblypodia Lapithis, *E. Doubleday, List Lep. Brit. Mus. pt. II. p. 23 (1847)*.

a. Moulmein. Presented by the Trustees of the British Museum.

M. Lapithis.—UPPER-SIDE glossy-brown; anal part of hind-wing tail, and ciliae, white, with a patch of light glistening blue at anal angle and between median veinlets. UNDER-SIDE, fore-wing chrome-yellow, darker on anterior half; a white fascia from middle of anterior margin, spreading widely across to posterior margin, and bordered by blackish; three separate blackish short lines across the anterior half of the wing; hind-wing white; extreme base and anterior angle yellow; at base and in centre of wing are disposed several spots and short lines of black; anal angle black; a black spot at angle on exterior margin, these being bounded anteriorly with metallic silvery-green, and then yellow. Expanse of wings $1\frac{1}{4}$ in.

80. *MYRINA LISIAS*, Fabricius Sp.

Papilio Lisias, *Fabricius, Mant. Ins. II. pp. 65, 261 (1787); Ent. Syst. III. pt. I. p. 261.*

Myrina Lisias, *Godart, Enc. Méth. IX. p. 593. Boisduval, Sp. Gén. I. t. 22, f. 2. E. Doubleday, List Lep. Brit. Mus. pt. II. p. 21. Westwood, in*

Doubleday and Hewitson's Diurnal Lep. p. 476,
No. 3.

a. ♀. Moulmein. Presented by the Trustees of the
British Museum.

81. *MYRINA TRIOPAS*, Cramer Sp.

Papilio Triopas, Cramer, *Pap. Exot.* IV. t. 320, f. G.H.
(1782).

Papilio Amor, Fabricius, *Mant. Ins.* II. p. 65 (1787).
Herbst, *Pap.* t. 302, f. 9, 10.

Myrina Amor, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 4.

Polyommatus Amor, Godart, *Enc. Méth.* IX. p. 620.

Zephyrius Amor, Guérin-Méneville, *Icon. Rég. An. Ins.*
p. 490, t. 81, f. 6.

a. Canara. Presented by S. N. Ward, Esq., E.I.C.C.S.

82. *MYRINA ETOLUS*, Fabricius Sp.

Papilio Etolus, Fabricius, *Mant. Ins.* II. p. 66 (1787);
Ent. Syst. III. pt. I. p. 264 (nec. Cramer, 1782).

Myrina Etolus, E. Doubleday, *List Lep. Brit. Mus.*
pt. II. p. 22. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 9.

Polyommatus Etolus, Godart, *Enc. Méth.* IX. p. 639.

Amblypodia Etolus, Horsfield, *Cat. Lep. Mus. E.I.C.*
p. 115, t. 1, f. 9.

a. ♂. Java. From Dr. Horsfield's Collection.

b. c. ♂ ♀. Dukhun. Presented by Colonel Sykes.

d. ♂. Silhet. Presented by Colonel Buckley.

83. *MYRINA AMYNTOR*, Herbst Sp.

Papilio Amyntor, Herbst, *Pap.* t. 300, f. 5, 6.

Myrina Amyntor, E. Doubleday, *List Lep. Brit. Mus.*
pt. II. p. 21. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 15.

a. b. c. ♂ ♀. N. India. From Mr. Argent's Collection.

d. ♀. Silhet. Presented by Colonel Buckley.

84. *MYRINA ERYLUS*, Godart Sp.

Polyommatus Erylus, Godart, *Enc. Méth.* IX. p. 633 (1819).

Myrina Erylus, E. Doubleday, *List Lep. Brit. Mus.* pt. II. p. 22. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 14.

Amblypodia Erylus, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 111.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. ♂. Silhet. Presented by the Trustees of the British Museum.

e. ♂. India. From Mr. Argent's Collection.

85. *MYRINA JALINDRA*, Horsfield Sp.

♀ *Amblypodia Jalindra*, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 109 (1829). Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 478, No. 21.

♀ *Polyommatus Nedymond*, Godart, *Enc. Méth.* IX. p. 634 (1819) (nec. Cramer).

a. ♀. Java (type). From Dr. Horsfield's Collection.

b. ♂. Silhet. Presented by Colonel Buckley.

c. ♀. Canara. Presented by S. N. Ward, Esq.

M. Jalindra.—Male, UPPER-SIDE varying violet-blue; exterior margin of fore-wing, and anterior, exterior, and abdominal margin of hind-wing, brown; anal angle whitish. UNDER-SIDE as in female.

86. *MYRINA NEDYMOND*, Cramer Sp.

♂ *Papilio Nedymond*, Cramer, *Pap. Exot.* III. t. 299, f. E.F. (1782), (nec. Godart, *Enc. Méth.*).

Myrina Nedymond, E. Doubleday, *List of Lep. Brit. Mus.* pt. II. p. 22. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 7.

Thecla Nedymond, Horsfield, *Cat. Lep. Mus. E.I.C.* p. 96 (1829).

a. b. ♂. Java. From Dr. Horsfield's Collection.

87. *MYRINA CHITRA*, Horsfield Sp.

♀ Thecla Chitra, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 97, t. 1, f. 5 (1829).

Myrina Chitra, *E. Doubleday*, *List Lep. Brit. Mus. pt. II.* p. 22. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 8.

a. b. ♀. Java. From Dr. Horsfield's Collection.

88. *MYRINA RAVINDRA*, Horsfield.

♂ ♀ Myrina Ravindra, *Horsfield*, *Cat. Lep. Mus. E. I. C.* p. 117, t. 1, f. 11, 11a (1829). *Doubleday, List Lep. Brit. Mus. pt. II.* p. 22. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 5.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

89. *MYRINA SUGRIVA*, Horsfield Sp. (Plate 1a, fig. 12).

Amblypodia Sugriva, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 105, t. 1, f. 10, 10a (1829).

Myrina Sugriva, *E. Doubleday*, *List Lep. Brit. Mus. pt. II.* p. 21. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 476, No. 2.

a. Java. From Dr. Horsfield's Collection.

Genus *LOXURA*, Horsfield.

LOXURA, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 119 (1829). *Westwood, Diurn. Lep.* p. 475.

MARMESSUS, p. Hübner.

MYRINA, p. Godart.

90. *LOXURA ATYMNUS*, Cramer Sp.

Papilio Atymnus, *Cramer*, *Pap. Exot. IV.* t. 331, f. D.E. (1782).

Loxura Atymnus, *Horsfield*, *Cat. Lep. Mus. E. I. C.* p. 121. *Boisduval*, *Sp. Gén. Lép. I.* t. 7, f. 3; t. 11, f. 6, 6a. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 475. *Kollar*, in *Hügel's Kaschmir*, p. 411.

- Hesperia Atymnus, *Fabricius*, *Mant. Ins.* II. p. 70 (1787); *Ent. Syst.* III. pt. I. p. 283.
 Myrina Atymnus, *Godart*, *Enc. Méth.* IX. p. 594.
E. Doubleday, *List Lep. Brit. Mus.* pt. II. p. 21.
 Papilio Atymnus, *Donovan*, *Ins. of India*, t. 39, f. 1.
 Marmessus Atymnus, *Hübner*, *Verz. bek. Schmett.* n. 828.
a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.
f. g. h. N. India. Presented by Colonel Buckley.
i. Canara. Presented by S. N. Ward, Esq.

The female of this species, according to the observation of Dr. Horsfield, "can be distinguished only by the structure of the anterior tarsi."

91. *LOXURA SURYA*, Moore (*Plate 1a, fig. 13*).

L. Surya, *n. sp.*—Differs from *L. Atymnus* in having the fulvous colour of the UPPER-SIDE more intense and brighter; the blackish-brown border of the *fore-wing* is also darker, and which extends to the base of the wing, occupying the whole space between the costal vein and anterior margin; a patch of blackish-brown near base of anterior margin of *hind-wing*, abdominal margin brown. UNDER-SIDE dark ochreous.

- a.* Canara. Presented by S. N. Ward, Esq.

92. *LOXURA PITA*, Horsfield.

- ♀ *Loxura Pita*, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 122 (1829). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* t. 74, f. 1, 2, p. 472.
Myrina Pita, *E. Doubleday*, *List Lep. Brit. Mus.* p. 21.
a. ♀. Java. From Dr. Horsfield's Collection.

Genus ANOPS, Boisduval.

ANOPS, *Boisduval*, *Sp. Gén. Lép.* I. p. 6 (1836). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 473.

PHÆDRA, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 124 (1829).

93. *ANOPS THETYS*, Drury *Sp.*

- ♀ *Papilio Thetys*, *Drury*, *Ill. Ins.* t. 9, f. 3, 4 (1770).
Cramer, *Pap. Exot.* III. t. 238, f. D.

- Anops Thetys, *Doubleday, List Lep. Brit. Mus.* II. p. 20.
♂ Papilio Phædrus, *Fabricius, Mant. Ins.* II. p. 79 (1787);
Syst. Ent. III. pt. I. p. 307.
Anops Phædrus, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 473.
Polyommatus Phædrus, *Godart, Enc. Méth. Hist. Nat.* IX. p. 675.
♂ Candalides Phædrus, *Hübner, Verz. bek. Schmett.* n. 720 (1816).
♀ Papilio Æsopus, *Fabricius, Mant. Ins.* II. p. 79; *Syst. Ent.* III. pt. I. p. 307.
Curetis Æsopus, *Hübner, Verz. n.* 1070 (1816).
♂ ♀ Phædra terricola, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 124 (1829).
a. b. ♂. N. India. Purchased.
c. ♀. Canara. Presented by S. N. Ward, Esq.

94. *ANOPS INSULARIS*, Horsfield Sp. (Plate 1a, fig. 14).

- ♂ Phædra insularis, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 125 (1829).
Anops insularis, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 473.

a. ♂. Java. From Dr. Horsfield's Collection.

95. *ANOPS CINYRA*, Cramer Sp.

- ♂ Papilio Cinyra, *Cramer, Pap. Exot.* III. t. 238, f. C. (1782).
Candalides Phædrus, *apud Hübner, Zuträge*, f. 263, 264 (*nec. Fabr.*).

a. b. ♂. Java. From Dr. Horsfield's Collection.

In the three preceding species the hind-wings are more rounded than in *A. Bulis* and *A. Santana*.

96. *ANOPS BULIS*, Boisduval.

- ♂ Anops Bulis, *Boisduval, MS.* (1847). *E. Doubleday, List Lep.* pt. II. p. 21. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 473, t. 75, f. 5 (1852).

- a.* ♂. Bootan. From Pemberton's Collection.
- b. c.* ♂. Silhet. Presented by Colonel Buckley.
- d.* ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- e.* ♂. N. India. Donor unknown.

The female of this species may be distinguished by the white patches on the *UPPER-SIDE* of both wings being larger; the *fore-wings* having a disco-cellular dentiform mark of black. *UNDER-SIDE* silvery creamy-white, and having a darkish band running from anterior angle of fore-wing in a straight line to anal angle of hind-wing, the whole surface of the under-side being covered with very minute dark-brown dots.

97. *ANOPS SANTANA*, Moore.

Anops Phædrus, *apud Boisduval*, *Spéc. Gén. Lép.* I.
t. 23, f. 1.

- a.* ♂. Java. From Dr. Horsfield's Collection.
- b. c.* ♂. Java. From Mr. Argent's Collection.

Stirps II., with CHILOGNATHIFORM or JULIFORM Larva.

CHILOGNATHIFORM OR JULIFORM (*part*), *Horsfield, Cat. Lep. Mus. E.I.C.* pp. 38, 127 (1828). *Swainson, Cabinet Cyclop.* pp. 58, 63 (1840).

A. PAPILIO (*part*), *Linnæus—DANAI CANDIDI.*

B. PAPILIO (*part*), *Linnæus—EQUITES, TROES et ACHIVI.*

A.

PAPILIONES (*part*), *Horsfield, Cat. Lep. Mus. E.I.C.* p. 59 (1828).

PIÉRIDES, *Boisduval, Hist. Gén. des Lép. Amér. Sep.* p. 2 (1829); id. *Spéc. Gén. Lép. I.* pp. 163, 402 (1836); id. *Index Méth. Europ. Lép.* p. 4 (1840). *Westwood, Introd. to Mod. Class. Ins. II.* p. 349; *Syn.* p. 87 (1840).

COLIADES, *Boisduval, Hist. Gén. des Lép. Amér. Sep.* p. 2 (1829).

PIERINÆ et COLIANÆ, *Swainson, Cabinet Cyclop.* pp. 87–9 (1840).

PIERIDÆ, *Dup. C.* p. 23 (1846). *E. Doubleday, Diurnal Lep.* p. 32.

RHODOCERIDÆ, *Dup. C.* p. 26 (1846).

PIERIDI, *Stephens, Cat. Brit. Lep. Brit. Mus.* p. 3 (1850). *Stainton, Manual Brit. Butt.* pp. 12, 15 (1856).

RHODOCERIDI, *Stephens, Cat. Brit. Lep. Brit. Mus.* p. 2 (1850).

LEP. HEXAPODA (*part*), *Latreille, Règ. Anim. V.* p. 374 (1829).

PAPILIONIDÆ (*part*), *Swainson, Cabinet Cyclop.* p. 86 (1840). *Stainton, Manual Brit. Butt.* pp. 10, 12 (1856).

DANAI FLAVI, MM. *Denis et Schieffermüller, Wien. Verz.* p. 164 (1776).

Genus CALLIDRYAS, *Boisduval.*

CALLIDRYAS, *Boisduval, Hist. Gén. et Icon. des Lép. et Chen. Amér. Sep.* p. 73 (1829); id. *Spéc. Gén. Lép. I.* p. 607. *Doubleday, Diurnal Lep.* p. 66.

COLIAS, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 128 (1829).

CATOPSILIA, MURTIA, PHÆBIS, COLOTIS, *Hübner, Verz.* (1816).

98. *CALLIDRYAS PYRANTHE*, *Linnæus Sp.*

- Papilio Pyranthe, *Linnæus, Syst. Nat.* II. p. 763 (1767).
Fabricius, Spec. Ins. II. p. 44 (1781).
Callidryas Pyranthe, *Boisduval, Spéc. Gén. Lép.* I. p. 611 (1836). *Doubleday, List Lep. Brit. Mus.* pt. I. p. 38 (1844); id. *Diurnal Lep.* p. 67.
Colias Pyranthe, *Godart, Enc. Méth.* IX. p. 97 (1819).
Horsfield, Cat. Lep. Mus. E.I.C. p. 129.
Papilio Chryseis, *Drury, Ins.* I. t. 12, f. 3, 4 (1770).
Papilio Gnoma, *Fabricius, Spec. Ins.* II. p. 50; id. *Syst. Ent. App.* p. 828 (1775); *Ent. Syst.* III. pt. I. p. 210.
Papilio Nephte, *Fabricius, Ent. Syst.* III. pt. I. p. 190 (1793).
Papilio Alcyone, *Cramer, Pap.* I. t. 58, f. A. B. C. (1779).
a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
c. d. ♂ ♀. Pinang. Presented by Dr. Cantor, 1844.

The larva and pupa of *C. Pyranthe* are figured on Plate 1, fig. 8, 8a, discovered in Ceylon by E. L. Layard, Esq., from whose drawings our figures are copied.

99. *CALLIDRYAS PHILIPPINA*, *Cramer Sp.*

- Papilio Philippina, *Cramer, Pap.* t. 361, f. C.D. (1782).
Callidryas Philippina, *Boisduval, Spéc. Gén. Lép.* I. p. 609 (1836). *Doubleday, List Lep. Brit. Mus.* pt. I. *App.* p. 147; *Diurnal Lep.* p. 67 (1847).
Colias Philippina, *Hübner, Verz. bek. Schmett.* p. 99 (1816). *Godart, Enc. Méth.* IX. p. 96 (1819).
Horsfield, Cat. Lep. Mus. E.I.C. p. 130 (1829).
a. b. c. d. ♂ ♀. India. From the Asiatic Society of Bengal.

100. *CALLIDRYAS ALCMEONE*, *Cramer Sp.*

- Papilio Alemeone, *Cramer, Pap.* II. t. 141, f. E. (1779).
Fabricius, Spec. Ins. II. p. 44 (1781); id. *Ent. Syst.* III. pt. I. p. 196 (1793).
Callidryas Alemeone, *Boisduval, Spéc. Gén. Lép.* p. 627.

*Doubleday, List Lep. Brit. Mus. pt. I. p. 40; id.
Diurnal Lep. p. 68.*

Colias Alcmeone, *Godart, Enc. Méth. IX. p. 97. Horsfield, Cat. Lep. Mus. E.I.C. p. 131.*

Catopsilia Alcmeone, *Hübner, Verz. bek. Schmett. p. 98 (1816).*

Var. ♀ Papilio Jugurtha, *Cramer, Pap. Exot. II. t. 187, f. E.F. (1779).*

Colias Jugurthina, *Godart, Enc. Méth. IX. p. 96. Horsfield, Cat. Lep. Mus. E.I.C. p. 132.*

? Var. ♀ Papilio Crocale, *Cramer, Pap. Exot. I. t. 55, f. C.D.*

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. e. f. Var. ♀. Java. From Dr. Horsfield's Collection.

g. h. ♂ ♀. N. India. From the Asiatic Society of Bengal.

i. j. Var. ♀. India. From the Asiatic Society of Bengal.

"This species is very abundant in Java."—(Horsfield.)

The larva and pupa of *C. Alcmeone* are figured on Plate 1, fig. 7, 7a, discovered in Ceylon by E. L. Layard, Esq.

101. *CALLIDRYAS HILARIA*, *Cramer Sp.*

♂ Papilio Hilaria, *Cramer, Pap. IV. t. 339, f. A.B. (1782).*

Callidryas Hilaria, *Boisduval, Spéc. Gén. Lép. I. p. 626. Doubleday, List Lep. Brit. Mus. pt. I. p. 39; id. Diurnal Lep. p. 68.*

♂ Colias Hilaria, *Godart, Enc. Méth. IX. p. 97. Horsfield, Cat. Lep. Mus. E.I.C. p. 132.*

Catopsilia Hilaria, *Hübner, Verz. bek. Schmett. p. 98.*

Papilio Titania, *Fabricius, Ent. Syst. V. p. 28 (1793).*

♀ Papilio Catilla, *Cramer, Pap. III. t. 229, f. D.E. (1782). Fabricius, Ent. Syst. III. pt. I. p. 209.*

Colias Catilla, *Godart, Encycl. Méth. IX. p. 95.*

Var. ♀ Papilio Pomona, *Fabricius, Ent. Syst. III. pt. I. p. 213.*

- a. b. c. d.* ♂ ♀. Java. From Dr. Horsfield's Collection.
e. ♂. Pinang. Presented by Dr. Cantor, 1844.
f. g. ♂ ♀. N. India. From the Asiatic Society of Bengal.
h. ♀. N. India. Presented by Colonel Buckley.

102. *CALLIDRYAS SCYLLA*, *Linnæus Sp.*

- Papilio Scylla, *Linnæus*, *Syst. Nat.* II. p. 763 (1767).
 Cramer, *Pap.* I. t. 12, f. C.D. Sulz., *Ins. edit.*
Ræm. t. 15, f. 6. Fabricius, *Spec. Ins.* II. p. 47 (♀);
 id. *Ent. Syst.* III. pt. I. p. 201. Donovan, *Ins. Ind.*
t..
 Callidryas Scylla, *Boisduval*, *Spéc. Gén. Lép.* I. p. 631.
Doubleday, *List Lep. Brit. Mus.* pt. I. p. 40; id.
Diurnal Lep. p. 68.
 Colias Scylla, *Hübner*, *Verz. bek. Schmett.* p. 99. Godart,
Enc. Méth. IX. p. 95. Horsfield, *Cat. Lep. Mus.*
E.I.C. p. 133.
 ♂ Papilio Cornelius, *Fabricius*, *Mant. Ins.* II. p. 21 (1787).
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

"In Java, the larva of this species feeds on various species of *Cassia*, particularly on the *C. fistula* and *C. obtusifolia*; it is, however, occasionally found on other plants. It is extremely abundant, particularly in the early part of the rainy season, after the renewal of the foliage of these plants. I must, however, remark, that the various species of *Colias* resemble each other so closely in the larva state, that the distinction of species is extremely difficult. The greatest number of specimens obtained by breeding belong to this species (*C. Scylla*); but in examining the series put up in Java in illustration of the drawings prepared during the progress of this research, I observed several specimens of *C. Jugurthina* and *C. Alcmeone*, a circumstance which shows their near resemblance in the larva state. Among these specimens is contained that remarkable variety of *C. Jugurthina* figured by Cramer with the name of *Papilio Crocale*."—(Horsfield.)

The larva and pupa of *C. Scylla* are figured on Plate 1, fig. 9, 9a, from Java.

Genus GONEPTERYX, Leach.

GONEPTERYX, *Leach, Edinb. Enc. IX.* p. 128 (1815).

ANTEOS, *Hübner, Verz.* p. 99 (1816).

GONOPTYRYX, *Boisduval, Spéc. Gén. Lép. I.* p. 598 (1836).

GONIAPTERYX, *Westwood, Introd. Syn.* p. 87 (1839).

RHODOCERA, *p. Boisduval.*

103. GONEPTERYX NIPALENSIS, E. Doubleday.

Gonepteryx Rhamni, G. R. Gray, Lep. Ins. of Nepal,
t. 5, f. 1 (nec. Linn.).

Gonepteryx nepalensis, E. Doubleday, Diurnal Lep.
p. 71, n. 9 (1847).

a. ♂. N. India. From the Asiatic Society of Bengal.

b. c. ♂ ♀. N. India. Presented by Colonel Hearsey.

d. ♂. Ladakh. Presented by Captain R. Strachey.

e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus DERCAS, Boisduval.

DERCAS, *Boisduval, MS.* (*Doubleday, Diurn. Lep.* p. 70, 1847).

104. DERCAS VERHUELLI, Van der Hoeven Sp.

Colias Verhuellii, Van der Hoeven, Tijdschrift voor Nat.
Gesch. t. 8, f. 3, 4 (1838).

Dercas Verhuellii, Boisduval, MS.

Gonepteryx Verhuellii, E. Doubleday, Diurnal Lep. t. 8,
f. 3, p. 71, n. 8; id. List Lep. Brit. Mus. App.
p. 10 (1848).

♀ Rhodocera Lycoreias, *E. Doubleday, Gray's Zool. Misc.*
p. 77 (1842).

a. b. ♂ ♀. N. India. Purchased.

c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

This genus differs from *Gonepteryx (Rhamni)* "in having the claws furnished with a distinct pulvillus, and in its much slenderer feet and antennæ."—(E. Doubleday.)

Genus HEBOMOIA, Hübner.

HEBOMOIA, Hübner, Verz. bek. Schmett. p. 95 (1816).

IPHIAS, Boisduval, Spéc. Gén. Lép. I. p. 595 (1836).

PIERIS, p. Godart.

COLIAS, p. Horsfield.

105. HEBOMOIA GLAUCIPPE, Linnæus Sp.

Papilio Glauceppe, Linnæus, Syst. Nat. II. p. 762 (1767).

Clerck, Icones, t. 35, f. 1. Drury, Ins. I. t. 10, f. 3, 4 (1770). Fabricius, Spec. Ins. II. p. 45 (1781); id. Ent. Syst. III. pt. I. p. 198. Cramer, Pap. II. t. 164, f. A.B.C.

Hebomoia Glauceppe, Hübner, Verz. bek. Schmett. p. 95.

Doubleday, Diurnal Lep. p. 63.

Pieris Glauceppe, Godart, Enc. Méth. IX. p. 119.

Colias Glauceppe, Horsfield, Cat. Lep. Mus. E.I.C. p. 130.

Iphias Glauceppe, Boisduval, Spéc. Gén. Lép. I. p. 596.

Doubleday, List Lep. Brit. Mus. pt. I. p. 37.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♂. Borneo. Presented by H. Lowe, Esq.

d. e. ♂ ♀. Dukhun. Presented by Colonel Sykes.

f. ♂. N. India. Presented by Colonel Buckley.

"This genus, of which as yet we know but two species, is easily known by its large size, the gradually thickening antennæ, the peculiar form of the palpi, and the presence of a disco-cellular nervule. The minute terminal joints of the palpi and the form of the pupa indicate an affinity to *Anthocharis*; its antennæ and the form of the larva point out an equal one to [*Colias*] *Callidryas*

"This insect is not unfrequent in Java; the larva feeds on a species of *Capparis*, distinguished by the native name of Wanwannan: February to April."—(Horsfield.)

The larva and pupa of *H. Glauceppe* are figured on Plate 1, fig. 10, 10a; from Java.

Genus ERONIA, Hübner.

ERONIA, Hübner, Samml. Exot. Schmett. (1806-27).

VALERIA, Horsfield, Cat. Lep. Mus. E.I.C. p. 139 (1829).

MYLOTHRIS, p. Hübner.

106. *ERONIA VALERIA*, Cramer Sp.

- ♂ Papilio Valeria, Cramer, *Pap. Exot.* I. t. 85, f. A. (1779).
Eronia Valeria, Doubleday, *Diurnal Lep.* p. 65, n. 1.
Mancipium Vorax Valeria, Hübner, *Samml. Exot. Schmett.*
p. (1806-27).
Acrea Valeria, Hübner, *Verz. bek. Schmett.* p. 93.
Pieris Valeria, Godart, *Enc. Méth.* IX. p. 154. *Boisduval,*
Spéc. Gén. Lép. I. p. 444. *Doubleday, List Lep.*
Brit. Mus. pt. I. p. 25.
Pontia (Valeria) Valeria, Horsfield, *Cat. Lep. Mus.*
E.I.C. p. 139.
Pieris Hippia, Fabricius, *Ent. Syst.* III. *pt. I.* p. 59
(1793).
Danais Hippia, Godart, *Enc. Méth.* IX. p. 193.
♀ *Danais Anais*, Bougainville, *Voy. de la Thétis*, t. 44, f. 2
(1837).
♀ *Pieris Indica*, E. Doubleday, *List Lep. Brit. Mus. pt. I.*
p. 25 (1844).
Var. ♀ *Papilio Philomela*, Fabricius, *Ent. Syst.* III. *pt. I.* p. 57.
a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
c. ♂. Borneo. Presented by H. Lowe, Esq.
d. e. ♂ ♀. Dukhun. Presented by Colonel Sykes.
f. ♀. N. India. From the Asiatic Society of Bengal.
g. h. ♀. N. India. Presented by Colonel Buckley.

107. *ERONIA AVATAR*, Moore (Plate IIa, fig. 1).

E. Avatar, n. sp.—Male, *UPPER-SIDE* greenish or bluish-white; *fore-wings*, with the anterior and exterior margins, and the veins, black; *hind-wings* along the exterior margin slightly undulated with black, which extends along the veinlets. *UNDER-SIDE* mother-o'-pearl-white, with the veinlets blackish.

- a. b. c. d.* ♂. N. India. Presented by Lieutenant Hugo James.
e. f. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

There appears to be no variation whatever in this species, eighteen specimens before me being all precisely alike.

Genus COLIAS, *Fabricius*.

COLIAS, *Fabricius* (*Illiger, Mag.* VI. p. 284, 1807). *Boisduval*,
Spéc. Gén. Lép. I. p. 633. *Doubleday, Diurnal Lep.*

COLOTIS, *ZERENE*, *Hübner, Verz.* (1816).

EUBYMMUS (*Swainson*). *Horsfield, Cat. Lep. Mus. E.I.C.* pp. 129, 134
 (1829); *Swains., Zool. Ill. 2nd ser. fol. 60* (1832).

108. *COLIAS EDUSA*, *Fabricius*.

Papilio Edusa, *Fabricius, Ent. Syst. III. pt. I.* p. 206
 (1793). *Hübner, Europ. Schmett. Pap. f.* 429–31.

Colias Edusa, *Ochsenheimer, Schmett. von Eur.* IV. pt. II.
 p. 32. *Godart, Enc. Méth. IX.* p. 101. *Boisduval*,
Spéc. Gén. Lép. I. p. 638. *E. Doubleday, List Lep.*
Brit. Mus. pt. I. p. 41; id. *Diurnal Lep.* p. 74, n. 7.

a. b. ♂. Bootan. From Pemberton's Collection.

c. d. e. f. ♂ ♀. N. India. Presented by Colonel
 Buckley.

g. ♀. N. India. Presented by Colonel Hearsey.

h. i. ♀. Darjeeling. From Pearson's Collection.

109. *COLIAS HYALE*, *Linnæus Sp.*

Papilio Hyale, *Linnæus, Syst. Nat.* II. p. 764 (1767).
Fabricius, Ent. Syst. III. pt. I. p. 207.

Euryalus Hyale, *Swainson. Horsfield, Cat. Lep. Mus.*
E.I.C. p. 134 (1829).

Pieris Hyale, *Latreille, Gén. Crust. et Ins.* XIV. p. 113
 (1805).

Colias Hyale, *Ochsenheimer, Schmett. von Europa*, IV.
 p. 32 (1816). *Godart, Enc. Méth. IX.* p. 33.
Boisduval, Spéc. Gén. Lép. I. p. 650. *Doubleday,*
List Lep. Brit. Mus. pt. I. p. 147; id. *Diurnal*
Lep. p. 75, n. 23.

Papilio Palæo, *Hübner, Eur. Schmett. Pap. f.* 438, 439.

a. b. c. ♂ ♀. Punjaub. Presented by Colonel
 Hearsey.

110. *COLIAS NERIENE*, *Fischer*.

Colias Neriene, *Fischer, Entomol. de la Russie* Lép. t. 11,
 f. 3, 4 (1820–22). *Boisduval, Spéc. Gén. Lép.* I.

p. 646. *Herr. Schaeffer*, f. 30, 31, 32 (1844). *Doubleday, List Lep. Brit. Mus. pt. I.* p. 147; id. *Diurnal Lep.* p. 74, n. 18.

- a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.
- d. e. f. g. ♂ ♀. Punjab. Presented by Colonel Hearsey.

Genus TERIAS, Swainson.

TERIAS, Swainson, Zool. Ill. 1st ser. text to t. 22 (1820). Horsfield.

XANTHIDIA, Boisduval et Leconte, Icon. Lép. Am. Sept. p. 48 (1828).

LEPTOSIA, EUREMA, ABAEIS, Hübner, Verz. (1816).

III. TERIAS HECABE, Linnaeus Sp.

Papilio Hecabe, Linnæus, Syst. Nat. II. p. 763 (1767).
Cramer, Pap. Exot. II. t. 124, f. B.C. *Fabricius, Spec. Ins.* II. p. 42; *Ent. Syst.* III. pt. I. p. 192.
Sulz., Ins. edit. Roem. t. 15, f. 7.

Terias Hecabe, Swainson, Zool. Ill. 1st ser. t. 22. Horsfield, Cat. Lep. Mus. E.I.C. p. 135. Boisduval, Spéc. Gén. Lép. I. p. 669. Doubleday, List Lep. Brit. Mus. pt. I. p. 44; id. Diurn. Lep. p. 79, n. 29.

Pieris Hecabe, Godart, Enc. Méth. IX. p. 134.

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
- c. d. ♂. N. India. From the Asiatic Society of Bengal.
- d. ♀. Bootan. From Pemberton's Collection.
- e. ♂. Darjeeling. From Pearson's Collection.
- f. ♀. Canara (S. India). Presented by S. N. Ward, Esq.

"This is one of the most common species of diurnal Lepidoptera, having a wide range through intratropical Asia and the Eastern Islands. In Java, it was frequently obtained by breeding, and the larva feeds on the *Aeschynomene Sesban*, and is found abundantly from January to April."—(Horsfield.)

The larva and pupa of *T. Hecabe* are figured on Plate 1, fig. 11, 11a; from Java.

112. *TERIAS SARI*, Horsfield.

Terias Sari, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 136 (1829).

Terias Hecabe, var. ♀, *Boisduval, Spéc. Gén. Lép. I.* p. 669. *Doubleday, Diurnal Lep.* p. 79, n. 29.

a. ♀. Java. From Dr. Horsfield's Collection.

b. c. Borneo. Presented by Hugh Lowe, Esq.

"Clearly distinguished from *T. Hecabe* by the breadth and interior outline of the posterior border of the fore-wings above, as well as by the large quadrangular apical spot and the peculiarity of the markings underneath."—(Horsfield.)

113. *TERIAS TILAHIA*, Horsfield.

Terias Tilaha, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 136 (1829). *Boisduval, Spéc. Gén. Lép. I.* p. 668. *E. Doubleday, Diurnal Lep.* p. 79, n. 28; id. *List Lep. Brit. Mus. App.* p. 11.

a. ♀. Java. From Dr. Horsfield's Collection.

114. *TERIAS BLANDA*, Boisduval.

Terias Blanda, *Boisduval, Spéc. Gén. Lép. I.* p. 672 (1836). *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 45; id. *Diurnal Lep.* p. 79, n. 35.

a. ♂. China. Presented by the Trustees of the British Museum.

b. ♂. Darjeeling. From the Indian Collection, Exposition Universelle at Paris, 1855.

115. *TERIAS HARINA*, Horsfield.

Terias Harina, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 137 (1829). *Boisduval, Spéc. Gén. Lep. I.* p. 668. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 44; id. *Diurnal Lep.* p. 79, n. 27.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. d. e. ♂. Darjeeling. From the Indian Collection, Exposition Universelle at Paris, 1855.

116. *TERIAS LÆTA*, Boisduval.

Terias Læta, Boisduval, Spéc. Gén. Lép. I. p. 674 (1836).

E. Doubleday, List Lep. Brit. Mus. pt. I. p. 45; id. Diurnal Lep. p. 79, n. 39.

- a. b. ♂ ♀. Bootan. From Pemberton's Collection.
- c. d. ♂ ♀. N. India. Presented by Colonel Buckley.
- e. ♂. N. India. Presented by Colonel Hearsey.
- f. g. Darjeeling. From the Indian Collection, Exposition Universelle at Paris, 1855.

117. *TERIAS VENATA*, Moore (Plate IIa., fig. 2).

T. Venata, n. sp.—Intermediate in shape between *T. Læta* and *T. Drona*, having the fore-wings less sharply produced at the anterior angle than in *T. Læta*, yet more so than in *T. Drona*; markings on the UPPER-SIDE of the fore-wings as in *T. Læta*, but the brown border extending to the posterior angle; hind-wings narrowly bordered with brown; UNDER-SIDE clear yellow, with only a disco-cellular dot, and one between each veinlet on the outer margin of both wings. Expanse of wings $1\frac{1}{4}$ in.

- a. b. c. N. India. Presented by Colonel Hearsey.

118. *TERIAS DRONA*, Horsfield.

Terias Drona, Horsfield, Cat. Lep. Mus. E.I.C. p. 137, t. 1, f. 13 (1829). Boisduval, Spéc. Gén. Lép. I. p. 675. E. Doubleday, Diurnal Lep. p. 79, n. 40.

- a. ♂. Java. From Dr. Horsfield's Collection.
- b. c. d. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus THESTIAS, Boisduval.

Thestias, Boisduval, Spéc. Gén. Lép. I. p. 590 (1836).

Ixiias, Hübner, Verz. bek. Schmett. p. 95 (1816).*

Pieris, p. Latreille, Godart.

Pontia, p. Horsfield.

* This name being so near that of *Ixia*, employed in Botany, cannot be retained.

119. *THESTIAS ÆNIPPE*, Cramer Sp.

- Papilio Ænippe, Cramer, Pap. II. t. 105, f. C.D. t. 229, f. B.C. (1779). Fabricius, Spec. Ins. II. p. 48; Ent. Syst. III. pt. I. p. 204.
 Thestias Ænippe, Boisduval, Spéc. Gén. Lép. I. p. 591. Doubleday, List Lep. Brit. Mus. pt. I. p. 36; id. Diurnal Lep. p. 61, n. 1.
 Ixias Ænippe, Hübner, Verz. bek. Schmett. p. 95.
 Pieris Ænippe, Godart, Enc. Méth. IX. p. 120.
 Pontia Ænippe, Horsfield, Cat. Lep. Mus. E.I.C. p. 143.
a. b. c. ♂ ♀. Bootan. From Pemberton's Collection.
d. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.
f. g. ♀ var. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

120. *THESTIAS MARIANNE*, Cramer Sp.

- Papilio Marianne, Cramer, Pap. III. t. 217, f. C.D.E. (1780).
 Thestias Marianne, Boisduval, Spéc. Gén. Lép. I. p. 592. Doubleday, List Lep. Brit. Mus. pt. I. p. 36; id. Diurnal Lep. p. 61, n. 2.
 Ixias Marianne, Hübner, Verz. bek. Schmett. p. 95.
 Pieris Marianne, Godart, Enc. Méth. IX. p. 120.
 Pontia Marianne, Horsfield, Cat. Lep. Mus. E.I.C. p. 142.
 Papilio Sesia, Fabricius, Spec. Ins. II. p. 47 (1781).
 Ixias Bebryce, Hübner, Verz. bek. Schmett. p. 95 (1816).
a. b. ♂ ♀. N. India. Presented by Colonel Hearsey.
c. ♂. N. India. Presented by Colonel Buckley.
d. e. ♂ ♀. Madras. From Captain J. M. Jones's Collection, 1856.

121. *THESTIAS PYRENE*, Linnæus Sp.

- Papilio Pyrene, Linnæus, Syst. Nat. II. p. 762 (1767). Cramer, Pap. II. t. 125, f. A.B.C.
 Thestias Pyrene, Boisduval, Spéc. Gén. Lép. I. p. 593. Doubleday, List Lep. Brit. Mus. pt. I. p. 36; id. Diurnal Lep. p. 61, n. 3.
 Ixias Pyrene, Hübner, Verz. bek. Schmett. p. 95.

Pieris Pyrene, *Godart, Enc. Méth.* IX. p. 120.
 Pontia Pyrene, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 143.
 Papilio Rhexia, *Fabricius, Mant. Ins.* II. p. 23 (1787).
 Ixias Anexibia, *Hübner, Verz. bek. Schmett.* p. 95 (1816).

- a. ♂. Silhet. Presented by Colonel Buckley.
- b. ♀. Assam. Donor unknown.
- c. ♀. Darjeeling. From Pearson's Collection.
- d. e. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

122. THESTIAS VENILIA, *Godart Sp.*

Pieris Venilia, *Godart, Enc. Méth.* IX. p. 121 (1819).
 Thestia Venilia, *Boisduval, Spéc. Gén. Lép.* I. p. 594.
Doubleday, List Lep. Brit. Mus. pt. I. p. 36; id.
Diurnal Lep. p. 61, n. 5.

Pontia Venilia, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 143.
 ♀ Papilio Ænippe, *apud Cramer*, II. t. 157, f. C. D.
 ♂ Thestias Balidia, *Boisduval, MS. Mus. E.I.C.*

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

Genus IDMAIS, *Boisduval.*

IDMAIS, *Boisduval, Spéc. Gén. Lép.* I. p. 59 (1836).
 MANCIPIUM, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 141 (1829).

123. IDMAIS CALAIS, *Cramer Sp.*

♂ ♀ Papilio Calais, *Cramer, Pap.* I. t. 53, f. C. D. (1779);
 t. 351, f. A—D.

Idmais Calais, *Boisduval, Spéc. Gén. Lép.* I. p. 587.
Doubleday, List Lep. Brit. Mus. pt. I. p. 36; id.
Diurnal Lep. p. 59, n. 4.

♂ Papilio Amata, *Fabricius, Ent. Syst.* III. pt. I. p. 202
 (1793).

Pieris Amata, *Godart, Enc. Méth.* IX. p. 131.

Pontia (Mancipium) Amata, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 141.

♀ Papilio Ciprea, *Fabricius, Ent. Syst.* III. pt. I. p. 202.

- a. b. c. d. India. Donor unknown.

e. f. Madras. From Captain J. M. Jones's Collection.

124. *IDMAIS PHISADIA*, Godart Sp.

- Pieris Phisadia, *Godart*, *Enc. Méth.* IX. p. 132 (1819).
 Idmais Phisadia, *Boisduval*, *Spéc. Gén. Lép.* I. p. 587.
Doubleday, Diurnal Lep. p. 59, No. 3.
 Pontia Arne, *Klug*, *Ehrenb. Symb. Phys. Ins.* t. 7, f. 1—4
 (1829—45).
a. b. c. d. N. India. Donor unknown.

125. *IDMAIS DYNAMENE*, Klug Sp.

- Pontia Dynamene, *Klug*, *Ehrenb. Symb. Phys. Ins.* t. 7,
 f. 5, 6 (1829—45).
 Idmais Dynamene, *Boisduval*, *Spéc. Gén. Lép.* I. p. 588.
Doubleday, Diurnal Lep. p. 59, n. 5.

a. b. c. Punjab. Presented by Colonel Hearsey.

126. *IDMAIS FAUSTA*, Olivier Sp.

- Pieris Fausta, *Oliv.*, *Voy. en Syrie*, t. 33, f. 4 (1801—7).
 Idmais Fausta, *Boisduval*, *Spéc. Gén. Lép.* I. p. 586.
Doubleday, Diurnal Lep. p. 59, n. 2.
 Pieris Fausta, *Godart*, *Enc. Méth.* IX. p. 132 (1819).
a. ♂. Punjab. Presented by Colonel Hearsey.
b. ♂. N. W. India. Donor unknown.

Genus *CALLOSUNE*, E. *Doubleday*.

- ANTHOCHARIS* (*CALLOSUNE*), *Doubleday, Diurnal Lep.* p. 57 (1847).
EUCHLOË, *APHRODITE*, p. *Hübner*.
MANCIPIUM, p. *Horsfield*.

127. *CALLOSUNE EUCHARIS*, Fabricius Sp.

- Papilio Eucharis, *Fabricius*, *Syst. Ent.* p. 472 (1775).
 Anthocharis (*Callosune*) Eucharis, *Doubleday, Diurnal Lep.* p. 57, n. 15.
 Pieris Eucharis, *Godart*, *Enc. Méth.* IX. p. 124.
 Anthocharis Eucharis, *Boisduval*, *Spéc. Gén. Lép.* I.
 p. 568. *Doubleday, List Lep. Brit. Mus. pt. I.* p. 35.
 Papilio Aurora, *Cramer*, *Pap.* III. t. 299, f. A—D.
 (1782).
 Pieris Titea, *Godart*, *Enc. Méth.* IX. p. 124 (1819).

Pontia (Mancipium) Titea, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 141.

Euchloë Coëneos, *Hübner, Verz. bek. Schmett.* p. 94 (1816).

a. b. India. Donor unknown.

c. Madras. Donor unknown.

d. Madras. From Captain J. M. Jones's Collection, 1856.

128. *CALLOSUNE DANAE*, *Fabricius Sp.*

Papilio Danaë, *Fabrieius, Mant. Ins. II.* p. 22; *Syst. Ent.* p. 476 (1775). *Donovan, Ins. of Ind. t.*

Anthocharis (*Callosune*) *Danaë*, *Doubleday, Diurnal Lep.* p. 57, n. 17.

Pieris Danaë, *Godart, Enc. Méth. IX.* p. 124.

Pontia (*Mancipium*) *Danaë*, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 141.

Anthocharis Danaë, *Boisduval, Spéc. Gén. I.* p. 570. *Doubleday, List Lep. Brit. Mus. pt. I.* p. 35.

Papilio Eborea, *Cramer, Pap. IV.* t. 352, f. C—F. (1782).

Aphrodite Eborea, *Hübner, Verz. bek. Schmett.* p. 94.

a. b. c. ♂ ♀. India. Donor unknown.

d. ♀. Madras. From Captain J. M. Jones's Collection, 1856.

129. *CALLOSUNE ETRIDA*, *Boisduval Sp.*

Anthocharis Etrida, *Boisduval, Spéc. Gén. Lép. I.* p. 576 (1836).

Anthocharis (*Callosune*) *Etrida*, *Doubleday, Diurnal Lep.* p. 57, n. 25.

a. b. c. d. e. f. ♂ ♀. N. India. Presented by Colonel Hearsey.

Genus PONTIA, *Fabricius.*

PONTIA, *Fabricius*. *McLeay. Boisduval, Spéc. Gén. Lép. I.* p. 430 (1836).

NINA, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 140 (1829).

130. *PONTIA NINA*, *Fabricius.*

Papilio Nina, *Fabricius, Ent. Syst. III. pt. I.* p. 194 (1793).

- Pontia Nina, *Boisduval*, *Spéc. Gén. Lép.* I. p. 431.
Doubleday, *List Lep. Brit. Mus. pt. I.* p. 24;
Diurnal Lep. p. 40, n. 2.
- Pieris Nina, *Godart*, *Enc. Méth. IX.* p. 162.
- Pontia (Nina) Nina, *Horsfield*, *Cat. Lep. Mus. E.I.C.*
p. 140.
- Papilio Xiphia, *Fabricius*, *Mant. Ins. II.* p. 20 (1787).
- Leptosia chlorographa (var.), *Hübner*, *Zuträge*, n. 47, 48
(1818).
- a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.
- d. ♂. Canara. Presented by S. N. Ward, Esq.
- e. ♀. N. India. Presented by Colonel Buckley.
- f. ♂. N. India. Presented by Colonel Hearsey.

Genus PIERIS, Schrank.

- PIERIS, C. p. *Schrank*, *F. B. II. I.* p. 164 (1801). *Boisduval*, *Spéc. Gén. Lép.* I. p. 434. *Doubleday*, *Diurnal Lep.* p. 42.
- MYLOTHRIS, APPIAS, PERRYHYBRIS, DELIAS, CATHÆMIA, PONTIA, BELENOIS, ACRAEA, ANAPHEA, CATOPHAGA, SYNCLOË, *Hübner*, *Verz.* (1816).
- LEUCONIA, *Donzel*, *Ann. Soc. France*, VI. (1837).
- PIERIS, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 127 (1829).
- PONTIA, p. *Horsfield*.

131. PIERIS NERO, *Fabricius Sp.*

- ♂ Papilio Nero, *Fabricius*, *Ent. Syst. III. pt. I.* p. 153 (1793). *Donovan*, *Ins. of India*, fig.
- Pieris Nero, *Boisduval*, *Spéc. Gén. Lép.* I. p. 485. *Doubleday*, *Cat. Lep. Brit. Mus.* p. 28; *Diurnal Lep.* p. 51, n. 155.
- ♀ Pieris Thyria, *Godart*, *Enc. Méth. IX.* p. 147 (1819). *Guérin*, *Icon. Rég. An. Ins. t. 77, f. 1.*
- Pontia Thyria, *Horsfield*, *Zool. Journ. IV.* p. 69, t. 4, f. 2 (1829).
- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
- c. d. Presented by T. Tatum, Esq.

132. PIERIS ROUXI, Boisduval.

Pieris Rouxii, Boisduval, Spéc. Gén. Lép. I. p. 481 (1836). Doubleday, Diurnal Lep. p. 50, n. 148.

Pieris Namouna, Doubleday, List Lep. Brit. Mus. pt. I. p. 281 (1844).

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. e. f. ♂ ♀. Borneo. Presented by H. Lowe, Esq.

g. Bootan. From Pemberton's Collection.

h. Madras. From Wight's Collection.

133. PIERIS PANDA, Godart Sp.

Pieris Panda, Godart, Enc. Méth. IX. p. 147 (1819).

Boisduval, Spéc. Gén. Lép. I. p. 485. Doubleday, List Lep. Brit. Mus. pt. I. p. 28; Diurnal Lep. p. 50, n. 154.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

134. PIERIS HIPPO, Cramer Sp.

♀ Papilio Hippo, Cramer, Pap. III. t. 195, f. B.C. (1782).

Pieris Hippo, Godart, Enc. Méth. IX. p. 143. Boisduval, Spéc. Gén. Lép. I. p. 534. Doubleday, List Lep. Brit. Mus. pt. I. p. 32.

♂ Papilio Phryne, Fabricius, Ent. Syst. III. pt. I. p. 196 (1793).

Pieris Phryne, Doubleday, List Lep. Brit. Mus. pt. I. p. 28; Diurnal Lep. p. 50, n. 149.

Pieris Eleonora, Boisduval, Spéc. Gén. Lép. I. p. 481 (1836).

? ♂ Papilio Lyncida, Cramer, Pap. II. t. 131, f. B. (1779).

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

f. ♂. Bootan. From Pemberton's Collection.

g. h. ♂ ♀. Silhet. Presented by Colonel Buckley.

i. j. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

135. *PIERIS NERISSA*, *Fabricius Sp.*

Papilio Nerissa, *Fabricius*, *Ent. Syst.* III. pt. I. p. 192 (1793).

Pieris Nerissa, *Godart*, *Enc. Méth.* IX. p. 142. *Boisduval*, *Spéc. Gén. Lép.* I. p. 535. *Doubleday, Diurnal Lep.* p. 50, n. 142.

Aeræa Nerissa, *Hübner*, *Verz. bek. Schmett.* p. 93.

a. ♂. Assam. Donor unknown.

136. *PIERIS EGA*, *Boisduval*.

Pieris Ega, *Boisduval*, *Spéc. Gén. Lép.* I. p. 536 (1836). *Doubleday, List Lep. Brit. Mus.* pt. I. p. 32.

a. b. Canara. Presented by S. N. Ward, Esq.

c. N. India. Donor unknown.

137. *PIERIS NEOMBO*, *Boisduval* (*Plate IIa.*, fig. 3, ♀).

Pieris Neombo, *Boisduval*, *Spéc. Gén. Lép.* I. p. 539 (1839). *Doubleday, Diurnal Lep.* p. 50, n. 135.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

138. *PIERIS MESENTINA*, *Cramer Sp.*

Papilio Mesentina, *Cramer*, *Pap.* III. t. 270, f. A.B. (1782).

Pieris Mesentina, *Godart*, *Enc. Méth.* IX. p. 130. *Boisduval*, *Spéc. Gén. Lép.* I. p. 501. *Doubleday, List Lep. Brit. Mus.* pt. I. p. 30; *Diurnal Lep.* p. 46, n. 50.

Pontia Mesentina, *Horsfield*, *Cat. Lep. Mus. E. I. C.* p. 142.

Papilio Aurota, *Fabricius*, *Ent. Syst.* III. pt. I. p. 197 (1793).

a. b. ♂ ♀. N. India. From the Asiatic Society of Bengal.

c. d. e. f. ♂ ♀. N. India. Presented by Colonel Hearsey.

g. ♀, with black borders. Madras. From Captain J. M. Jones's Collection, 1856.

139. *PIERIS LIBYTHEA*, Fabricius Sp.

Papilio Libythea, *Fabricius*, *Ent. Syst.* III. pt. I. p. 190 (1793). *Donovan*, *Ins. Ind. t.*

Pieris Libythea, *Doubleday*, *Diurnal Lep.* p. 46, n. 44.

Pieris Libitina, *Godart*, *Enc. Méth.* IX. p. 133 (1819). *Boisduval*, *Spéc. Gén. Lép.* I. p. 499.

? *Papilio Zelmira*, *Cramer*, *Pap.* IV. t. 320, f. C—F. (1782).

a. b. ♂ ♀. India. Donor unknown.

c. ♂. N. India. Presented by Colonel Hearsey.

140. *PIERIS PANDIONE*, Hübner Sp.

Hiposcritia Pandione, *Hübner*, *Zuträge*, f. 651—2 (1826).

Pieris Pandione, *Boisduval*, *Spéc. Gén. Lép.* I. p. 537. *Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 32; *Diurnal Lep.* p. 50, n. 136.

a. ♂. Java. From Dr. Horsfield's Collection.

b. c. d. ♂ ♀. N. India (?). Purchased.

141. *PIERIS PAULINA*, Cramer Sp.

Papilio Paulina, *Cramer*, *Pap.* II. t. 110, f. E.F. (1779). *Fabricius*, *Ent. Syst.* III. pt. I. p. 189.

Pieris Paulina, *Godart*, *Enc. Méth.* IX. p. 142. *Boisduval*, *Spéc. Gén. Lép.* I. p. 538. *Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 33; *Diurnal Lep.* p. 50, n. 137.

Catophaga Leis, *Hübner*, *Zuträge*, f. 771, 772 (1827).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Borneo. Presented by H. Lowe, Esq.

f. g. ♂ ♀. N. India. Purchased.

h. i. ♂ ♀. Darjeeling. From the Messrs. Schlagintweit's Collection.

142. *PIERIS DURVASA*, Moore.

Pieris Durvasa, *Moore*, *Proc. Zool. Soc.* (1857), p. ; *Annulosa*, t. , f. 6.

a. b. c. d. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

e. f. ♀. Assam. From McClelland's Collection.

Pieris Durvasa, n. sp.—UPPER-SIDE white; male, *fore-wing*, from middle of costal margin, curving transversely, apically, and scolloped to near end of outer margin, black, and having near the apex some white marks, generally three, the outer ones being sometimes indistinct; base of costal margin and body greenish; on the middle of disco-cellular veinlets is a round black spot, and another more quadrate spot between the first and second median veinlets, the latter spot slightly touching the scolloped black outer margin. Female, black colour broader; the quadrate spot larger and broadly confluent at the outer angle's angles with the outer margin, thus forming a white spot on middle of outer margin; *hind-wing* of female with a marginal row of blackish spots, the extreme margin and anal angle being yellowish-white. UNDER-SIDE, apex of *fore-wing* pale yellow, the black forming only a curved transverse bar, besides the two black spots; *hind-wing* wholly pale yellow, and having a small disco-cellular black spot; body yellowish. Form of wings as in *P. Paulina* and *P. Pandione*. Expanse of wings $2\frac{1}{4}$ in. to $3\frac{3}{8}$ in.

143. PIERIS INDRA, Moore.

Pieris Indra, Moore, Proc. Zool. Soc. (1857), p. ;
Annulosa, t. , f. 5.

a. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

Pieris Indra, n. sp.—UPPER-SIDE dark brown; *fore-wing* with a central longitudinal space of white from the base, also two small white spots near anterior angle; *hind-wing* with the anterior base brownish greenish-white, also with two white spots near anterior angle. UNDER-SIDE, *fore-wing* with a broad irregular fascia from middle of anterior margin to posterior angle; anterior angle chrome-yellow, with some white parting the two colours; basal half white, tinged with straw-yellow along discoidal cell; *hind-wing* chrome-yellow, minutely irrorated with brown; anterior half of discoidal cell and space between each veinlet near outer margin white, also a dark brown dot on disco-cellular veinlet. Wings shaped as in *P. Paulina*, Fabr. Expanse of wings 3 in.

Remark.—The nearest ally of *Pieris Indra* appears to be *P. Lalage*, Doubleday, Diurnal Lep. t. VI. fig. 5.

144. *PIERIS NIPALENSIS*, G. R. Gray.

Pieris Nipalensis, G. R. Gray, *Lep. Ins. Nepal*, p. 9, t. 6, f. 1, 3 (1832).

Pieris Brassicæ, *Linnæus*, var. *E.* *Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 32.

a. ♂. Bootan. From Pemberton's Collection.

b. ♀. N. India. Presented by Colonel Hearsey.

c. ♂. Madras. From Wight's Collection.

145. *PIERIS GLICIRIA*, Cramer Sp.

Papilio Gliciria, Cramer, *Pap. Exot. II.* t. 171, f. E. F. (1779).

Pieris Gliciria, *Boisduval*, *Spéc. Gén. Lép. I.* p. 524. *Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 32; *Diurnal Lep.* p. 48, n. 102.

Catophaga Gliciria, *Hübner*, *Verz. bek. Schmett.* p. 93.

Pieris Glaphyra, *Godart*, *Enc. Méth. IX.* p. 160 (1819).

a. ♂. Assam. From McClelland's Collection.

b. ♀. Bootan. From Pemberton's Collection.

c. ♀. N. India. Presented by Colonel Buckley.

d. ♀. Chusan. Presented by Dr. Cantor, 1844.

146. *PIERIS DAPLIDICE*, Linnæus Sp.

Papilio Daplidice, *Linnæus*, *Syst. Nat. II.* p. 760 (1767).

Fabricius, *Ent. Syst. III. I.* p. 191. *Hübner*, *Eur. Schmett. Pap. f. 414, 415.*

Pieris Daplidice, *Godart*, *Enc. Méth. IX.* p. 128. *Boisduval*, *Spéc. Gén. Lép. I.* p. 544. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 33; id. *Diurnal Lep.* p. 48, No. 95.

Synchloë Daplidice, *Hübner*, *Verz. bek. Schmett.* p. 94.

a. ♀. N. India. Presented by Colonel Hearsey.

Several other specimens of *P. Daplidice* are in the collections of Colonel Hearsey and J. O. Westwood, Esq.

147. *PIERIS REMBA*, Moore.

Pieris Remba, n. sp.—UPPER-SIDE pale yellowish-white; anterior half of fore-wing blackish-brown; some blackish on outer margin of

hind-wing. UNDER-SIDE, broadly along costal margin and anterior half of fore-wing dull greenish-yellow, smeared with blackish; rest white; hind-wing dull greenish-yellow, smeared with darker colour. Shape of wings as in *P. Coronis*, Fabr. Expanse of wings $2\frac{2}{5}$ in.

a. Canara, Malabar Coast. Presented by S. N. Ward, Esq.

148. *PIERIS NAMA*, *Doubleday*.

δ Pieris Nama, *Doubleday, List Lep. Brit. Mus. pt. I.* p. 28 (1844); *Diurnal Lep.* p. 46, n. 38. *Moore, Proc. Zool. Soc.* (1857); *Annulosa, t. , f. 1, 2,* δ φ .

a. δ . Silhet. Presented by the Trustees of the British Museum.

b. δ . Darjeeling. From Pearson's Collection.

c. δ . Bootan. From Pemberton's Collection.

d. δ . e. f. g. φ . Darjeeling. From Messrs. Schlagintweit's Collection.

Pieris Nama.—Male, UPPER-SIDE white; fore-wing with a narrow brown line along anterior margin, curving and widening across near the middle of wing, and again tapering to posterior angle; hind-wing tinged with blackish (as if from intensity of that colour on the under-side) along outer margin, where the veinlets are dark brown. Female, brown, with three longitudinal white streaks in middle of fore-wing, and two in the hind-wing, these streaks in some specimens being confluent, and occupying nearly the whole of the middle of both fore and hind-wings. UNDER-SIDE, along anterior, and widening to the outer margin of fore-wing, greenish-yellow, the rest white; hind-wing greenish yellow, darker on the veins, and nearly white along discoidal cell towards anterior angle. Expanse of wings from $2\frac{1}{4}$ to 3 in.

149. *PIERIS CORONIS*, *Cramer Sp.*

Papilio Coronis, *Cramer, Pap. I. t. 44, f. B.* (1779).

Fabricius, Spec. Ins. II. p. 45; Ent. Syst. III. pt. I. p. 198.

Pieris Coronis, *Godart, Enc. Méth. IX. p. 132. Boisduval, Spéc. Gén. Lép. I. p. 471. Doubleday, List Lep. Brit. Mus. pt. I. p. 27; Diurnal Lep. p. 46, n. 35.*

Pontia Coronis, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 144, t. 4, f. 9, *larva*,—9a, *pupa*.

Var. *Papilio Evagete*, *Cramer, Pap. III.* t. 221, f. F.G. (1782).

Var. *Papilio Zeuzippe*, *Cramer, Pap. IV.* t. 362, f. E.F. (1782).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♀. Dukhun. Presented by Colonel Sykes.

f. g. ♂ ♀. N. India. From the Asiatic Society of Bengal.

The larva and pupa of *P. Coronis* are figured on Plate 1, fig. 12, 12a, from Java.

150. PIERIS JUDITH, *Fabricius Sp.*

Papilio Judith, *Fabricius, Mant. Ins. II.* p. 22 (1787); *Ent. Syst. III. pt. I.* p. 202. *Donovan, Ins. Ind. t.*

Pieris Judith, *Godart, Enc. Méth. IX.* p. 121. *Boisduval, Spéc. Gén. Lép. I.* p. 468. *Doubleday, List Lep. Brit. Mus. pt. I.* p. 27; *Diurnal Lep.* p. 45, n. 30.

Pontia Judith, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 144.

Acræa Judith, *Hübner, Zuträge*, f. 669, 670.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♀. Pinang. Presented by Dr. Cantor, 1844.

"This species is very abundant in Java."—(Dr. Horsfield.)

151. PIERIS LEA, E. *Doubleday*.

Pieris Lea, *Doubleday, List Lep. Brit. Mus. pt. I.* p. 27 (1844); *Ann. N. H. XVII.* p. 23 (1846); *Diurnal Lep.* p. 45, n. 31.

Pieris Clemanthe, *Doubleday, Diurnal Lep.* t. 6, f. 3 (1847).

a. ♂. Borneo. Presented by the Trustees of the British Museum.

b. c. d. ♂ ♀. Borneo. Presented by Hugh Lowe, Esq.

PIERIS, Horsfield.

152. PIERIS THESTYLIS, E. Doubleday.

Pieris Thestylis, E. Doubleday, *Gray's Zool. Misc.* p. 76 (1842); *List Lep. Brit. Mus.* pt. I. p. 25; *Diurnal Lep.* p. 44, n. 1.

- a. b. ♂ ♀. Assam. Presented by Colonel Buckley.
- c. ♀. Bootan. From Pemberton's Collection.
- d. e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

153. PIERIS SETA, Moore. ♀. ♀ *Thestylis*.

Pieris Seta, Moore, *Proc. Zool. Soc.* (1857), p. ; *Annulosa*, t. 44, f. 3.

- a. ♀. Bootan. From Pemberton's Collection.

Pieris Seta, n. sp.—UPPER-SIDE blackish-brown; fore-wing with two rows of narrowish white marks, two lengthened marks between median and submedian veinlets, and four small spots within discoidal cell; hind-wing with a marginal row of whitish spots; another row from anterior margin widening towards the anal angle; abdominal margin broadly whitish, the latter tinged with yellow; also a white linear mark in discoidal cell. UNDER-SIDE as in the upper-side, but with all the markings on the hind-wings yellow. Wings shaped as in *P. Thestylis*, Doubleday. Expanse $3\frac{7}{8}$ in.

154. PIERIS BELLADONNA, Fabricius Sp.

Papilio Belladonna, Fabricius, *Ent. Syst.* III. pt. I. p. 180 (1793). *Donovan, Nat. Rep.* t. 35 (1823).

Pieris Belladonna, Boisduval, *Spéc. Gén. Lép.* I. p. 447. *Doubleday, List Lep. Brit. Mus.* pt. I. p. 25; *Diurnal Lep.* p. 44, n. 2.

Pieris Horsfieldii, G. R. Gray, *Gray's Zool. Misc.* p. 32 (1832); *Lep. Ins. Nepal*, p. 7, t. 8, f. 2. *Boisduval, Spéc. Gén. Lép.* I. p. 448. *Herr. Schaeffer, Pap. Exot. liv.* I. f. 13, 14 (1853).

- a. b. c. ♂ ♀. Darjeeling. From Pearson's Collection.
- d. ♂. N. India. Presented by Colonel Buckley.
- e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

155. *PIERIS SANACA*, Moore.

Pieris Sanaca, *Moore*, *Proc. Zool. Soc.* (1857), *p.* ;
Annulosa, *t.* , *f.* 4.

a. b. Darjeeling Presented by Colonel Buckley.

Pieris Sanaca, *n. sp.*—UPPER-SIDE white; *fore-wing* with the veins and veinlets broadly clouded with black, leaving only a row of lanceolate white spots on the outer margin, and another row of more linear marks extending across the disc; *hind-wing* with the veins and veinlets sharply defined with black; discoidal and median veins clouded with black, the latter broadly so; also a marginal row of angular lunate marks; anterior base and anal angle bright yellow. UNDER-SIDE, *fore-wing* as in the upper-side, but the white markings more clearly defined, those near the anterior angle being yellowish; *hind-wing* with the dark colour broader, and the white spaces nearly covered with yellow. Shape of wings as in *P. Belladonna*. Expanse $3\frac{1}{4}$ in.

156. *PIERIS CRITHOË*, Boisduval.

Pieris Crithoë, *Boisduval*, in *Guérin et Perch, Gén. Lép.*
t. , *f.* (18—); *id. Spéc. Gén. Lép.* I. *p.* 450
 (1836). *Doubleday, List Lep. Brit. Mus.* pt. I.
App. p. 7; *Diurnal Lep.* *p. 44, n. 3.*

a. ♀. Java. From Dr. Horsfield's Collection.

157. *PIERIS EGIALEA*, Cramer Sp.

Papilio Egialea, *Cramer, Pap.* II. *t.* 189, *f. D.E.* (1779);
id. III. t. 253, f. E.F.

Pieris Egialea, *Boisduval, Spéc. Gén. Lép.* I. *p.* 450.
Doubleday, List Lep. Brit. Mus. pt. I. *App. p. 7*;
Diurnal Lep. *p. 44, n. 4.*

Delias Apriate, *Hübner, Verz. bek. Schmett.* *p. 91* (1816).
Pieris Pasithoë, var., *Godart, Enc. Méth.* IX. *p. 148.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

158. *PIERIS PASITHOË*, Linnæus Sp.

Papilio Pasithoë, *Linnæus, S. N.* II. *p. 755* (1767).
Fabricius, Ent. Syst. III. *pt. I. p. 179.*
Pieris Pasithoë, *Godart, Enc. Méth.* IX. *p. 148. Bois-*

duval, *Spéc. Gén. Lép.* I. p. 451. *Doubleday, List Lep. Brit. Mus. pt.* I. p. 26; *Diurnal Lep.* p. 44, n. 5.

Delias Pasithoë, *Hübner, Verz. bek. Schmett.* p. 91.

Papilio Porsenna, *Cramer, Pap.* I. t. 43, f. D.E. (1779);
id. IV. t. 352, f. A.B.

a. b. ♂ ♀. Dukhun. Presented by Colonel Sykes.

159. *PIERIS THISBE*, *Cramer Sp.*

Papilio Thisbe, *Cramer, Pap.* III. t. 233, f. C. (1782).

Pieris Thisbe, *Boisduval, Spéc. Gén. Lép.* I. p. 449.
Doubleday, List Lep. Brit. Mus. pt. I. p. 25;
Diurnal Lep. p. 44, n. 6. *G. R. Gray, Lep. Ins. of Nepal*, p. 8, t. 7, f. 1.

Pieris Acalis, *Godart, Enc. Méth.* IX. p. 148 (1819).

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.

160. *PIERIS EUCHARIS*, *Drury Sp.*

Papilio Eucharis, *Drury, Ins.* II. t. 10, f. 5, 6 (1773).

Cramer, Pap. III. t. 201, f. B.C.; t. 202, f. C.

Pieris Eucharis, *Doubleday, Diurnal Lep.* p. 44, n. 10.

Pontia Eucharis, *Hübner, Verz. bek. Schmett.* p. 92.

Papilio Hyparete, *Fabricius, Ent. Syst.* III. pt. I. p. 178 (1793), nec. *Linn.*

Pieris Epicharis, *Godart, Enc. Méth.* IX. p. 153. *Boisduval, Spéc. Gén. Lép.* I. p. 456. *Doubleday, List Lep. Brit. Mus. pt.* I. p. 26.

a. b. ♂ ♀. N. India. From the Asiatic Society of Bengal.

c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

The larva and pupa of *P. Eucharis* are figured on Plate 1, fig. 13, 13a, from Canara, Southern India, where it was discovered by S. N. Ward, Esq., E.I.C.C.S., and from whose drawing our figures were copied.

161. *PIERIS HYPERATE*, *Linnæus Sp.*

Papilio Hyparete, *Linn.*, *S. N.* II. p. 763 (1767).
Clerck, Icon. t. 38, f. 2, 3.

Pieris Hyparete, *Godart*, *Enc. Méth.* IX. p. 153. *Boisduval*, *Spéc. Gén. Lép.* I. p. 455. *Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 26; *Diurnal Lep.* p. 44, n. 9.

Pontia Hyparete, *Hübner*, *Verz. bek. Schmett.* p. 92.

Papilio Antonoë, *Cramer*, *Pap.* II. t. 187, f. C.D. (1779); id. t. 320, f. A.B. (nec. *Stoll*).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

162. PIERIS HIERTE, *Hübner Sp.*

Pontia Hierte, *Hübner*, *Verz. bek. Schmett.* p. 92 (1816); id. *Zuträge*, f. 77, 78.

Papilio Antonoë, *Stoll*, *Suppl. à Cramer Pap.* t. 33, f. 2, 2B. (1791), nec. *Cramer* (1779).

Pieris Antonoë, *Boisduval*, *Spéc. Gén. Lép.* I. p. 454. *Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 26; *Diurnal Lep.* p. 44, n. 8.

Pieris Hyparete, var., *Godart*, *Enc. Méth.* IX. p. 153.

a. b. c. ♂ ♀. N. India. Presented by Colonel Sykes.

163. PIERIS AGOSTINA, *Hewitson.*

Pieris Agostina, *Hewitson*, *Exot. Butt.* pt. III.; *Pieridæ*, t. 1, f. 1, 2 (1852).

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

P. Agostina.—UPPER-SIDE cream-colour; hind-wings more tinted than the fore-wings; *fore-wing* with the costal margin grey; the apex, to beyond the middle of the outer margin, black, marked with a row of five white spots; *hind-wing* with the border of the under-side seen through. UNDER-SIDE, *fore-wing* with all the veins broadly black; *hind-wing* of a rich orange-yellow, with a submarginal line of black, and marginal spots of the same colour. Expanse of wings $2\frac{8}{10}$ in.

164. PIERIS DESCOMBESI, *Boisduval.*

Pieris Descombesi, *Boisduval*, *Spéc. Gén. Lép.* I. p. 465 (1836). *Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 27; *Diurnal Lep.* p. 45, n. 19.

a. b. ♂ ♀. Silhet. Presented by the Trustees of the British Museum.

c. ♀. Bootan. From Pemberton's Collection.

d. e. ♂ ♀. N. India. Presented by Colonel Sykes.

✓ 165. *PIERIS BELISAMA*, Cramer Sp.

Papilio Belisama, Cramer, Pap. III. t. 258, f. A. B. C. D. (1782).

Pieris Belisama, Godart, Enc. Méth. IX. p. 147. Boisduval, Spéc. Gén. Lép. I. p. 464. Doubleday, List Lep. Brit. Mus. pt. I. p. 27; Diurnal Lep. p. 45, n. 18.

Cathæmia Belisama, Hübner, Verz. bek. Schmett. p. 92 (1816).

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *P. Belisama* are figured on Plate 1, fig. 14, 14a, from Java.

"The larva of this species feeds on a species of *Dioscorea*, and was very abundant during the rainy season, from December to February."
—(Horsfield, MS.)

166. *PIERIS AUTOTHISBE*, Hübner Sp. (Plate IIa., fig. 4).

Delias Autothisbe, Hübner, Samml. Exot. Schmett. p. (1806-27).

Pieris Autothisbe, Boisduval, Spéc. Gén. Lép. I. p. 452. Doubleday, List Lep. Brit. Mus. pt. I. p. 26; Diurnal Lep. p. 45, n. 29.

a. ♀. Java. Presented by the Trustees of the British Museum.

167. *PIERIS CORONEA*, Cramer Sp.

Papilio Coronea, Cramer, Pap. I. t. 68, f. B. C. (1779); id. t. 361, f. G. H. Fabricius, Ent. Syst. III. pt. I. p. 201.

Pieris Coronea, Godart, Enc. Méth. IX. p. 151. Boisduval, Spéc. Gén. Lép. I. p. 474. Doubleday, List Lep. Brit. Mus. pt. I. p. 28; Diurnal Lep. p. 46, n. 52.

Anaphæis Coronea, *Hübner*, *Verz. bek. Schmett.* p. 93 (1816).

♀ *Papilio Deiopeia*, *Donovan*, *Ins. of New Holl.* t. (1805).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

168. *PIERIS VISHNU*, *Moore* (Plate IIa., fig. 5). *3d.*

Pieris Vishnu, n. sp.—UPPER-SIDE, base of fore-wing white; anterior half black, with a short row of small white spots across anterior angle; hind-wing, basal two-thirds white, the rest black; body whitish. UNDER-SIDE, fore-wing as in the upper, but the base yellowish; hind-wing with the basal half fine yellow, anterior half black, with a marginal row of eight red lunate spots. Wings shaped as in *P. Coronea*, Cramer. Expanse of wings $2\frac{1}{4}$ in.

a. ♂. Java. From Dr. Horsfield's Collection.

Genus APORIA, *Hübner*.

APORIA, *Hübner*, *Verz. bek. Schmett.* (1816).

PIERIS, p. *Doubleday*, *Diurnal Lep.* *near* *Lamprospilus*

169. *APORIA AGATHON*, *G. R. Gray* Sp.

Pieris Agathon, *G. R. Gray*, *Gray's Zool. Misc.* p. 33 (1832); *List Lep. Ins. Nepal*, p. 8, t. 8, f. 1. *Boisduval*, *Spéc. Gén. Lép.* I. p. 447. *Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 25; *Diurnal Lep.* p. 48, n. 85.

Var. *Pieris Phryxe*, *Boisduval*, *Spéc. Gén. Lép.* I. p. 446 (1836). *Blanchard*, *Voy. de Jacquemont*, *Ins.* t. 2, f. 1 (1840).

a. b. ♂ ♀. Bootan. From Pemberton's Collection.

c. ♀. Cherra Poonjee. Presented by Colonel Buckley.

170. *APORIA SORACTA*, *Boisduval* Sp.

Pieris Soracta, *Boisduval*, *MS. Doubleday*, *Diurnal Lep.* p. 48, n. 86 (1847); *List Lep. Brit. Mus. pt. I. App.* p. 7.

a. India. Donor unknown.

b. Darjeeling. Reserved from Indian Collection, *Exposition Universelle* at Paris, 1855.

Aporia Soracta, Boisd.—UPPER-SIDE white as in *A. Cratægi*; fore-wing with the veinlets on the anterior half and outer margin broadly clouded with black, confining the white within a series of oval spaces; hind-wing, with the veinlets for a short distance *only* from the margin, and ciliae sharply defined with black; between each veinlet from anterior margin to anal angle a lanceolate black mark. UNDER-SIDE with all the veins and veinlets sharply and broadly defined with black, also the lanceolate marks; base of wing yellow. Expanse of wings $2\frac{1}{4}$ in.

B.

PAPILIONES (*part*), Horsfield, *Cat. Lep. Mus. E.I.C.* p. 59 (1828).

PAPILIONIDÆ, Leach, *Edinb. Encycl. IX.* p. 127 (1815). *Dup. C.*

p. 21. E. Doubleday, Diurnal Lep. p. 1. *Westwood (Introd.).*
Stephens. G. R. Gray. Stainton.

PAPILIONIDES, Boisduval, *Spéc. Gén. Lép. I.* pp. 163, 171 (1836); et

Index Méth. Eur. Lép. p. 1 (1840). *Westwood (Introd.).*
Blanchard, Hist. Nat. des Ins. II. p. 325 (1845).

PAPILIONINÆ et PARNASSINÆ, Swainson, *Cabinet Cyclop.* pp. 87, 90
(1840).

PAPILIONIDI, Stephens, *Cat. Brit. Lep. Brit. Mus. pt. I.* p. 1 (1850).
Stainton, Manual Brit. Butt. pp. 12, 14 (1856).

ARCHONTES HEROICI, A. DOMINANTES, et A. SAPIENTES, Hübner.

PAP.-PARNASSII (*part*), Linnæus et Fabricius.

LEP. HEXAPODA (*part*), Latreille, *Règ. Anim. V.* p. 374 (1829).

PAPILIONIDÆ (*part*), Swainson, *Cab. Cyclop.* p. 86 (1840).

PAPILIONII (*part*), Blanchard, *Hist. Nat. des Ins. II.* pp. 322, 324
(1845).

CHEVALIERS, Boisduval, *Hist. Gén. des Lép. Amér. Sep.* p. 2 (1829).

Genus PARNASSIUS, Latreille.

PARNASSIUS, Latreille, *H. N. des Crust. et Ins. XIV.* p. 110 (1805).

DORITIS, Fabricius, *Syst. Gloss.* (*ined.*).

PARNASSIS, Hübner, *Verz. bek. Schmett.* p. 90 (1816).

PIERIS, A. Schrank, *F. B. II. I.* p. 160 (1801).

171. PARNASSIUS HARDWICKI, G. R. Gray.

Parnassius Hardwickii, G. R. Gray, *Gray's Zool. Misc.*
p. 32 (1832); *Lep. Ins. Nepal*, p. 7, t. 4, f. 1, 1a.

Boisduval, *Spéc. Gén. Lép.* I. p. 400. *Doubleday, List Lep. Brit. Mus.* pt. I. p. 21; *Diurnal Lep.* p. 27, n. 10. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 76, n. 352, t. 12, f. 8—11; id. *List Lep. Brit. Mus.* pt. I. p. 91.

a. b. ♂. N. India. Presented by Colonel Hearsey.

This species was taken by Major Charlton in Chinese Tartary, at an elevation of 15,000 feet.

Genus LEPTOCIRCUS, Swainson.

LEPTOCIRCUS, *Swainson, Zool. Ill. 2nd ser. t. 106* (1832).

IPHICLIDES, *pt. Hübner, Zuträge, f. 645—6* (1832).

LAMPROSURA, *Boisduval*.

172. LEPTOCIRCUS CURIUS, Fabricius Sp.

Papilio Curius, Fabricius, Mant. Ins. II. p. 9 (1787); *Ent. Syst. III. pt. I. p. 28. Donovan, Ins. Ind. t. 47, f. 1.*

Leptocircus Curius, E. Doubleday, Zoologist, III. cumfig.; List Lep. Brit. Mus. pt. I. p. 19; Diurnal Lep. p. 23, n. 1, t. 4, f. 1. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 73, n. 336; id. List Lep. Brit. Mus. pt. I. p. 86.*

a. b. ♂ ♀. N. India. Presented by Colonel Sykes.

c. d. ♂ ♀. Cherra Poonjee. Presented by Colonel Buckley.

173. LEPTOCIRCUS MEGES, Zinken-Sommer Sp.

Papilio Meges, Zinken-Sommer, Nova Acta Acad. Nat. Cur. XV. p. 161 (1831).

Leptocircus Meges, E. Doubleday, Zoologist, III. p. 23, fig.; List Lep. Brit. Mus. pt. I. p. 20; Diurnal Lep. p. 23, n. 2. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 73, n. 337.

Leptocircus Curius, apud Swainson, Zool. Ill. 2nd ser. t. 106. Boisduval, Spéc. Gén. Lép. I. p. 381; t. 3, B.f. 1; et t. 1, C.f. 3.

Erycina Curius, apud Godart, Enc. Méth. IX. p. 827.

Iphicrides Curius, apud Hübner, Zuträge, f. 645—6.

a. b. ♂ ♀. Java. From Mr. Argent's Collection.

Genus TEINOPALPUS, *Hope.**TEINOPALPUS*, *Hope*, *Trans. Linn. Soc. XIX.* p. 131 (1843).174. *TEINOPALPUS IMPERIALIS*, *Hope.*

♂ *Teinopalpus imperialis*, *Hope*, *Trans. Linn. Soc. XIX.*
p. 131, t. 11, f. 1, 2. *Westwood, Arcana Ent. t. 59.*
Doubleday, List Lep. Brit. Mus. I. p. 1; *Diurnal*
Lep. p. 2. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.*
p. 80, n. 367; *id. List Lep. Brit. Mus. pt. I.* p. 95.

♀ *Teinopalpus Parryiæ*, *Hope*, *Trans. Linn. Soc. XIX.*
p. 132. *Westwood, Arcana Ent. t. 60.* *Doubleday,*
List Lep. Brit. Mus. pt. I. p. 2.

a. b. ♂. Darjeeling. From Pearson's Collection.

c. ♂. Darjeeling. Presented by Colonel Buckley.

d. e. f. ♂ ♀. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

"This beautiful genus, of which one species only is yet known, may be distinguished at a glance from the others of this family, by its long porrect palpi. There is little else in its structure to separate it from *Ornithoptera* or *Papilio*, though some of its peculiarities indicate an approach to *Thais*, a genus in which the palpi are more developed than in any other of the *Papilionidæ*, with the exception of *Teinopalpus*. The posterior wings differ materially in the two sexes; in the male they are dentate, one-tailed; in the female, three-tailed.

Nothing is known of its habits beyond the fact of its inhabiting the highest ranges of the Himalayas, especially towards the frontiers of Assam, and there it seems to be of very rare occurrence."—
(Doubleday, Diurnal Lep.)

Genus ORNITHOPTERA, *Boisduval.**ORNITHOPTERA*, *Boisduval, Faune de l'Océan.* t. 4, f. 1 (1832).*TROIDES*, *Hübner, Verz. bek. Schmett.* 87 (1816).*AMPHRISIUS*, *Swainson, Zool. Ill. 2nd ser.* t. 98 (1833).*PAPILIO* (*ORNITHOPTERA*), *G. R. Gray, Cat. Lep. Brit. Mus.* (1852).175. *ORNITHOPTERA RICHMONDIA*, *G. R. Gray.**Papilio* (*Ornithoptera*) *Richmondia*, *G. R. Gray, Cat.*

*Lep. Brit. Mus. pt. I. p. 2, t. 2, f. 1, 2 (1852);
List of Lep. Brit. Mus. pt. I. p. 2.*

a. b. ♂ ♀. Richmond River, New South Wales.
Collected by F. Strange, Esq. From Captain Parry's Collection.

176. ORNITHOPTERA DARSIUS, G. R. Gray.

Papilio (Ornithoptera) Darsius, G. R. Gray, *Cat. Lep. Brit. Mus. pt. I. p. 5, n. 11 (1852); List Lep. Brit. Mus. pt. I. Pap. p. 4.*

Ornithoptera Amphimedon, *apud E. Doubleday, Gen. Diurnal Lep. t. 1, f. 2, ♂.*

a. b. ♂. Ceylon. Donor unknown.

c. ♀. Drawing. Ceylon. From Jonville's Collection.

The larva and pupa of *Ornithop. Darsius* are figured on Plate II., fig. 2, 2a, from Ceylon, where discovered by E. L. Layard, Esq., from whose drawings our figures were copied.

177. ORNITHOPTERA POMPEUS, Cramer Sp.

*♀ Papilio Pompeus, Cramer, Pap. Ex. t. 25, f. A. (1775).
Esper, Ausl. Schmett. t. 24, f. 2, ♀.*

Ornithoptera Pompeus, *E. Doubleday, Diurnal Lep. p. 4, n. 9.*

Papilio (Ornithoptera) Pompeus, G. R. Gray, *Cat. Lep. Brit. Mus. pt. I. p. 5, n. 13; List Lep. Brit. Mus. pt. I. p. 5.*

♀ Papilio Astenous, Fabricius, Syst. Ent. II. p. 448 (1775); Ent. Syst. III. I. p. 19.

Troides Astenous, *Hübner, Verz. bek. Schmett. p. 88.*

♀ Papilio Minos, Cramer, Pap. Ex. t. 195, f. A. (1782).

Troides Minos, *Hübner, Verz. bek. Schmett. p. 88.*

♂ ♀ Papilio Heliacon, Fabricius, Ent. Syst. III. I. p. 19 (1793). Donovan, Ins. of Ind. t. 22, f. 1; ex Jones, Icon. ined. I. t. 42.

Ornithoptera Heliacon, *Boisduval, Spéc. Gén. Lép. I. p. 178. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 2.*

Amphrisius Nymphalides, *Swainson, Zool. Ill. 2nd ser. t. 98.*

Papilio Amphrisius, *pt. Godart, Enc. Méth.* IX. *p. 27.*
Papilio Amphrisius, *apud Horsfield, Cat. Lep. Mus. E.I.C.*
t. IV. f. 13, larva et pupa.

- a. b. c. d. e. f. ♂ ♀.* Java. From Dr. Horsfield's Collection.
- g. ♂.* Darjeeling. From Pearson's Collection.
- h. i. ♂.* Cherra Poonjee. Presented by Colonel Buckley.
- j. ♂.* Darjeeling. From Messrs. Schlagintweit's Collection.

The larva and pupa of *Ornithop. Pompeus* are figured on Plate II., fig 1, 1a, from Java, where it "feeds on a species of *Aristolochia*, bearing the native name of *Wallas*. March. Scarce."—(Horsfield, MS.)

178. ORNITHOPTERA RHADAMANTHUS, Boisduval.

Ornithoptera Rhadamanthus, Boisduval, Spéc. Gén. Lép.
I. p. 180 (1836). E. Doubleday, Diurnal Lep.
p. 4, n. 10.

Papilio (Ornithoptera) Rhadamanthus, G. R. Gray, Cat.
Lep. Brit. Mus. pt. I. p. 6, n. 14; List Lep. Brit.
Mus. pt. I. p. 5.

- ♂ *Papilio Amphrisius, apud Lucas, Lep. Exot. t. 2, f. 1.*
- a. ♂.* N. India. Presented by Colonel Sykes.
- b. ♀.* Canara? Presented by S. N. Ward, Esq.
- c. ♀.* Darjeeling. From Pemberton's Collection.
- d. e. ♂ ♀.* Darjeeling. From Messrs. Schlagintweit's Collection.
- f. ♀.* Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

179. ORNITHOPTERA AMPHRISIUS, Cramer Sp.

♂ *Papilio Amphrysus, Cramer, Pap. Ex. t. 219, f. A.*
(1782).

Ornithoptera Amphrisius, Boisduval, Spéc. Gén. Lép. I.
p. 178. De Haan, Verh. Nat. Gesch. Ned. p. 19?
E. Doubleday, Diurnal Lep. p. 4, n. 8.

Papilio Amphrisius, Fabricius, Ent. Syst. III. I. p. 11.
Godart, Enc. Méth. IX. p. 27, pt. n. 7.

Papilio Amphrisus, *Herbst*, *Pap.* t. 1, f. 3. *Esper*, *Ausl. Schmett.* t. 34, f. 1.

Troides Amphrysus, *Hübner*, *Verz. bek. Schmett.* p. 88.

Papilio (Ornithoptera) Amphrisius, *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 6, n. 15; *List Lep. Brit. Mus. pt. I.* p. 6.

a. b. c. d. ♂ ♀. Borneo. Presented by H. Lowe, Esq.

Genus PAPILIO, *Linnæus.*

PAPILIO, *Linnæus*. *Fabricius*. *Godart*. *Horsfield*. *Boisduval*.
Doubleday. *G. R. Gray*.

ZELIMA, *Fabricius*. *Zinken-Sommer*.

AMARYSSUS, *Dalm.*, *P. S.* p. 85 (1816).

IPHICLIDES, JASONIADES, EUPHÆADES, HERACLIDES, LAERTIADES, MENELAIDES, ACHILLIDES, IDAIDES, ZETIDES, ORPHEIDES, NESTORIDES, CALAIDES, PRIAMIDES, PARIDES, ITHOBALUS, ILIADES, ARISBE, ZELIDES, HECTORIDES, ASCANIDES, AMAURIS, *Hübner*, *Verz.* (1816).

CLYTIA, POLYDORUS, THOAS, CHLORISSES, PROTESILAUS, PODALIRIUS, *Swainson*, *Zool. Ill. 2nd ser.* (1829-33).

180. PAPILIO NOX, *Swainson.*

Papilio Nox, *Swainson*, *Zool. Ill. 1st ser.* t. 102 (1822).
Horsfield, *Cat. Lep. Mus. E.I.C.* t. 1, f. 15. *Boisduval*, *Spéc. Gén. Lép.* I. p. 277. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 11; *Diurnal Lep. p. 9, n. 12*. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 8, n. 28; *id. List Lep. Brit. Mus. pt. I.* p. 9.

Papilio Memercus, *Godart*, *Enc. Méth. IX. Suppl.* p. 809 (1823).

Papilio Neesius, *Zinken-Sommer*, *Nov. Act. Acad. Nat. Cur. XV.* t. 14, f. 4 (1831).

Papilio Nox, var. ♀, *De Haan*, *Verh. Nat. Gesch. Ned. Overz. Bez. Ins.* p. 41, t. 5, f. 3.

a. ♀. Java. From Dr. Horsfield's Collection.

181. PAPILIO ASTINA, *Horsfield.*

Papilio Astina, *Horsfield*. *Westwood*, *Cabinet Orient. Ent.* p. 20, t. 9, f. 3; *id. Doubleday's Diurnal Lep.*

Add. p. 529. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 72, n. 335; id. *List Lep. Brit. Mus.* pt. I. p. 85.

a. ♂. (type). Java. From Dr. Horsfield's Collection.

182. *PAPILIO MACAREUS*, Godart.

Papilio Macareus, Godart, *Enc. Méth. IX.* p. 76 (1819).

Horsfield, Cat. Lep. Mus. E.I.C. t. 5, f. 1. *Bois-duval, Spéc. Gén. Lép.* I. p. 374. *Lucas, Lep. Exot.* t. 23, f. 1. *E. Doubleday, Diurnal Lep.* p. 21, n. 259; *List Lep. Brit. Mus.* pt. I. App. p. 4.

Papilio striatus, Zinken-Sommer, *Nova Acta Acad. Nat. Cur.* XV. t. 14 (1831).

Papilio Xenocles, var. a., *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 71; *List Lep. Brit. Mus.* pt. I. p. 83.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

183. *PAPILIO MEGARUS*, Westwood.

Papilio Megarus, Westwood, *Arcana Ent.* p. 98, t. 72, f. 3 (1845). *E. Doubleday, Diurnal Lep.* p. 20, n. 256; *List Lep. Brit. Mus.* pt. I. App. p. 4. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 71, n. 326; id. *List Lep. Brit. Mus.* pt. I. p. 83.

a. ♂. Silhet. Presented by the Trustees of the British Museum.

184. *PAPILIO XENOCLES*, E. Doubleday.

Papilio Xenocles, E. Doubleday, *Gray's Zool. Misc.* p. 74 (1842); *List Lep. Brit. Mus.* pt. I. p. 19; *Diurnal Lep.* p. 20, n. 258. *Westwood, Arcana Ent.* p. 127, t. 92, f. 1. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 71, n. 327; id. *List Lep. Brit. Mus.* pt. I. p. 83.

Papilio Pollux, apud Erichson, *Wiegmanns Archiv. für Naturg.* II. p. 248 (1843).

a. b. c. ♂ ♀. Silhet. Presented by Colonel Buckley.

d. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

185. *PAPILIO DELESSERTI*, Guérin.

- Papilio Delessertii, *Guérin, Rev. Zool.* (1839), p. 233 ;
Deless. Souv. Voy. dans l'Inde, Ins. t. 17. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 19; Diurnal Lep. p. 20, n. 257. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 71, n. 325; id. List Lep. Brit. Mus. pt. I. p. 82.
Papilio Laodocus, *De Haan, Verh. Naturl. Gesch.* p. 42, t. 8, f. 5 (1839).
Papilio Melanides (*De Haan*), *Erichson, Wieg'm. Archiv. für Naturg.* II. p. 248 (1843).
a. ♀. Java (?). From Dr. Horsfield's Collection (?).

186. *PAPILIO AGESTOR*, G. R. Gray.

- Papilio Agestor, *G. R. Gray, Lep. Ins. of Nepal*, t. 4, f. 2 (1831). *J. E. Gray's Zool. Misc.* (1832), p. 32.
Boisduval, Spéc. Gén. Lép. I. p. 376. *Westwood, Arcana Ent.* p. 59, t. 16, f. 2. *Kollar, in Hügel's Kaschmir, pt. II.* p. 406. *E. Doubleday, List Lep. Brit. Mus. pt. I. p. 19; Diurnal Lep.* p. 21, n. 262. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 71, n. 323; id. List Lep. Brit. Mus. pt. I. p. 82.*

- a. ♂. N. India. Presented by Colonel Buckley.
b. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

"This species is one of the earliest of the genus that is on the wing, being found in woody situations in April and May, dancing lightly over the tops of low bushes and trees, with a sailing kind of flight, gliding along without moving the wings. It is by no means rare at Masuri."—(Capt. T. Hutton, *Trans. Ent. Soc.* V. p. 50.)

187. *PAPILIO DISSIMILIS*, Linnæus.

- ♂ Papilio dissimilis, *Linn., S.N.* II. p. 782 (1767). *Clerck, Icon.* t. 16, f. 2. *Cramer, Pap.* I. t. 82, f. C.D. *Fabricius, Ent. Syst.* III. I. p. 38. *Herbst, Pap.* t. 126, f. 2, 3. *Godart, Enc. Méth.* IX. p. 175. *Boisduval, Spéc. Gén. Lép.* I. p. 377. *Lucas, Lep. Exot.* t. 23, f. 2. *E. Doubleday, List Lep. Brit. Mus. pt. I. p. 19; Diurnal Lep.* p. 21, n. 263.

- G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 71,*
n. 330; id. List Lep. Brit. Mus. pt. I. p. 84.
- Clytia dissimilis, Swainson, Zool. Ill. 2nd ser. t. 120.*
- ♀ *Papilio Panope, Linnæus, S. N. II. p. 782.* *Cramer,*
Pap. III. t. 295, f. E.F. Fabricius, Ent. Syst. III.
I. p. 59. Boisduval, Spéc. Gén. Lép. I. p. 373.
Kollar, in Hügel's Kaschmir, pt. II. p. 406. E.
Doubleday, List Lep. Brit. Mus. pt. I. p. 19.
- Princeps Dominans Panope, Hübner, Samml. Exot. I.*
t. 132.
- Arisbe Panope, Hübner, Verz. bek. Schmett. p. 89.*
- Papilio Panopes, Godart, Enc. Méth. IX. p. 75.*
- Var. *a.* *Papilio Clytia, Linnæus, S. N. II. p. 781.* *Fabricius,*
Ent. Syst. III. I. p. 127.
- Papilio Echidna, De Haan, Verh. Naturl. Gesch. p. 42,*
t. 8, f. 6 (18—).
- Var. *b.* *Papilio Palephates, Westwood, Arcana Ent. p. 127,*
pl. 76, f. 1. E. Doubleday, Diurnal Lep. p. 21,
n. 265.
- a. b. c. ♂ ♀.* N. India. Presented by Colonel
 Buckley.
- d. ♀.* Canara. Presented by S. N. Ward, Esq.
- e. ♀,* with anterior wings *wholly* brown. Donor un-
 known.

The larva and pupa of *P. dissimilis* are figured on Plate II., fig. 3, 3a, 3b, and were copied from the drawings of General Hardwicke, now in the Library of the British Museum. This species has been also reared in Ceylon by E. L. Layard, Esq. The transformations were also observed by Lady Gilbert, among whose drawings figures are given. Her ladyship states that "it feeds on a plant called by the natives *Maike*. The larva was obtained on the 26th July; two days after it commenced its transformation, and on the following day the chrysalis resembled a dried twig, adhering to the frame only by the extremity of the tail, and supported on each side by two fine threads: in this state it died."

"Occurs in warm glens as well as in the Dhoon, but it does not appear to be very numerous."—(Captain T. Hutton, Trans. Ent. Soc. V. p. 49.)

188. *PAPILIO CASTOR*, Westwood.

- ♂ Papilio Castor, *Westwood*, *Ann. N. H.* IX. p. 37 (1842);
Arcana Entom. p. 129, t. 80, f. 1, 2. *E. Doubleday*,
List Lep. Brit. Mus. pt. I. p. 4; *Diurnal Lep.*
p. 12, n. 72. *G. R. Gray*, *Cat. Lep. Brit. Mus.*
pt. I. p. 21, n. 93; id. *List Lep. Brit. Mus.* pt. I.
p. 28.
- ♀ Papilio Pollux, *Westwood*, *Ann. N. H.* IX. p. 37 (1842);
Arc. Ent. p. 129, t. 90, f. 1. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 19; *Diurnal Lep.* p. 21,
n. 264.
- a. ♂. Cherra Poonjee. Presented by Colonel Buckley.
- b. ♂. Silhet. Presented by the Trustees of the British Museum.
- c. ♀. N. India. Donor unknown.

189. *PAPILIO HECTOR*, Linnaeus.

- Papilio Hector, *Linnaeus*, *S. N.* II. p. 745 (1767).
Clerck, Icon. t. 33, f. 1. *Cramer, Pap.* II. t. 141,
f. A. Fabricius, Ent. Syst. III. I. p. 3. *Herbst, Pap.* t. 13, f. 2. *Esper, Ausl. Schmett.* t. 1, f. 2.
Godart, Enc. Méth. IX. p. 70. *Boisduval, Spéc. Gén. Lép.* I. p. 269. *Lucas, Lep. Exot.* t. 4, f. 2.
De Haan, Verh. Naturl. Gesch. p. 39. *Westwood, Arcana Ent.* p. 9, t. 3. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 10; *Diurnal Lep.* p. 9, n. 25. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 11, n. 41; id. *List Lep. Brit. Mus.* pt. I. p. 13.

Menelaides Hector, *Hübner, Verz. bek. Schmett.* p. 84.
Princeps Heroicus Hector, *Hübner, Samml. Exot.* I. t. 109.

- a. b. ♂ ♀. Calcutta. Presented by Colonel Buckley.
- c. d. ♂. N. India. Donor unknown.
- e. f. ♀. Madras. In exchange.
- g. Ceylon. Drawing. From Jonville's Collection.

The larva and pupa of *Pap. Hector* are figured on Plate II., fig. 4, 4a, 4b, from General Hardwicke's drawings.

"The larva feeds on *Aristolochia indica*."—(Hardwicke.)

190. *PAPILIO DIPHILUS*, Esper.

Papilio Diphilus, *Esper*, *Ausl. Schmett.* t. 40, f. 1 (1801).

E. Doubleday, *Diurnal Lep.* p. 9, n. 20. *G. R. Gray*,
Cat. Lep. Brit. Mus. pt. I. p. 10, n. 34; id. *List Lep. Brit. Mus. pt. I.* p. 11.

Polydorus Thoas, *Swainson*, *Zool. Ill. 2nd ser.* t. 100
 (1833).

Papilio Polydorus, *apud Cramer*, *Pap. II.* t. 128, f. A.B.
Herbst, *Pap.* t. 15, f. 3. *Godart*, *Enc. Méth.* IX.
 p. 71 (part). *Horsfield*, *Cat. Lep. Mus. E.I.C.*
 t. 3, f. 17. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.*
 p. 10.

Menelaides Polydorus, *Hübner*, *Verz. bek. Schmett.* p. 84.

Princeps Heroicus Polydorus, *Hübner*, *Sammel. Exot.* I.
 t. 110.

Papilio Polydorus, var., *Boisduval*, *Spéc. Gén. Lép.* I.
 p. 267. *De Haan*, *Verh. Naturl. Gesch.* p. 38, t. 8,
 f. 1.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. N. India. From the Asiatic Society of Bengal.

g. h. ♀. Calcutta. Presented by Colonel Buckley.

The larva and pupa of *Pap. Diphilus* are figured on Plate II., fig. 5, 5a, from Java.

191. *PAPILIO ANTIPHUS*, Fabricius.

Papilio Antiphus, *Fabricius*, *Ent. Syst.* III. pt. I. p. 10.

Donovan, *Ins. of Ind.* t. 15, f. 2; ex *Jones*, *Icon. ined.* I. t. 14. *Godart*, *Enc. Méth.* IX. p. 71. *Boisduval*, *Spéc. Gén. Lép.* I. p. 266. *De Haan*, *Verh. Naturl. Gesch.* p. 49, t. 8, f. 2, ♀. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 9; *Diurnal Lep.* p. 9, n. 21. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 11, n. 37; id. *List Lep. Brit. Mus. pt. I.* p. 12.

Papilio Polygius, *Godart*, *Enc. Méth.* IX. *Suppl.* 8, n. 11, 129-30 (1823).

? ♀ *Papilio Theseus*, *Cramer*, *Pap. II.* t. 180, f. B. (1776).
Fabricius, *Ent. Syst.* III. pt. I. p. 2. *Herbst*, *Pap.*

t. 14, f. 3. Esper, Ausl. Schmett. t. 36, f. 3. Godart, Enc. Méth. IX. p. 71. Boisduval, Spéc. Gén. Lép. I. p. 276. E. Doubleday, Diurnal Lep. p. 11, n. 63.

Menelaides Theseus, Hübner, Verz. bek. Schmett. p. 84.

Var. *a.*, *G. R. Gray.*

Papilio Kotzebuæ, Eschscholtz, Voy. Kotzebue, t. 1, f. 2.

Papilio Antiphus, ♂, De Haan, Verh. Naturl. Gesch. pp. 5, 40.

a. ♂. Java. Presented by the Trustees of the British Museum.

Var. *b.*, *G. R. Gray*, Cat. Lep. Brit. Mus. pt. I. p. 11.

a. b. ♂. Borneo. Presented by H. Lowe, Esq.

192. PAPILIO COON, Fabricius.

Papilio Coon, Fabricius, Ent. Syst. III. I. p. 10 (1793).
Donovan, Ins. of China, t. 24, f. 1; ex Jones, Icon. ined. I. t. 36. Godart, Enc. Méth. IX. p. 65. Lucas, Lep. Exot. t. 6, f. 2. Boisduval, Spéc. Gén. Lép. I. p. 201. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 3; Diurnal Lep. p. 10, n. 41. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 16, n. 60; id. List Lep. Brit. Mus. pt. I. p. 19.

Papilio Hypenor, Godart, Enc. Méth. IX. p. 65 (1819).

Achillides Hypenor, Hübner, Samml. Exot. III. t. 491.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. ♂. N. India. Presented by Colonel Buckley.

e. ♂. China. Presented by Dr. Cantor, 1844.

193. PAPILIO ALCINOUS, Klug.

Papilio Aleinus, Klug, Neue Schmett. der Insekt.-Samml. Berlin (1836), t. 1. De Haan, Verh. Nat. Gesch. Insect. p. 26. E. Doubleday, Diurnal Lep. p. 9, n. 19. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 12, n. 45; List Lep. Brit. Mus. pt. I. p. 14.
Hab. Japan.—(Siebold.)

Var. *a.*, *G. R. Gray*, Cat. Lep. Brit. Mus. pt. I. p. 12, t. 4, f. 2, ♂, 3, ♀; List. Lep. Brit. Mus. pt. I. p. 14.

a. b. ♂. Bootan. From Pemberton's Collection.

The larva of *Pap. Alcinous* is figured on Plate II., fig. 6, copied from De Haan's published drawing.

194. *PAPILIO PHILOXENUS*, G. R. Gray.

♀ Papilio Philoxenus, G. R. Gray, *Lep. Ins. of Nepal*, p. 5, t. 2 (1831). *J. E. Gray's Zool. Misc.* (1832), p. 32. *Boisduval, Spéc. Gén. Lép.* I. p. 264. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 9; *Diurnal Lep.* p. 9, n. 14. *Westwood, Cab. Orient. Ent.* t. 40, f. 2. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 9, n. 31; *id. List Lep. Brit. Mus. pt. I.* p. 10.

Var. *a.*, G. R. Gray.

♂ Papilio Polyeuctes, *E. Doubleday, Gray's Zool. Misc.* p. 74 (1842); *Diurnal Lep.* p. 9, n. 15, t. 2, f. 3. *Westwood, Cab. Orient. Ent.* t. 40, f. 3.

Papilio Boötes, *pt.*, Erichson.

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.

c. d. ♂ ♀. Cherra Poonjee. Presented by Colonel Buckley.

e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

195. *PAPILIO DASARADA*, Moore.

♂ Papilio Philoxenus, var., *Westwood, Cab. Orient. Ent.* t. 40, f. 5.

Papilio Philoxenus, var. B., G. R. Gray, *Cat. Lep. Brit. Mus. pt. I.* p. 9; *List Lep. Brit. Mus. pt. I.* p. 10.

a. b. ♂ ♀. Cherra Poonjee. Presented by Colonel Buckley.

196. *PAPILIO RAVANA*, Moore.

♀ Papilio Philoxenus, var., *Westwood, Cab. Orient. Ent.* t. 40, f. 4.

Papilio Philoxenus, var. C., G. R. Gray, *Cat. Lep. Brit. Mus. pt. I.* p. 9; *List Lep. Brit. Mus. pt. I.* p. 10.

a. b. ♂ ♀. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

197. *PAPILIO MINEREUS*, G. R. Gray.

♀ Papilio Minereus, *G. R. Gray, Lep. Ins. of Nepal*, p. 5, t. 1 (1831). *J. E. Gray's Zool. Misc.* (1832), p. 32; *Cat. Lep. Brit. Mus. pt. I.* p. 9, n. 32; id. *List Lep. Brit. Mus. pt. I.* p. 10. *Westwood, Cab. Orient. Ent.* t. 40, f. 1. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 9; *Diurnal Lep.* p. 9, n. 16.

Papilio Latreillii, Donovan, Nat. Reposit. vol. IV. t. 140.

Papilio Philoxenus, ♀, *Boisduval, Spéc. Gén. Lép.* I. p. 264.

Papilio Philoxenus, var., *Westwood*.

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

198. *PAPILIO JANAKA*, Moore.

Papilio Janaka, Moore, Proc. Zool. Soc. (1857); *Annulosa*, t. , f. 1, 1a.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

b. c. Darjeeling. From Messrs. Schlagintweit's Collection.

Pap. Janaka, n. sp.—UPPER-SIDE black; hind-wing with a white patch on the disc, which is divided by three of the veinlets, thus forming four separate patches, the outer one on each side being the shortest, and the two nearest the abdominal margin being tinged with red; three sub-marginal and three marginal lunules and circular mark at anal angle red; tail with two red spots. UNDER-SIDE black; fore-wing with the base red; hind-wing with patch on disc as on upper-side, but the one nearest abdominal margin nearly covered with red, which colour is continued upwards and downwards, occupying the base of the wing and the whole space between the third median veinlet and sub-median vein; lunules as above, but are larger, and a fourth sub-marginal one appears between the discoidal and first median veinlets; tail spotted as above; ciliae between the angles white; head, neck, body beneath, and sides, red. Wings shaped as in *P. Boötes*, Westwood, Arc. Ent. t. 31. Expanse of wings 5 in.

Remark.—*Papilio Boötes* appears to be a near ally of *P. Janaka*.

199. *PAPILIO VARUNA*, *White*.

Papilio Varuna, *White*, *Entomologist* (1842), p. 280.

Westwood, *Ann. N. H.* IX. n. s. p. 37. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 11; *Diurnal Lep. p. 9, n. 11*. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I. p. 8, n. 27*; *id. List Lep. Brit. Mus. pt. I. p. 9*.

♂ *Papilio Astorion*, *Westwood*, *Ann. N. H.* (1842), p. 37; *Arc. Ent. p. 69, t. 66, f. 1*.

♀ *Papilio Chara*, *Westwood*, *Arc. Ent. p. 69, t. 66, f. 2* (1845).

a. ♂. N. India. Presented by the Trustees of the British Museum.

b. ♀. N. India. Presented by Lieutenant Hugo James.

c. d. e. f. ♂ ♀. Darjeeling. Reserved from the Indian Collection, Exposition Universelle at Paris, 1855.

200. *PAPILIO RHETENOR*, *Westwood*.

Papilio Rhetenor, *Westwood*, *Arcana Ent. p. 59, t. 16*,

f. 1, 1a (1842). *Kollar*, *in Hügel's Kaschmir, pt. II. p. 403*. *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 3*; *Diurnal Lep. p. 10, n. 35*. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I. p. 15, n. 54*; *id. List Lep. Brit. Mus. pt. I. p. 18*.

a. b. ♀. Darjeeling. From Pearson's Collection.

c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

201. *PAPILIO PROTENOR*, *Cramer*.

♂ *Papilio Protenor*, *Cramer*, *Pap. I. t. 49, f. A. B.* (1775).

Fabricius, *Ent. Syst. III. I. p. 13*. *Herbst*, *Pap. t. 7, f. 1, 2*. *Esper*, *Ausl. Schmett. t. 33, f. 2*; *t. 29, f. 2?* *Godart*, *Enc. Méth. IX. p. 30*. *Boisduval*, *Spéc. Gén. Lép. I. p. 198*. *De Haan*, *Verh. Naturl. Gesch. p. 25*. *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 3*; *Diurnal Lep. p. 10, n. 34*. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I. p. 14, n. 52*; *id. List Lep. Brit. Mus. pt. I. p. 17*.

- Iliades Protenor, *Hübner, Verz. bek. Schmett.* p. 89.
 ♀ Papilio Laomedon, *Fabricius, Ent. Syst.* III. I. p. 12.
Donovan, Ins. of China, t. 25; ex Jones, Icon. ined. I. t. 10.
 a. ♂. Bootan. From Pemberton's Collection.
 b. c. ♂ ♀. N. India. Presented by Colonel Buckley.
 d. ♀. N. India. In exchange.

"Not an uncommon species in the warm glens of the Masuri Hills during the summer months, and it is common in the Dhoon. Its flight is somewhat heavy and unsteady."—(Captain T. Hutton, *Trans. Ent. Soc.* V. p. 49.)

202. PAPILIO MEMNON, Linnæus.

- ♂ Papilio Memnon, *Linnæus, S. N.* II. p. 747 (1767).
Cramer, Pap. Exot. I. t. 91, f. C. *Fabricius, Ent. Syst.* III. I. p. 12. *Herbst, Pap.* t. 6, f. 2, 3. *Esper, Ausl. Schmett.* t. 20, f. 3. *Godart, Enc. Méth.* IX. p. 29. *Swainson, Zool. Ill. 2nd ser.* t. 95. *De Haan, Verh. Naturl. Gesch.* p. 23, t. 3, f. 1. *Boisduval, Spéc. Gén. Lép.* I. p. 192. *E. Double-day, List Lep. Brit. Mus. pt. I.* p. 2; id. *Diurnal Lep.* p. 10, n. 30. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 13, No. 47; id. *List Lep. Brit. Mus. pt. I.* p. 14.

Iliades Memnon, *Hübner, Verz. bek. Schmett.* p. 89.

Papilio Arbates, *Zinken-Sommer, Nova Acta Acad. Nat. Cur.* XV. p. 151 (1831).

- a. b. c. ♂. Java. From Dr. Horsfield's Collection.

Var. A. ♂ Papilio Androgeos, *Cramer, Pap. Exot.* I. t. 91, f. A. B. (1775).

Papilio Memnon, *Esper, Ausl. Schmett.* t. 8, f. 1.

Iliades Mestor, *Hübner, Verz. bek. Schmett.* p. 89 (1816).

- d. ♂. Pinang. Presented by Dr. Cantor, 1844.

e. ♂. Darjeeling. Reserved from Indian Collection, *Exposition Universelle at Paris, 1855.*

Var. B. ♀ Papilio Anceus, *Cramer, Pap. Exot.* III. t. 222, f. A. B. (1780).

Iliades Anceus, *Hübner, Verz. bek. Schmett.* p. 88.

Var. C. ♀ Papilio Laomedon, *Cramer, Pap. Exot.* I. t. 50, f. A. B. (1775).

Iliades Laomedon *Hübner, Verz. bek. Schmett.* p. 89.

Papilio Memnon (*pt.*), *Fabricius, Ent. Syst.* III. I. p. 12.

f. ♀. Java. From Dr. Horsfield's Collection.

Var. D. Papilio Memnon, var. d., *G. R. Gray.*

g. h. ♀. Cherra Poonjee. Presented by Colonel Buckley.

Var. E. ♀ Papilio Laomedon, var., *De Haan, Verh. Nat. Gesch.* p. 24, t. 3, f. 2.

Var. F. Papilio Memnon, var. f., *G. R. Gray.*

Var. G. ♀ Papilio Agenor, *Linnæus, Syst. Nat.* II. p. 747 (1767).
Clerck, Icon. t. 15. *Cramer, Pap. Exot.* I. t. 32, f. A. B. *Fabricius, Ent. Syst.* III. I. p. 13. *Herbst, Pap.* t. 8, f. 3. *Esper, Ausl. Schmett.* t. 26, f. 1. *Donovan, Ins. of China,* t. 24, f. 2. *Godart, Enc. Méth.* IX. p. 28.

Iliades Agenor, *Hübner, Verz. bek. Schmett.* p. 89.

i. j. ♀. Java. From Dr. Horsfield's Collection.

Var. H. ♀ Papilio Achates, var., *De Haan, Verh. Nat. Gesch.* p. 24, t. 3, f. 3.

Var. I. ♀ Papilio Achates, *Cramer, Pap. Exot.* II. t. 182, f. A. B.; t. 243, f. A. (1777). *Fabricius, Ent. Syst.* III. I. p. 9. *Herbst, Pap.* t. 15, f. 1. *Esper, Ausl. Schmett.* t. 28, f. 1. *Godart, Enc. Méth.* IX. p. 64.

Achillides Achates, *Hübner, Verz. bek. Schmett.* p. 85.

Papilio Achatiades, *Esper, Ausl. Schmett.* t. 28, f. 2, t. 29, f. 1.

k. l. m. n. ♀. Java. From Dr. Horsfield's Collection.

Var. K. ♀ Papilio Alcanor, *Cramer, Pap. Exot.* II. t. 166, f. A. (1776). *Esper, Ausl. Schmett.* t. 34, f. 2.

Achillides Alcanor, *Hübner, Verz. bek. Schmett.* p. 85.

Papilio Alphenor, *Fabricius*, *Spec. Ins.* II. p. 4. *Herbst*,
Pap. t. 16, *f.* 1.

a. ♀. Cherra Poonjee. Presented by Colonel Buckley.

The larva and pupa of *Papilio Memnon* are figured on Plate II.,
 fig. 7, 7a, from Java.

203. *PAPILIO POLYMNESTOR*, Cramer.

Papilio Polymnestor, *Cramer*, *Pap.* I. *t.* 53, *f.* A. B.
 (1775). *Fabricius*, *Ent. Syst.* III. I. *p.* 18. *Herbst*,
Pap. t. 7, *f.* 3, 4. *Esper*, *Ausl. Schmett.* *t.* 19, *f.* 1.
Donovan, *Ins. of Ind.* *t.* 23, *f.* 2. *Godart*, *Enc. Méth.* IX. *p.* 29. *Lucas*, *Lep. Exot.* *t.* 12, *f.* 1.
Boisduval, *Spéc. Gén. Lép.* I. *p.* 191. *De Haan*,
Verh. Naturl. Gesch. *p.* 23. *E. Doubleday*, *List Lep. Brit. Mus.* *pt.* I. *p.* 2; *Diurnal Lep.* *p.* 10,
n. 29. *G. R. Gray*, *Cat. Lep. Brit. Mus.* *p.* 12,
n. 46; *id. List Lep. Brit. Mus.* *pt.* I. *p.* 14.

Iliades Polymnestor, *Hübner*, *Verz. bek. Schmett.* *p.* 88.

a. ♂. Calcutta. Presented by Colonel Buckley.

b. ♀. N. India. From the Asiatic Society of
 Bengal.

c. Ceylon. Drawing. From Jonville's Collection.

The larva and pupa of *Pap. Polymnestor* are figured on Plate III.,
 fig. 1, 1a, from Ceylon, where discovered by E. L. Layard, Esq.

204. *PAPILIO ISWARA*, White.

Papilio Iswara, *White*, *Entomologist* (1842), *p.* 280.
E. Doubleday, *List Lep. Brit. Mus.* *pt.* I. *p.* 4;
Diurnal Lep. *p.* 11, *n.* 58, *t.* 2, *f.* 1. *G. R. Gray*,
Cat. Lep. Brit. Mus. *pt.* I. *p.* 19, *n.* 78; *id. List Lep. Brit. Mus.* *pt.* I. *p.* 23.

a. ♀. Pinang. Presented by the Trustees of the
 British Museum.

205. *PAPILIO HELENUS*, Linnæus.

Papilio Helenus, *Linnæus*, *S. N.* II. *p.* 754 (1767).
Clerck, *Icon.* *t.* 13, *f.* 2. *Cramer*, *Pap.* *t.* 153,
f. A. B. *Fabricius*, *Ent. Syst.* III. I. *p.* 2. *Herbst*,
Pap. t. 14, *f.* 2. *Esper*, *Ausl. Schmett.* *t.* 2, *f.* 2.

Godart, Enc. Méth. IX. p. 68. *Lucas, Lep. Exot.* t. 15, f. 2. *Boisduval, Spéc. Gén. Lép.* I. p. 211. *De Haan, Verh. Naturl. Gesch.* p. 30. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 4; *Diurnal Lep.* p. 11, n. 57. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 18, n. 77; id. *List Lep. Brit. Mus.* pt. I. p. 23.

Achillides Helenus, Hübner, Verz. bek. Schmett. p. 85.

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
- c. ♂. Darjeeling. From Pearson's Collection.
- d. ♀. Darjeeling. Presented by Colonel Buckley.
- e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

The larva and pupa of *Pap. Helenus* are figured on Plate III., fig. 2, 2a, from Canara, where discovered by S. N. Ward, Esq., E. I. C. civil service.

206. *PAPILIO CHAON*, Westwood.

Papilio Chaon, Westwood, Arcana Ent. t. 72, f. 1, 2 (1845). *E. Doubleday, List Lep. Brit. Mus.* pt. I. App. p. 2; *Diurnal Lep.* p. 11, n. 55. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 18, n. 76; id. *List Lep. Brit. Mus.* pt. I. p. 23.

- a. ♂. Assam. From the Asiatic Society of Bengal.
- b. ♀. Cherra Poonjee. Presented by Colonel Buckley.
- c. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.
- d. ♂. Var. Borneo. Presented by H. Lowe, Esq.

207. *PAPILIO ROMULUS*, Cramer.

Papilio Romulus, Cramer, Pap. I. t. 43, f. A. (1775). *Esper, Ausl. Schmett.* t. 25, f. 2. *E. Doubleday, Diurnal Lep.* p. 10, n. 26; id. *List*, p. 13. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 11, n. 42.

Menelaides Romulus, Hübner, Verz. bek. Schmett. p. 84.

Papilio Mutius, Fabricius, Ent. Syst. III. I. p. 3 (1793).

Godart, Enc. Méth. IX. p. 70. *Boisduval, Spéc. Gén. Lép.* I. p. 270. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 10.

Papilio Mutius, var.,? *De Haan, Verh. Naturl. Gesch.*
p. 6.

♀ Papilio Astyanax, *Fabricius, Ent. Syst.* III. I. p. 13
(1793). *Donovan, Ins. of Ind.* t. 23, f. 1; ex *Jones, Icon. ined.* I. t. 20. *Godart, Enc. Méth.* IX. p. 72.
Boisduval, Spéc. Gén. Lép. I. p. 271.

a. b. c. ♂ ♀. N. India. Presented by the Asiatic Society of Bengal.

d. ♀. Calcutta. Presented by Colonel Buckley.

208. PAPILIO POLYTES, Linnæus.

Papilio Polytes, *Linnæus, S. N.* II. p. 746 (1767).
Clerck, Icon. t. 14, f. 1. *Cramer, Pap.* III. t. 265,
f. A. B. C. *Herbst, Pap.* t. 15, f. 2, ♀, t. 20;
f. 3, 4, ♂. *Fabricius, Ent. Syst.* III. pt. I. p. 2.
Esper, Ausl. Schmett. t. 3, f. 1; t. 12, f. 1. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 20, n. 83; id.
List Lep. Brit. Mus. pt. I. p. 25.

Menelaides Polytes, *Hübner, Verz. bek. Schmett.* p. 85.

Princeps Heroicus Stichius, *Hübner, Samml. Exot.* I.
t. 3.

Papilio Pammon ♀, *Boisduval, Spéc. Gén. Lép.* I. p. 272.
De Haan, Verh. Nat. Gesch. p. 41. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 11; *Diurnal Lep.* p. 11, n. 62.

Var. Papilio Numa, *Weber, Observ. Entom. Kiel.* (1801),
p. 106.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Calcutta. From the Asiatic Society of Bengal.

g. h. ♂ ♀. N. India. Presented by Colonel Buckley.
i. ♀. Chusan. Presented by Dr. Cantor, 1842.

"The larva and pupa of *Pap. Polytes* are figured on Plate III., fig. 3, 3a, from Java, where it feeds on a species of *Citrus*, bearing the native name of *Juruk*, from January to May."—(Horsfield.)

"Not uncommon in the Dhoon during the rainy season, and at Rajpore, at the foot of the Hills, it is frequently met with."—(Captain T. Hutton, Trans. Ent. Soc. V. p. 49.)

"At Chusan, the perfect insect appears during August and September."—(Dr. Cantor's Notes.)

209. *PAPILIO PAMMON*, *Linnæus*.

Papilio Pammon, *Linn.*, *S. N. I.* p. 746 (1767). *Clerck*,
Icon. t. 14, f. 2, ♂. *Cramer*, *Pap.* t. 141, f. B.
Herbst, *Pap.* t. 19, f. 4. *Fabricius*, *Ent. Syst.* III.
pt. I. p. 7. *Esper*, *Ausl. Schmett.* t. 4; f. 1; t. 40,
f. 1. *Godart*, *Enc. Méth.* IX. p. 74. *Boisduval*,
Spéc. Gén. Lép. I. p. 272. *E. Doubleday*, *List Lep.*
Brit. Mus. *pt. I.* p. 10; *Diurnal Lep.* p. 11, n. 62.
G. R. Gray, *Cat. Lep. Brit. Mus.* *pt. I.* p. 19, n. 82;
id. *List Lep. Brit. Mus.* *pt. I.* p. 24.

Laertias Pammon, *Hübner*, *Verz. bek. Schmett.* p. 84.

Princeps Heroicus Pammon, *Hübner*, *Samml. Exot.* I.
t. 108.

Var. A., *G. R. Gray*.

Papilio Cyrus, *Fabricius*, *Ent. Syst.* III. *pt. I.* p. 7
(1793).

Laertias Cyrus, *Hübner*, *Samml. Exot.* II. t. 96.

Papilio Pammon, var., *Stoll*, *Cramer Pap. Suppl.* t. 33,
f. 1.

Var. C.

Papilio Ledebouria, *Eschscholtz*, *in Kotzebue Voy.* III.
t. 3, f. 7.

Papilio Alphenor (*Boisd.*), ♂, *De Haan*, *Verh. Nat.*
Gesch. p. 41.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. N. India. In exchange.

g. h. ♂ ♀. Chusan. Presented by Dr. Cantor, 1842.

i. ♀. N. India. Donor unknown.

The larva and pupa of *Pap. Pammon* are figured on Plate III., fig. 4, 4a, from Java, "and feeds on a species of *Citrus*, bearing the native name of *Juruk*,—November to April. Note.—Every variety of *Juruk*, or Orange (*Gulug*, *Kina*, *Pichet*, &c.), produces a different butterfly, although scarcely any difference appears in the caterpillar."—(Horsfield, MS. Notes.)

R. Templeton, Esq., states that "the male of *P. Pammon* has a

little white mark near the anal angle of the posterior wing; the female an ocellus exactly resembling that of *P. Polytes*; all the other markings are the same in both [sexes,] except in the females they are larger and longer. The larvæ are particularly destructive to the lime-trees in Ceylon, as are also those of *P. Polymnestor*, *Polydorus*, and *Hector*.”—(Ent. Trans. V. p. 44.)

“At Chusan, the perfect insect was observed from July to September, inclusive.”—(Dr. Cantor’s Notes.)

“This is the most common species of the genus, being sometimes seen in dozens in the same field at Rajpore, and elsewhere in the Dhoon, during the months of August and September; nor are they uncommon in the glens of the mountains.”—(Captain T. Hutton, Trans. Ent. Soc. V. p. 50.)

210. *PAPILIO DEMOLION*, Cramer.

Papilio Demolion, Cramer, Pap. I. t. 89, f. A. B. (1775).

Herbst, Pap. t. 49, f. 1, 2. *E. Doubleday*, *Diurnal Lep.* p. 13, n. 94. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 25, n. 116; id. *List Lep. Brit. Mus. pt. I.* p. 34.

Papilio Cresphontes, *Fabricius*, *Ent. Syst.* III. I. p. 33 (1793). *Godart*, *Enc. Méth.* IX. p. 61. *Boisduval*, *Spéc. Gén. Lép.* I. p. 220. *Lucas*, *Lep. Exot. t. 15, f. 1.* *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 4.*

Heraclides Cresphontes, *Hübner*, *Verz. bek. Schmett.* p. 84.

a. b. ♂ ♀. Java. From Dr. Horsfield’s Collection.

c. ♂. Pinang. Presented by Dr. Cantor.

d. e. ♂ ♀. Borneo. Presented by H. Low, Esq.

The larva and pupa of *Pap. Demolion* are figured on Plate III., fig. 5, 5a, from Java; “and feeds on a species of *Fagara*, which bears the native name of *Sampang*,—March and April.”—(Horsfield.)

211. *PAPILIO ERITHONIUS*, Cramer.

Papilio Erithonius, Cramer, Pap. III. t. 232, f. A. B. (1780). *Herbst*, Pap. t. 36, f. 5, 6. *E. Doubleday*, *Diurnal Lep.* p. 12, n. 71. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 21, n. 92; id. *List Lep. Brit. Mus. pt. I.* p. 28.

Princeps Dominans Erithonius, *Hübner, Samml. Exot.* I.
t. 116.

Papilio Epius, *Fabricius, Ent. Syst.* III. I. p. 35 (1793).

Jones, Icon. ined. I. t. 61. *Donovan, Ins. of China*,
t. 29, f. 2. *Godart, Enc. Méth.* IX. p. 43. *Bois-
duval, Spéc. Gén. Lép.* I. p. 238. *E. Doubleday,
List Lep. Brit. Mus. pt.* I. p. 7.

Orpheides Epius, *Hübner, Verz. bek. Schmett.* p. 86.

Papilio Demoleus, *apud Esper, Ausl. Schmett.* t. 50,
f. 1, 4.

a. *Canara.* Presented by S. N. Ward, Esq.

b. c. ♂ ♀. *Calcutta.* Presented by Colonel
Buckley.

d. *Pinang.* Presented by Dr. Cantor.

The larva and pupa are figured on Plate III., fig. 6, 6a, the discovery of which we owe to S. N. Ward, Esq., of the Madras civil service, from whose drawing our figures were copied.

Among the drawings made by Lady Gilbert in India, the transformations of this species are given, and her ladyship states the larva was brought to her on a branch of orange-tree on the 3rd July, and on the 8th it changed to a chrysalis, the perfect insect emerging on the 17th.

"The male of *P. Erithonius* is without the blue lunule, the female has it; and both sexes vary in having or not having one or two spots outside the closing vein of the discoidal cell of the lower wing; beneath, the markings in both are nearly identical, lunule and all." —(R. Templeton, Esq., *Trans. Ent. Soc.* V. p. 44.)

"Occurs in the Deyrah Dhoon, and likewise in the Hills during summer. I have received it from Madras, and frequently took it at Neemuch, in Western India. The larva feeds on the Citron, and is green, with a reddish or orange-coloured head; the fourth segment of the body is also bordered with the same colour, and there is a lateral oblique stripe on the hinder parts, which is blackish and edged with white; the spiracles are black: there are two short tentacular horns projecting from the anterior segment and two others from the anal segment, beneath which latter is a whitish stripe, running obliquely forwards and downwards; a white lateral stripe above the legs, which are yellowish." —(Captain T. Hutton, *Proc. Ent. Soc.* V. p. 48.)

212. *PAPILIO ARJUNA*, Horsfield.

Papilio Arjuna, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 1, f. 4 (1828). *Boisduval*, *Spéc. Gén. Lép.* I. p. 209. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 4; *Diurnal Lep.* p. 11, n. 53. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 17, n. 69; id. *List Lep. Brit. Mus. pt. I.* p. 21.

Papilio Paris, var., *Godart*, *Enc. Méth.* IX. p. 67.

Papilio Paris, *Zinken-Sommer*, *Nova Acta Acad. Nat. Cur.* XV. p. 142.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Pap. Arjuna* are figured on Plate III., fig. 7, 7a, from Java. "Feeds on a species of *Citrus*,—December." —(Horsfield.)

213. *PAPILIO PARIS*, Linnæus.

Papilio Paris, *Linnæus*, *S. N. II.* p. 745 (1767). *Clerck*, *Icones*, t. 13, f. 1. *Cramer*, *Pap.* . t. 103, f. A.B. *Drury*, *Ins. I.* t. 12, f. 1, 2. *Fabricius*, *Ent. Syst. III.* I. p. 1. *Herbst*, *Pap.* t. 14, f. 1, 2. *Esper*, *Ausl. Schmett.* t. 2, f. 1. *Donovan*, *Ins. of China*, t. 23. *Godart*, *Enc. Méth.* IX. p. 67. *Boisduval*, *Spéc. Gén. Lép.* I. p. 208. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 4; *Diurnal Lep.* p. 11, n. 52. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 17, n. 68; id. *List Lep. Brit. Mus. pt. I.* p. 21. *Achillides Paris*, *Hübner*, *Verz. bek. Schmett.* p. 85.

a. b. ♂. Cherra Poonjee. Presented by Colonel Buckley.

c. ♂. Darjeeling. From Pearson's Collection.

d. ♀. N. India. Presented by Lieutenant Hugo James.

e. ♂. Var. Canara. Presented by S. N. Ward, Esq.

214. *PAPILIO GANESA*, E. Doubleday.

Papilio Ganesa, *E. Doubleday*, *Gray's Zool. Misc.* p. 73 (1842); *List Lep. Brit. Mus. pt. I.* p. 4; *Diurnal*

Lep. p. 11, n. 48. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 16, n. 66; id. *List Lep. Brit. Mus.* pt. I. p. 20.

- a. ♂. Darjeeling. From Pearson's Collection.
- b. ♂. N. India. Presented by the Trustees of the British Museum.
- c. d. e. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

215. *PAPILIO ARCTURUS*, *Westwood*.

Papilio Arcturus, *Westwood*, *Ann. N. H.* IX. p. 37 (1842); *Arcana Ent.* t. 27. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 3; *Diurnal Lep.* p. 11, n. 47. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 17, n. 67; id. *List Lep. Brit. Mus.* pt. I. p. 21.

- a. b. ♂. Darjeeling. Presented by Colonel Buckley.
- c. ♀. Darjeeling. From Dr. Pearson's Collection.
- d. ♀. Darjeeling. From the Asiatic Society of Bengal.

216. *PAPILIO KRISHNA*, *Moore* (*Plate IIa.*, fig. 6).

Papilio Krishna, *n. sp.*—UPPER-SIDE black, irrorated with golden-green atoms; *fore-wing* with a well-defined transverse fascia of deep golden-green atoms from anterior margin, commencing at one-third of the length of the wing from the apex to posterior margin near the angle; *hind-wing* with a brilliant sky-blue sinuated patch near anterior angle, and connected with abdominal margin by a broad well-defined sinuous line of deep golden-green atoms; near the margin is a series of five crimson lunules, margined inwardly by a purplish-blue line, the one on the abdominal margin inclosing at the anal angle a black spot, which is bounded at the angle with ferruginous; the space between the sinuous line and patch and the lunules is pure black, without irrorations; the space from the lunules and down middle of tail with golden-green irrorations; indentations margined with white, and tinged inwardly with ferruginous. UNDER-SIDE black; both pair of wings basally irrorated with golden-green and whitish atoms; fascia of *fore-wing* creamy-white; *hind-wing* with creamy-white fascia from anterior margin near angle to middle of the wing, then branching upwards to abdominal margin; lunules

seven in number, brighter and broader than in upper-side; the indentations and anal angle with broad pale ferruginous lunules.

a. ♂. Bootan. From Pemberton's Collection.

b. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

217. *PAPILIO POLYCTOR*, Boisduval.

Papilio Polyctor, *Boisduval*, *Spéc. Gén. Lép.* I. p. 205 (1836). *Blanchard*, *Voy. de Jacquemont*, *Ins. t. 1*, f. 1, 2. *Kollar*, in *Hügel's Kaschmir*, pt. II. p. 403. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 3; *Diurnal Lep.* p. 11, n. 46. *G. R. Gray*, *Cat. Lep. Brit. Mus.* pt. I. p. 16, n. 65; id. *List Lep. Brit. Mus.* pt. I. p. 20.

a. b. c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

218. *PAPILIO CRINO*, Fabricius.

Papilio Crino, *Fabricius*, *Ent. Syst.* III. I. p. 5 (1793). *Donovan*, *Ins. of China*, t. 22; ex *Jones*, *Icon. ined.* I. t. 53. *Godart*, *Enc. Méth.* IX. p. 66. *Boisduval*, *Spéc. Gén. Lép.* I. p. 207. *Guérin*, *Rev. Zool.* (1840), p. 43, t. 1, f. 1. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. *App.* p. 1; *Diurnal Lep.* p. 11, n. 50. *G. R. Gray*, *Cat. Lep. Brit. Mus.* pt. I. p. 18, n. 73; id. *List Lep. Brit. Mus.* pt. I. p. 22.

Papilio Palinurus, var. (?), *De Haan*, *Verh. Natur. Gesch.* pp. 6, 28, t. 7, f. 3.

a. ♂. Ceylon. Donor unknown.

b. ♀. Ceylon. From T. Tatum, Esq.

219. *PAPILIO PERANTHUS*, Fabricius.

Papilio Peranthus, *Fabricius*, *Ent. Syst.* III. I. p. 15 (1793). *Donovan*, *Ins. of China*, t. 26. *Godart*, *Enc. Méth.* IX. p. 66. *Lucas*, *Lep. Exot.* t. 12, f. 2. *Boisduval*, *Spéc. Gén. Lép.* I. p. 203. *De Haan*, *Verh. Natur. Gesch.* p. 27. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 3; *Diurnal Lep.* p. 11,

n. 44. *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 16, *n.* 63; id. *List Lep. Brit. Mus. pt. I.* p. 20.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

220. *PAPILIO BIANOR*, Cramer.

Papilio Bianor, *Cramer, Pap. II. t. 103, f. C.* (1776).

Fabricius, Ent. Syst. III. I. p. 1. *Esper, Ausl. Schmett. t. 35, f. 2.* *Boisduval, Spéc. Gén. Lép. I.* p. 205. *De Haan, Verh. Natur. Gesch.* p. 28, *t.* 5, *f.* 1, ♂; *f.* 2, ♀. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 3; *Diurnal Lep.* p. 11, *n.* 45. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 16, *n.* 64; id. *List Lep. Brit. Mus. pt. I.* p. 20.

Achillides Bianor, *Hübner, Verz. bek. Schmett.* p. 85.

Papilio Paris ♀, *Godart, Enc. Méth. IX.* p. 67.

a. ♀. N. India. Presented by the Trustees of the British Museum.

221. *PAPILIO GYAS*, Westwood.

Papilio Gyas, *Westwood, Arcana Ent. t. 11, f. 1* (1842).

E. Doubleday, List Lep. Brit. Mus. pt. I. p. 3; *Diurnal Lep.* p. 10, *n.* 43. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 16, *n.* 62; id. *List Lep. Brit. Mus. pt. I.* p. 20.

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.

222. *PAPILIO PAYENI*, Boisduval.

Papilio Payeni, *Boisduval, Spéc. Gén. Lép. I.* p. 235 (1836). *Van der Hoeven, Tijdsch. voor Nat. Gesch.* V. *t.* 8, *f.* 1, 2, 6. *E. Doubleday, Diurnal Lep.* p. 14, *No.* 103. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 27, *n.* 125.

Papilio Evan, *E. Doubleday, Ann. Nat. Hist. XVI.* *n. s.* pp. 235, 304 (1845); *List Lep. Brit. Mus. pt. I.* *App.* p. 2; *Diurnal Lep.* p. 14, *No.* 104, *t.* 2, *f.* 1. *Westwood, Cab. Orient. Ent. t. 31, f. 1.*

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

223. *PAPILIO XUTHUS*, Linnæus.

Papilio Xuthus, *Linnæus*, *S. N. II.* p. 751 (1767).
Cramer, *Pap.* I. t. 73, *f. A. B.* *Drury*, *Ins. II.* t. 9,
f. 2. *Fabricius*, *Ent. Syst. III. I.* p. 32. *Herbst*,
Pap. t. 49, *f. 3, 4.* *Godart*, *Enc. Méth. IX.* p. 58.
Boisduval, *Spéc. Gén. Lép. I.* p. 327, *t. 1, f. 1, 2.*
Lucas, *Lep. Exot.* t. 19, *f. 1.* *Herr. Schaeff. Suppl.*
f. 411—413. *De Haan*, *Verh. Natur. Gesch.* p. 41.
E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 15;
Diurnal Lep. p. 16, *n. 157.* *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 36, *n. 178*; *id. List Lep. Brit. Mus. pt. I.* p. 49.

Jasoniades Xuthus, *Hübner*, *Verz. bek. Schmett.* p. 83.

a. b. ♂. N. India. In exchange.

c. ♀. Chusan. Presented by Dr. Cantor, 1844.

The larva of *Pap. Xuthus* is figured on Plate IV., fig. 1, copied from De Haan's figure.

"At Chusan, the perfect insect appears during August and September."—(Dr. Cantor's Notes.)

224. *PAPILIO MACHAON*, Linnæus.

Papilio Machaon, *Linnæus*, *S. N. II.* p. 750 (1767).
Ræsel, *Ins. I. II.* t. 1. *Fabricius*, *Ent. Syst. III. I.* p. 30. *Herbst*, *Pap.* t. 45, *f. 1, 2.* *Hübner*, *Eur. Schmett.* *f. 390, 391.* *Godart*, *Enc. Méth. IX.* p. 57. *Boisduval*, *Spéc. Gén. Lép. I.* p. 328. *G. R. Gray*, *Lep. Ins. of Nepal*, p. 6, *t. 3, f. 1.* *Kollar*, *in Hügel's Kaschmir*, *pt. II.* p. 406. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 15; *Diurnal Lep.* p. 16, *n. 158.* *G. R. Gray*, *Cat. Lep. Brit. Mus. pt. I.* p. 37, *n. 180*; *id. List Lep. Brit. Mus. pt. I.* p. 50.

Pieris Machaon, *Schränk*.

Jasoniades Machaon, *Hübner*, *Verz. bek. Schmett.* p. 83.

Amaryssus Machaon, *Dalm.*, *Königl. Vet. Acad. Holm. XXXVII.* p. 85 (1816).

Var. ♀ *Papilio Machaon*, *De Haan*, *Verh. Natur. Gesch.* p. 42, *t. 5, f. 2.*

Var. *Papilio Sphyrus*, *Hübner*, *Europ. Schmett.* *f. 776, 777.*

- a.* ♂. Himalaya. From the Asiatic Society of Bengal.
- b. c.* ♂ ♀. N. India. Presented by Colonel Buckley.
- d.* ♂. Kumaon. Presented by Dr. McClelland.
- e.* ♀. Bootan. From Dr. Pemberton's Collection.

The transformations of *Pap. Machaon* are copied from Duponchel's work, and are figured on Plate IV., fig. 2, 2a.

Captain T. Hutton remarks: " *P. Machaon* is seen on the wing at Deyrah, in the valley of the Dhoon, as early as February, and in April its larvae are abundant there on the Carrot. At Masuri, in the Hills, it appears in the latter end of March, and continues till October. The larva is green, with a black velvety transverse band across each segment, bearing four spots of bright orange; it possesses the orange-coloured retractile process in the head, from which exudes a liquid drop of a strong aromatic scent, when the insect is touched, precisely as in the European larva. The food is the wild and garden Carrot, and the leaves and flowers of the Radish. I have taken the larvae at Masuri early in May, and the pupa on the 18th May. It is abundant about Simla, and extends far into the interior." —(Ent. Trans. V. p. 47.)

225. PAPILIO CLOANTHUS, Westwood.

Papilio Cloanthus, Westwood, Arcana Ent. p. 42, t. 11, f. 2 (1842). Kollar, in Hügel's Kaschmir, pt. II. p. 405. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 6; Diurnal Lep. p. 14, n. 116. G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 28, n. 136; id. List Lep. Brit. Mus. pt. I. p. 40.

- a. b.* ♂ ♀. N. India. Presented by Colonel Buckley.
- c.* ♀. N. India. From T. Tatum, Esq.

" This is very common in fine warm weather, flitting with great rapidity over the tops of the loftiest trees. It usually selects some lofty oak, over the summit of which it continues to dance with a jerking flight, like that of *P. Sarpedon*, until its domain is invaded by another individual, when a rapid chase round and round the tree takes place: one while they dart away from the tree down the side of the steep mountain, but ever and anon return to the favourite tree, until one is fairly driven off, when the other resumes its dance as before. It is difficult to capture, from its high and rapid flight.

It appears at the end of April, and continues throughout the summer."—(Captain T. Hutton, Tr. Ent. Soc. V. p. 51.)

226. *PAPILIO SARPEDON*, Linnæus.

Papilio Sarpedon, *Linnaeus, S. N. II.* p. 747 (1767). *Ræsel, Ins. IV.* t. 6, f. 1. *Cramer, Pap. II.* t. 122, f. D. *E. Fabricius, Ent. Syst. III.* pt. I. p. 14. *Herbst, Pap. t. 10,* f. 4, 5. *Esper, Ausl. Schmett. t. 8,* f. 2. *Godart, Enc. Méth. IX.* p. 46. *Lucas, Lep. Exot. t. 5,* f. 1. *Boisduval, Spéc. Gén. Lép. I.* p. 235. *De Haan, Verh. Natur. Gesch.* p. 34. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 6; *Diurnal Lep. p. 14,* n. 115. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 28, n. 135; *id. List Lep. Brit. Mus. pt. I.* p. 39.

Zetides Sarpedon, *Hübner, Verz. bek. Schmett.* p. 85; *id. Samml. Exot. III.* t. 471.

Chlorisses Sarpedon, *Scainson, Zool. Ill. 2nd ser. t. 89.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Canara. Presented by S. N. Ward, Esq.

f. N. India. Presented by Colonel Buckley.

The larva and pupa of *Pap. Sarpedon* are figured on Plate III., fig. 8, 8a; the larva copied from De Haan's figure, and the pupa from E. L. Layard's drawing, from Ceylon.

Mr. Edgar L. Layard states that the larva of *Pap. Sarpedon*, as observed in Ceylon, feeds on Cinnamon and Sour-sop.

"At Masuri this is one of the commonest, but not the least beautiful, of our butterflies; it appears early in May, and is found till the end of the rains in September. It usually frequents the top of oak-trees, where it flits about with a jumping or jerking flight, and is somewhat difficult to capture, from its quickness, and the height at which it keeps."—(Captain T. Hutton, Proc. Ent. Soc. V. p. 51.)

227. *PAPILIO EURYPYLUS*, Linnæus.

Papilio Eurypylus, *Linnæus, S. N. II.* p. 754 (1767). *Clerck, Icon. t. 28,* f. 2. *Cramer, Pap. II.* t. 122, f. C. D. *Fabricius, Ent. Syst. III.* I. p. 20. *Herbst, Pap. t. 37,* f. 5, 6. *Esper, Ausl. Schmett. t. 33,* f. 1. *Godart, Enc. Méth. IX.* p. 45. *Boisduval, Spéc.*

Gén. Lép. I. p. 233. *De Haan, Verh. Natur. Gesch.* p. 33. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 6; *Diurnal Lep.* p. 14, n. 113. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 28, n. 133; id. *List Lep. Brit. Mus. pt. I.* p. 38.

Zetides Eurypylius, Hübner, Verz. bek. Schmett. p. 86; id. *Samml. Exot.* II. t. 106.

Papilio Jason, apud Esper, Ausl. Schmett. t. 58, f. 5.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♂. Silhet. Presented by Colonel Buckley.

d. e. ♂ ♀. N. India. From the Asiatic Society of Bengal.

228. *PAPILIO BATHYCLES*, Zinken-Sommer.

Papilio Bathycles, Zinken-Sommer, Nova Acta Acad. Nat. Cur. XV. t. 14, f. 6, 7 (1831). *Boisduval, Spéc. Gén. Lép.* I. p. 232. *Lucas, Lep. Exot.* t. 5, f. 2. *De Haan, Verh. Natur. Gesch.* p. 33. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 6; *Diurnal Lep.* p. 14, n. 111. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 28, n. 132; id. *List Lep. Brit. Mus. pt. I.* p. 38.

a. b. ♀. N. India. Presented by the Trustees of the British Museum.

c. d. ♂ ♀. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

229. *PAPILIO AGAMEMNON*, Linnæus.

Papilio Agamemnon, Linnæus, S. N. II. p. 748 (1767). *Fabricius, Ent. Syst.* III. I. p. 33. *Herbst, Pap. t. 48, f. 1, 2.* *Esper, Ausl. Schmett.* t. 46, f. 1, 2, 3. *Donovan, Ins. of China,* t. 27, f. 2. *Godart, Enc. Méth.* IX. p. 46. *Boisduval, Spéc. Gén. Lép.* I. p. 230. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 5; *Diurnal Lep.* p. 14, n. 109. *G. R. Gray, Cat. Lep. Brit. Mus. pt. I.* p. 27, n. 130; id. *List Lep. Brit. Mus. pt. I.* p. 37.

Papilio Ægistus, vel Agamemnon, Cramer, Pap. t. 106, f. C. D.

Iphicrides Agamemnon, *Hübner, Verz. bek. Schmett.*
p. 82.

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
- c. d. ♂. Silhet. Presented by Colonel Buckley.
- e. ♀. Dukhun. Presented by Colonel Sykes.

The larva and pupa of *Papilio Agamemnon* are figured on Plate III., fig. 9, 9a, from Java, "where it feeds on a species of *Uvaria*, bearing the native name of *Kalak*;—December."—(Horsfield.)

230. *PAPILIO NOMIUS*, Esper.

Papilio Nomius, *Esper, Ausl. Schmett. t. 52, f. 3* (1785).

Boisduval, Spéc. Gén. Lép. I. p. 650. *E. Doubleday, Diurnal Lep. p. 14, n. 119.*

Papilio Orestes, *Fabricius, Ent. Syst. III. I. p. 34* (1793); ex *Jones, Icon. ined. I. t. 79.* *Godart, Enc. Méth. IX. p. 48.* *Boisduval, Spéc. Gén. Lép. I. p. 378.* *Westwood, Arcana Ent. p. 154.* *E. Doubleday, List Lep. Brit. Mus. pt. I. p. 8.* *G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 29,* n. 139; id. *List Lep. Brit. Mus. pt. I. p. 40.*

Papilio Niamus, *Godart, Enc. Méth. IX. p. 51* (1816).
Swainson, Zool. Ill. 2nd ser. t. 32.

Princeps Heroicus *Meges, Hübner, Samml. Exot. I. t. 106.*

Iphicrides Meges, *Hübner, Verz. bek. Schmett. p. 82.*

- a. b. c. ♂ ♀. N. India. From the Asiatic Society of Bengal.

231. *PAPILIO ANTICRATES*, E. Doubleday.

Papilio Anticrates, *E. Doubleday, Ann. N. H. XVIII. p. 371* (1846); *Diurnal Lep. p. 14, n. 120;* *List Lep. Brit. Mus. pt. I. App. p. 2.* *G. R. Gray, Cat. Lep. Brit. Mus. pt. I. p. 29, n. 140, t. 3, f. 3, 3a;* id. *List Lep. Brit. Mus. pt. I. p. 41.*

- a. ♂. N. India. Donor unknown.
- b. c. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

232. *PAPILIO ANTIPHATES*, Cramer.

Papilio Antiphates, Cramer, *Pap.* I. t. 72, f. A. B. (1775). *Fabricius*, *Ent. Syst.* III. I. p. 24; *Jones*, *Icon. ined.* t. 84. *Herbst*, *Pap.* t. 43, f. 1, 2. *Godart*, *Enc. Méth.* IX. p. 49. *Boisduval*, *Spéc. Gén. Lép.* I. p. 248. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 8; *Diurnal Lep.* p. 15, n. 125. *G. R. Gray*, *Cat. Lep. Brit. Mus.* pt. I. p. 31, n. 147; id. *List Lep. Brit. Mus.* pt. I. p. 42.

Iphicrides Antiphates, *Hübner*, *Verz. bek. Schmett.* p. 82. *Papilio Alcibiades*, *Fabricius*, *Ent. Syst.* III. pt. I. (1793), p. 25. *Godart*, *Enc. Méth.* IX. p. 49.

Papilio Pompilius, *Fabricius*, *Ent. Syst.* III. I. p. 25 (1793). *Godart*, *Enc. Méth.* IX. p. 49. *Lucas*, *Lep. Exot.* t. 22, f. 1.

Podalirius Pompilius, *Swainson*, *Zool. Ill. 2nd ser.* t. 105.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. d. ♂. Silhet. Presented by Colonel Buckley.

e. ♀. Borneo. Presented by H. Low, Esq.

The larva and pupa of *Pap. Antiphates* are figured on Plate III., fig. 10, 10 α , from Java, "where it feeds on the same plant as *Pap. Agamemnon*."

233. *PAPILIO AGETES*, Westwood.

Papilio Agetes, *Westwood*, *Arcana Ent.* t. 56, f. 1, 2 (1844). *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 7; *Diurnal Lep.* p. 15, n. 123. *G. R. Gray*, *Cat. Lep. Brit. Mus.* pt. I. p. 31, n. 145; id. *List Lep. Brit. Mus.* pt. I. p. 41.

h. ♀: N. India. Presented by the Trustees of the British Museum.

b. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

234. *PAPILIO GLYCERION*, G. R. Gray.

Papilio Glycerion, G. R. Gray, *Lep. Ins. of Nepal*, p. 6, t. 3, f. 2 (1831). *J. E. Gray's Zool. Misc.* (1832), p. 32. *Boisduval*, *Spéc. Gén. Lép.* I. p. 247. *West-*

wood, Arcana Ent. t. 55, f. 3. *E. Doubleday, List Lep. Brit. Mus.* pt. 1, p. 8; *Diurnal Lep.* p. 15, n. 122. *G. R. Gray, Cat. Lep. Brit. Mus.* pt. I. p. 31, n. 144; id. *List Lep. Brit. Mus.* pt. I. p. 41.

- a. b. ♂ ♀. N. India. Presented by Colonel Buckley.
 c. d. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.
-

REMARKS ON THE CHILOGNATHIFORM, OR JULIFORM, STIRPS.

For the description of the larva and pupa of this Stirps, I refer the reader to the 12th page of the Introductory Remarks.

The Imago, or perfect insect, exhibits in the typical group the most perfect production of the whole order. The beauty of many of the large Papiliones is a subject of universal admiration: several of them are unrivalled in form and colouring.

Linnaeus divides the genus *Papilio* into several phalanges. The first in order of these phalanges are designated *Nobiles*, the last *Plebeii*. The former of these, the *Equites*, is subdivided into—*a*, *Troes*, and *b*, *Achivi*. In these groups he perpetuates in Entomology, by a beautiful allegorical comparison, the names of the most distinguished heroes of Homer's Iliad, besides many other names renowned in ancient history.

Of the character of the wings no description can be given which would apply to the whole stirps. In the Pieridæ, the anterior wings are in general triangular, and the posterior wings short and obtuse. In the typical group, *Papilio*, the posterior wings are slightly falcate on the inner-margin, to give a free passage to the body. In many cases they are provided with tails, which are spatulate or contracted at the base. The *Pap. Machaon*, swallow-tail, is an example of this division.

The palpi, which in the Vermiform stirps are long, slender, and projecting, are in this stirps diversified in the different groups. In the Pieridæ they are of moderate length; and in the genus *Terias*, a diminution of their size is observed; in the typical group, with the exception of the genus *Teinopalpus* of Hope, they are shorter than the head, closely applied to it, and, in the true Papiliones, never exerted.

The antennæ, in the genus *Papilio*, exhibit three subdivisions.

First, antennæ filiform at the base, marked along their entire length with prominent annular rings, and terminated by a cylindrical club, which is attenuated at both ends. This division is exemplified in the following species of this Catalogue, viz.:—

Pap. Memnon (No. 202).	Pap. Demolion (No. 210).
„ Polymnestor (No. 203).	„ Erithonius (No. 211).
„ Helenus (No. 205).	„ Arjuna (No. 212).
„ Polytes (No. 208).	„ Xuthus (No. 223).
„ Pammon (No. 209).	„ Machaon (No. 224).

And by the metamorphoses of these species, viz.:—

Pap. Alcinous, Plate II. f. 6	Pap. Demolion, Plate III. f. 5, 5a
„ Memnon, „ II. f. 7, 7a	„ Erithonius, „ III. f. 6, 6a
„ Polymnestor, „ III. f. 1, 1a	„ Arjuna, „ III. f. 7, 7a
„ Helenus, „ III. f. 2, 2a	„ Xuthus, „ IV. f. 1
„ Polytes, „ III. f. 3, 3a	„ Machaon, „ IV. f. 2, 2a
„ Pammon, „ III. f. 4, 4a	

The larva of this division is characterized by a smooth surface, and by a swelling or intumescence of the fourth or fifth segment of the body, from which it tapers more abruptly to the head, and in a gradual manner to the anal extremity. It is typical of the whole stirps, and represents, *par excellence*, the Chilognathiform division.

In the *second* division of the genus *Papilio*, the antennæ agree in form and outline with those of the first, but the annuli, or rings, along their entire length are comparatively obscure. This division is exemplified in the following species of this Catalogue, viz.:—

Ornithoptera Darsius (No. 176).	Papilio Hector (No. 189).
„ Pompeus (No. 177).	„ Diphilus (No. 190).
Papilio dissimilis (No. 187).	

And the metamorphoses of the same species, which are figured on Plate II., figs. 1, 2, 3, 4, and 5.

The larva of this division is cylindrical, very slightly attenuated at the ends, somewhat thick and fleshy, smooth on the surface, and provided with short obtuse tubercles along its entire length. It represents the Chilopodiform modification of the genus *Papilio*.

In the *third* division of the genus *Papilio*, the antennæ are slender, filiform at the base, but terminated by an oval, comparatively enlarged compressed club, on which the annuli are closely arranged and

crowded. This division is exemplified in the following species of this Catalogue, viz. :—

Papilio Sarpedon (No. 226). | Papilio Agamemnon (No. 229).
Papilio Antipathes (No. 232).

And the metamorphoses of the same species, which are figured on Plate III., figs. 8, 9, and 10.

The larva of this division is smooth, slightly attenuated towards the extremities, somewhat broader or arched in the middle, and marked with regular transverse bands; but its chief character consists in two projecting points from the abdomen. It forms the Thysanuriform division of the genus Papilio.

The three divisions of the genus Papilio which have here been defined from the materials in the Museum, afford an indication of several distinct types of form, which in the larva state have an evident affinity to the Chilognathiform, Chilopodiform, and the Thysanuriform Ametabola; while, in each of these separate groups, the imago manifests a character which reciprocally confirms the distinct character of these divisions.

Whether in collections of greater extent and from more extensive regions, the Vermiform and Anopluriform representatives may be discovered, is a subject which remains for future research.

The general principle on which these divisions have been established, is in perfect consistence with the law above cited; namely, that in groups of consecutive minor value, the same principle is often found to prevail: of which, Messrs. De Haan and Vigors afford ample confirmation.

Without entering on further details, I submit this point to the consideration of Entomologists.—(Horsfield.)

Stirps III., with CHILOPODIFORM or SCOLOPENDRIFORM
Larva.

CHILOPODIFORM or SCOLOPENDRIFORM, *Horsfield, Cat. Lep. Mus. E.I.C. p. 44 (1828).*

PAP. DANAI FESTIVI (*part*), *Linnæus.*

PAP. NYMPHALES, GEMMATI et PHALERATI (*part*), *Linnæus.*

PAP. HELICONII, *Linnæus.* *Fabricius.*

DANAIDES, *Boisduval, Icon. I. p. 84 (1832); id. Spéc. Gén. Lép. I. p. 165 (1836); id. Index Méth. Eur. Lép. p. 15 (1840). Blanchard, Hist. Nat. des Ins. II. p. 330 (1845).*

DANAITES, *Blanchard, Hist. Nat. des Ins. II. p. 330 (1845).*

DANAIDÆ, *E. Doubleday, Diurnal Lep. p. 84 (1847).*

HELICONIDES, *Boisduval, Spéc. Gén. Lép. I. p. 165 (1836).*

HELICONIDÆ, *Westwood, Introd. to Entom. II. p. 351 (1840). E. Doubleday, Diurnal Lep. p. 96 (1847).*

HELICONIITES, *Blanchard, Hist. Nat. des Ins. II. p. 330 (1845).*

ACREIDÆ, *E. Doubleday, Diurnal Lep. p. 137 (1848).*

ARGYNNITES, *Blanchard, Hist. Nat. des Ins. II. p. 331 (1845).*

ARGYNNIDÆ, *Dup. C. p. 2 (1846).*

ARGYNNIDI, *Stephens, List Brit. Lep. Brit. Mus. pt. I. p. 13 (1850). Stainton, Manual Brit. Butt. pp. 21, 41.*

VANESSIDÆ, *Dup. C. p. 6 (1846).*

VANESSIDI, *Stephens, List Brit. Lep. Brit. Mus. pt. I. p. 11 (1850). Stainton, Manual Brit. Butt. pp. 21, 36.*

LEP. TETRAPODA (*part*), *Latreille, Règ. Anim. V. p. 378 (1829).*

NYMPHALIDÆ (*part*), *Swainson; id. Cab. Cyclop. pp. 86, 92 (1840). Stephens, Brit. Ent. Haust. I. p. 27 (1827); id. List Brit. Lep. Brit. Mus. pt. I. p. 6 (1850). Westwood, Introd. to Ent. II. p. 353 (1840). Dup. C. p. 8 (1846). Doubleday et Westwood, Diurnal Lep. p. 144 (1852). Stainton, Manual Brit. Butt. pp. 10, 21 (1856).*

NYMPHALIDES (*part*), *Boisduval, Spéc. Gén. Lép. I. p. 165 (1836); id. Index Méth. Eur. Lép. p. 16. Blanchard, Hist. Nat. des Ins. II. p. 331.*

NYMPHALII et NYMPHALITES (*part*), *Blanchard, Hist. Nat. des Ins. II. pp. 322, 329, 332 (1845).*

NYMPHALIDI (*part*), Stephens, *List Brit. Lep. Brit. Mus. pt. I.* p. 10 (1850). Stainton, *Manual Brit. Butt.* pp. 21, 33 (1856). ? PERIDROMIDÆ, Boisduval, *Spéc. Gén. Lép. I.* p. 164.

Genus DANAI_S, Latreille.

DANAIDA, DANAUS, or DANAI_S, Latreille (1805).

DANAI_S, Godart, *Enc. Méth. IX.* (1819). Boisduval. E. Double-day, *Diurnal Lep.* p. 89 (1847).

AMAUERIS, HESTIA (*part*), EUPLÉA, ANOSIA, Hübner, *Verz. bek. Schmett.* (1816).

235. DANAI_S LIMNIACÆ, Cramer Sp.

Papilio Limniacæ, Cramer, *Pap. I. t. 59, f. C. D.* (1775).

Danais Limniacæ, Godart, *Enc. Méth. IX.* p. 191. E. Double-day, *List Lep. Brit. Mus. pt. I.* p. 49; *Diurnal Lep.* p. 93, n. 31.

Papilio similis, *apud Fabricius*, *Ent. Syst. III.* I. p. 58.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. N. India. From the Asiatic Society of Bengal.

g. ♂. N. India. Presented by Colonel Sykes.

The larva and pupa of *Danais Limniacæ* are figured on Plate IV., fig. 3, 3a, from Java, "where it feeds on a species of *Epibatherium*, bearing the native name of *Tayungan*,—April."—(Horsfield, MS.)

The transformations of this species were also discovered by General Hardwicke.

236. DANAI_S MELISSA, Cramer Sp.

Papilio Melissa, Cramer, *Pap. Exot. IV. t. 377, f. C. D.* (1782). Herbst, *Pap. t. 125, f. 3, 4.*

Danais Melissa, Godart, *Enc. Méth. IX.* p. 192. E. Double-day, *List Lep. Brit. Mus. pt. I.* p. 49; *Diurnal Lep.* p. 92, n. 28.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. N. India. Presented by Lieutenant Hugo James.

f. g. ♂. N. India. Presented by Colonel Buckley.

237. *DANAIS SIMILIS*, Linnæus Sp.

- Papilio similis, *Linnæus*, *Syst. Nat.* I. p. 479 (1758)
 id. *Linn. Syst. Nat.* II. p. 782. *Clerck, Icones*,
 t. 16, f. 3.
Danais similis, *E. Doubleday, List Lep. Brit. Mus. pt. I.*
 p. 49; *Diurnal Lep.* p. 92, n. 30.
Papilio Aventina, *Cramer, Pap. Exot.* I. t. 59, f. F.
 (1775).
Danais Aventina, *Godart, Enc. Méth.* IX. p. 191.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. ♂. N. India. Presented by the Trustees of the British Museum.
f. ♂. Borneo. Presented by H. Low, Esq.

238. *DANAIS GRAMMICA*, Boisduval.

- Danais Grammica*, *Boisduval, Spéc. Gén. Lép.* I. t. 11, f. 10
 (1836). *E. Doubleday, List Lep. Brit. Mus. pt. I.*
 p. 50; *Diurnal Lep.* p. 92, n. 25.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. ♀. N. India. Presented by Colonel Sykes.

239. *DANAIS AGLEA*, Cramer Sp.

- ♂ *Papilio Aglea*, *Cramer, Pap. Exot.* IV. t. 377, f. E.
 (1782).
Danais Aglea, *Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 50; *Diurnal Lep.* p. 92, n. 22.
Danais similis, *apud Godart, Enc. Méth.* IX. p. 190.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. ♂. Bootan. From Pemberton's Collection.
f. g. ♂. N. India. Presented by Colonel Sykes.
h. i. ♀. N. India. Presented by Colonel Buckley.

240. *DANAIS JUVENTA*, Cramer Sp.

- Papilio Juventa*, *Cramer, Pap. Exot.* II. t. 188, f. B.
 (1780).

Danais Juventa, *Godart*, *Enc. Méth.* IX. p. 193. *E.*
Doubleday, *List Lep. Brit. Mus. pt. I.* p. 50;
Diurnal Lep. p. 93, n. 34.

Hestia Juventa, *Hübner*, *Verz. bek. Schmett.* p. 15 (1816).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Borneo. Presented by H. Low, Esq.

The larva and pupa of *Danais Juventa* are figured on Plate IV., fig. 4, 4a, from Java, "and feeds on a plant bearing the native name of *Simbukan-rambat*,—April and May."—(Horsfield, MS.)

241. DANAIS CLEONA, Cramer Sp.

~~distinct~~ ♂ *Papilio Cleona*, *Cramer*, *Pap. Exot. IV.* t. 377, f. F. (1782).

Danais Cleona, *Godart*, *Enc. Méth.* IX. p. 190. *E.*
Doubleday, *List Lep. Brit. Mus. pt. I.* p. 50; *id.*
Diurnal Lep. p. 92, n. 27.

Hab. Moluccas (*Cramer*).

Var. *Euploea Philomela*, *Zinken-Sommer*, in *Nova Acta Acad. Curios.* XV. t. 16, f. 17 (1831).

Danais Dorippa, *Boisduval*, *MS.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Borneo. Presented by H. Low, Esq.

242. DANAIS MELANEUS, Cramer Sp.

Papilio Melaneus, *Cramer*, *Pap. Exot. I.* t. 30, f. D. (1775).

Danais Melaneus, *Godart*, *Enc. Méth.* IX. p. 192. *E.*
Doubleday, *List Lep. Brit. Mus. pt. I.* p. 50;
Diurnal Lep. p. 93, n. 36.

a. b. ♂. N. India. Presented by Lieutenant Hugo James.

c. ♀. Pinang. Presented by Dr. Cantor.

d. ♂. Java. From Mr. Argent's Collection.

243. DANAIS TYTIA, G. R. Gray Sp.

Euploea Tytia, *G. R. Gray*, *Lep. Ins. of Nepal*, p. 9, t. 9, f. 2 (1846).

Danais Tytia, *E. Doubleday, List Lep. Brit. Mus. pt. I.*
p. 50; Diurnal Lep. p. 93, n. 35, t. 12, f. 4.

- a. ♂. Bootan. From Pemberton's Collection.
- b. c. ♀. Darjeeling. From Pearson's Collection.
- d. ♂. N. India. Presented by Colonel Buckley.
- e. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

244. *DANAIS ABLATA*, Zinken-Sommer Sp.

Euplœa ablata, *Zinken-Sommer, in Nova Acta Acad. Curios. XV. t. 16, f. 16* (1831).

Danais ablata, *E. Doubleday, List Lep. Brit. Mus. pt. I.*
p. 50; Diurnal Lep. p. 93, n. 37.

- a. ♀. Java. From Dr. Horsfield's Collection.
- b. c. d. ♂ ♀. Java. From Mr. Argent's Collection.

245. *DANAIS PHILENE*, Cramer Sp.

Papilio Philene, *Cramer, Pap. Exot. IV. t. 375, f. A. B.*
(1782).

Danais Philene, *Godart, Enc. Méth. IX. p. 187. E.*
Doubleday, List Lep. Brit. Mus. pt. I. p. 50;
Diurnal Lep. p. 92, n. 17.

Euplœa Philene, *Hübner, Verz. bek. Schmett. p. 15.*

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Danais Philene* are figured on Plate IV., fig. 5, 5a, from Java, "where it feeds on a species of *Cissus*, bearing the native name of *Galing*,—December."—(Horsfield, MS.)

246. *DANAIS PLEXIPPUS*, Linnæus Sp.

Papilio Plexippus, *Linnæus, Syst. Nat. II. p. 767* (1767).
Fabricius, Ent. Syst. III. I. p. 49.

Danais Plexippus, *Godart, Enc. Méth. IX. p. 186. E.*
Doubleday, List Lep. Brit. Mus. pt. I. p. 51;
Diurnal Lep. p. 92, n. 16.

Euplœa Plexippus, *Hübner, Verz. bek. Schmett. p. 15.*

Papilio Genutia, *Cramer, Pap. Exot. III. t. 206, f. C.D.*
(1780).

- a. b. ♂.* N. India. From the Asiatic Society of Bengal.
- c. ♂.* Assam. From Dr. McClelland's Collection.
- d. ♂.* Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

The larva and pupa of *Danais Plexippus* are figured on Plate IV., fig. 6, 6a, copied from General Hardwicke's drawings.

247. *DANAIS MELANIPPUS*, Cramer Sp.

Papilio Melanippus, Cramer, Pap. Exot. II. t. 127, f. A. B. (1777).

Danais Melanippus, Godart, Enc. Méth. IX. p. 189. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 50.

- a. b. c. d. ♂ ♀.* Java. From Dr. Horsfield's Collection.

Var. Papilio Hegesippus, Cramer, Pap. Exot. II. t. 180, f. A. (1779). Fabricius, Ent. Syst. III. I. p. 52.

Euploea Hegesippus, Hübner, Verz. bek. Schmett. p. 15.

Danais Hegesippus, Godart, Enc. Méth. IX. p. 189. G. R. Gray, Lep. Ins. of Nepal, p. 10, t. 9, f. 1. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 50; id. Diurnal Lep. p. 92, n. 19.

Danais Chirona, G. R. Gray, Lep. Ins. of Nepal, p. 10 (1833).

e. f. g. ♀. N. India. Presented by Colonel Buckley.

h. ♂. Pinang. Presented by Dr. Cantor.

248. *DANAIS LOTIS*, Cramer Sp.

Papilio Lotis, Cramer, Pap. Exot. III. t. 230, f. D. E. (1780).

Danais Lotis, Godart, Enc. Méth. IX. p. 189. E. Doubleday, Diurnal Lep. p. 92, n. 21; List Lep. Brit. Mus. App. 1848, p. 13.

Hestia Thoë, Hübner, Verz. bek. Schmett. p. 15 (1816).

Danais Edmondi, Bougainville, Voy. de la Corvette Thétis, t. 44, f. 3, 3 bis (1837).

a. ♂. Borneo. Presented by Hugh Low, Esq.

249. *DANAIS CHRYSIPPUS*, Linnæus Sp.

Papilio Chrysippus, Linnæus, *Syst. Nat.* II. p. 767 (1767). Cramer, *Pap. Exot.* II. t. 118, f. B. C. Fabricius, *Ent. Syst.* III. I. p. 50.

Danais Chrysippus, Godart, *Enc. Méth.* IX. p. 187. E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 51; *Diurnal Lep.* p. 92, n. 14.

Euplœa Chrysippus, Hübner, *Verz. bek. Schmett.* p. 15.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. Darjeeling. From Pearson's Collection.

g. h. ♂ ♀. Madras. In exchange from Captain J. M. Jones.

The larva and pupa of *Danais Chrysippus* are figured on Plate IV., fig. 7, 7a, from Java, "where it feeds on *Asclepias gigantea*, the native name of which is *Widuri*,—February and March. Not common."—(Horsfield, MS.)

Genus EUPLŒA, Fabricius.

EUPLŒA, *Fabricius* (*Illiger's Mag.* 1807). Horsfield. Boisduval. Doubleday, *Diurnal Lep.* p. 86.

TERPSICHROIS, *Crastia*, *Salpinx*, *Didonis*, Hübner, *Verz. bek. Schmett.* (1816).

DANAUS, *p.* Latreille.

DANAIS, *p.* Godart.

250. *EUPLŒA RHADAMANTHUS*, Fabricius Sp.

♂ Papilio Rhadamanthus, *Fabricius*, *Ent. Syst.* III. I. p. 42 (1793). Jones, *Icon. t. 45, f. 2.*

Euplœa Rhadamanthus, E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 48; *Diurnal Lep.* p. 88, n. 23.

♀ Papilio Diocletianus, *Fabricius*, *Ent. Syst.* III. I. p. 40.

♀ Danais Diocletianus, Godart, *Enc. Méth.* IX. p. 181.

♂ Terpsichrois Thoösa, Hübner, *Samml. Exot. Schmett.* (1806–27).

♂ Danais Aleidice, Godart, *Enc. Méth.* IX. p. 181 (1819).

♂ Danais Rhadamia, Godart, *Enc. Méth.* IX. p. 180 (1819).

- a. ♂. Java. From Dr. Horsfield's Collection.
- b. c. ♂. N. India. Presented by Lieut. Hugo James.
- d. ♂. Cherra Poonjee. Presented by Colonel Buckley.
- e. ♀. Pinang. Presented by Dr. Cantor.
- f. ♂. Borneo. Presented by Hugh Low, Esq.

251. *EUPLOEA MITRA*, Moore.

Euploea Mitra, Moore, *Trans. Ent. Soc.* 1857, p. ,
t. , f. .

- a. ♂. Habitat and donor not preserved.

Euploea Mitra, n. sp.—UPPER-SIDE pale chocolate-brown; fore-wing with a broad oblique transverse white fascia from anterior margin (beyond discoidal cell) to near middle of exterior margin, and there terminated by a separated round white spot; three small white spots near the apex of the wing, and a sub-marginal row of three or four rather indistinct spots: hind-wing with the anterior portion glaucous or silvery-whitish (somewhat as in *Euploea Rhadamanthus*). UNDER-SIDE slightly paler; fore-wing tinged with glaucous posteriorly; fascia as in upper-side, and having a sub-marginal row of ten or twelve white spots, and a longitudinal oval mark of dull brown between the sub-median vein and median veinlet; hind-wing with two rows of sub-marginal white spots, and some spots about the disc, two being within discoidal cell. Expanse of wings $2\frac{3}{4}$ inches.

252. *EUPLOEA ELUSINE*, Cramer Sp.

Papilio Elusine, Cramer, *Pap. Exot.* III. t. 266, f. D. (1780).

Euploea Elusine, Doubleday, *List Lep. Brit. Mus.* pt. I. p. 47; *Diurnal Lep.* p. 87, n. 9.

Terpsichrois Elusine, Hübner, *Verz. bek. Schmett.* p. 16 (1816).

Danais Elusine, Godart, *Enc. Méth.* IX. p. 177 (1819).

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

253. *EUPLOEA MAZARES*, E. Doubleday, MS.

Euploea Mazares, E. Doubleday, *Diurnal Lep.* p. 87, No. 10 (1847). Moore, *Trans. Ent. Soc.* 1857, p. , t. , f. .

Salpinx Elusine, *apud Hübner, Samml. Exot. Schmett.*
(1806-27).

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

Euplœa Mazares.—Male: UPPER-SIDE dark brown, paler on hind-wings; *fore-wing* with a brilliant blue gloss; a sub-marginal curved row of bluish-white spots from anterior margin near the apex, the spots lessening in size downwards; also a rather indistinct bluish spot on anterior margin near the middle: *hind-wing* with a creamy-white patch in anterior half, near the middle, and two small white spots near anterior angle. UNDER-SIDE paler brown, with a sub-marginal row of white spots, and also a marginal row of very minute spots: a costal white spot; a larger white spot near the discoidal cell, between the second and third median veinlets; also a creamy-white patch near posterior margin: *hind-wing* with a marginal half-row of minute white spots, commencing from anal angle, also a sub-marginal row curving from middle of anterior margin; some spots also at the base of the wings. Female as in male, but paler; the blue gloss less vivid, and the creamy-white patches obsolete. Shape of wings as in *Euplœa Elusine*. Expanse $2\frac{1}{4}$ inches.

254. *EUPLŒA MELINA*, Godart Sp.

Danais Melina, Godart, *Enc. Méth. IX.* p. 179 (1819).

Euplœa Melina, Boisduval, *Faune de l'Océanie*, p. 98.

E. Doubleday, Diurnal Lep. p. 88, n. 17.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

255. *EUPLŒA HÜBNERI*, Boisduval MS.

Euplœa Hübneri, Moore, *Trans. Ent. Soc.* 1857, p. ,
t. , f. :

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

Euplœa Hübneri, n. sp.—Male: UPPER-SIDE dark velvety brown, palest exteriorly: *fore-wing* with a semicircular row of six small white spots, commencing from anterior margin near apex, and curving outwards; also a less distinct white spot near discoidal cell, between first and second median veinlets; an impressed dull-brown longitudinal streak between sub-median vein and median veinlet: *hind-wing* with only a row of three spots near anterior angle. UNDER-

SIDE pale brown: *fore-wing* with the semicircular row of spots, and a row outside discoidal cell, and one spot within it, that, and the lowest spot, being the largest: *hind-wing* with two short rows of spots near anterior angle, the outer row the smallest; and small spots outside and one within, discoidal cell; some spots also at the base of wings. Female: UPPER-SIDE paler brown: *fore-wing* with only three spots near apex, the indistinct spot near discoidal cell, and impressed brown mark: *hind-wing* without spots. UNDER-SIDE paler: *fore-wing* with spots near apex, as above; also one on middle of costal margin, one within and two outside discoidal cell, one being between each median veinlet; also the longitudinal streak, which is white: *hind-wing* with sub-marginal row of very small spots, also a row outside, and one spot within, discoidal cell. Wings shaped as in *Euplœa Elusine*. Expanse 3½ inches.

256. *EUPLŒA CRAMERI*, Boisduval MS.

Euplœa Crameri, Moore, Trans. Ent. Soc. 1857, p. .
t. , f. .

a. ♂. Borneo. Presented by Hugh Low, Esq.

Euplœa Crameri, n. sp.—UPPER-SIDE wholly velvety blackish-brown, having a bluish tint, and slightly paler on exterior margins: *fore-wing* with a short transverse row of six white spots from anterior margin, close to apex, the two first small, the third large and longest, fourth same size as first, fifth and sixth very small; also a small spot near discoidal cell, between first and second median veinlets: *hind-wing* with two small white spots near anterior angle. UNDER-SIDE clear deep brown; *fore-wing* with white spots, as in upper-side, also with three spots along costal margin, and some on the disc: *hind-wing* with a sub-marginal row of small white spots, and some on the disc. Expanse of wings 3½ inches.

257. *EUPLŒA EUNICE*, Godart Sp.

Danais Eunice, Godart, Enc. Mith. IX. p. 177 (1819).
Euplœa Eunice, Boisduval, Faune de l'Océanie, p. 94 (1832). Doubleday, List Lep. Brit. Mus. pt. I. p. 47; Diurnal Lep. p. 87, n. 1.

? Limnas mutabilis Nemertes, Hübner, Samml. Exot. Schmett. (1806–27).

Salpinx Nemertes, Hübner, Verz. bek. Schmett. p. 17 (1816).

- a. ♂. Pinang. Presented by Dr. Cantor.
- b. ♂. Java. Presented by the Trustees of the British Museum.

258. *EUPLEA KLUGII*, Boisduval MS.

Euplea Klugii, Moore, *Trans. Ent. Soc.* 1857, p. ,
t. , f. .

- a. b. c. ♂ ♀. Bootan. From Dr. Pemberton's Collection.
- d. ♂. N. India. Presented by Colonel Sykes.

Euplea Klugii, n. sp.—Male: UPPER-SIDE deep brown, having on the *fore-wings* a brilliant blue gloss, a sub-marginal row of small bluish-white spots, and an inner parallel row of larger spots, also a bluish spot on costal margin; one within discoidal cell; two linear bluish marks, one between each discoidal veinlet; and a rather indistinct bluish mark between sub-median vein and median veinlet: *hind-wing* paler brown, darkest and glossed with blue in the middle; a sub-marginal row of white spots and a short inner row from anterior margin; also a patch of creamy-white near middle of the wing. UNDER-SIDE brown; *fore-wing* with the two rows of white spots smaller; one spot on costal margin, one small narrow spot between second discoidal and first median veinlets, and a larger spot between second and third median veinlets; a creamy-white patch on posterior base of the wing: *hind-wing* with two rows of white spots. Female: UPPER-SIDE nearly as in male, the *fore-wing* with the sub-marginal row of spots obsolete; *hind-wing* pale brown, darkest and slightly glossed with blue in the middle; two white spots only of inner row distinct, the rest of two rows indistinct; without the creamy-white patch. UNDER-SIDE nearly as in male, but not having the creamy-white patch. Shape of wings as in *Euplea superba*, Herbst. Expanse $3\frac{5}{8}$ inches.

259. *EUPLEA PROTHOË*, Godart Sp.

Danais Prothoë, Godart, *Enc. Méth.* IX. p. 177
(1819).

Euplea Prothoë, E. Doubleday, *List Lep. Brit. Mus.*
pt. I. p. 47; *Diurnal Lep.* p. 87, n. 12.

♀ *Papilio Midamus*, apud Cramer, *Pap. Exot.* III. t. 266,
f. A. B. (1780).

Euplœa Pavettæ, Zinken-Sommer, in *Nova Acta Acad. Nat. Curios.* XV. p. 189 (1831).

? *Terpsichrois Alea*, Hübner, *Verz. bek. Schmett.* p. 16 (1816).

a. ♂. Java. Presented by the Trustees of the British Museum.

The larva and pupa of *Euplœa Prothoë* are figured on Plate IV., fig. 8, 8a, from Ceylon, where discovered by E. L. Layard, Esq., from whose drawings our figures were copied.

260. *EUPLOEA SUPERBA*, Herbst Sp.

Papilio superba, Herbst, Pap. t. 119, 120 (1783-95).

Euplœa superba, E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 47; *Diurnal Lep.* p. 87, n. 14.

Danais Alopia, Godart, *Enc. Méth.* IX. p. 177 (1819).

Limnas mutabilis Midamis, Hübner, *Sammel. Exot. Schmett.* (1806-27).

a. ♂. Darjeeling. From Pearson's Collection.

b. ♂. China. Presented by Dr. Cantor.

c. ♀. China. In exchange.

261. *EUPLOEA ALCATHOË*, Godart Sp.

Danais Alcathoë, Godart, *Enc. Méth.* IX. p. 178 (1819).

Euplœa Alcathoë, Boisduval, *Faune de l'Océanie*, p. 99. E. Doubleday, *Diurnal Lep.* p. 87, n. 16.

Euplœa Anthrax, Boisduval MS.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Silhet. Presented by the Trustees of the British Museum.

f. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

262. *EUPLOEA CORE*, Cramer Sp.

Papilio Core, Cramer, Pap. Exot. III. t. 266, f. E. F. (1780).

Euplœa Core, E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 49; *Diurnal Lep.* p. 88, n. 31.

Crastia Core, *Hübner*, *Verz. bek. Schmett.* p. 16 (1816).
Papilio Corus, *Fabricius*, *Ent. Syst.* III. I. p. 41 (1793).
Danais Coreta, *Godart*, *Enc. Méth.* IX. p. 182 (1819).

a. b. c. ♂. N. India. Presented by Colonel Buckley.
d. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

The larva and pupa of *Euplæa Core* are figured on Plate IV., fig. 9, 9a, which were discovered in Canara by S. N. Ward, Esq., and from whose drawings our figures were copied.

263. EUPLEA DEIONE, Westwood.

Euplæa Deione, *Westwood*, *Cab. Orient. Entom.* p. 76, t. 37, f. 3 (1847).

a. ♂. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

264. EUPLEA OCHSENHEIMERI, Boisduval MS.

Euplæa Ochsenheimeri, *Moore*, *Trans. Ent. Soc.* 1857,
p. , t. , f. .

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. ♀. Java. Presented by the Trustees of the British Museum.

Euplæa Ochsenheimeri, *n. sp.*—Male: UPPER-SIDE silky brown, darkest on the hind-wings, and having a golden-greenish metallic lustre in some lights: *fore-wing* with a sub-marginal row of small white spots (two between each veinlet); two other parallel rows of somewhat larger spots (one between each veinlet in each row); also one spot between first and second branches of subcostal vein, and one within the discoidal cell; narrow ciliæ between indentations spotted with white: *hind-wing* with two short rows of four very small white spots near anterior angle; narrow ciliæ spotted with white. UNDER-SIDE as above, but the hind-wing having the sub-marginal row of white spots continued along the whole margin; also a row outside discoidal cell; some white spots also at the base of the wings. Female paler, the rows of white spots on upper-side of *fore-wing* disposed exactly the same, but are larger, those on the *hind-wing* more straggling, and but few being distinct. UNDER-SIDE as above, the

fore-wing having also a longitudinal club-shaped streak between submedian vein and median veinlet : *hind-wing* with three rows of white spots, also one spot within discoidal cell, and some spots disposed at base of the wings. Shape of wings as in *Euplœa Midamus*. Expanse of wings $3\frac{5}{8}$ inches.

265. EUPLŒA MIDAMUS, Linnæus Sp.

Papilio Midamus, *Linnæus, Syst. Nat.* II. p. 756 (1767).

Fabricius, Ent. Syst. III. I. p. 39.

Euplœa Midamus, *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 47 ; *Diurnal Lep.* p. 87, n. 11.

Terpsichrois Midamus, *Hübner, Verz. bek. Schmett.* p. 16 (1816).

Danaïs Midamus, *Godart, Enc. Méth.* IX. p. 179.

Papilio Muleiber, *Cramer, Pap. Exot.* II. 127, f. C.D. (1776).

♀ Papilio Claudius, *Fabricius, Ent. Syst.* III. I. p. 40 (1793).

Danaïs Claudia, *Godart, Enc. Méth.* IX. p. 180.

♀ Papilio Basilissa, *Cramer, Pap. Exot.* III. t. 266, f. C. (1780).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Darjeeling. From Pearson's Collection.

g. Pinang. Presented by Dr. Cantor.

k. ♂. Borneo. Presented by H. Low, Esq.

The larva and pupa of *Euplœa Midamus* are figured on Plate IV., fig. 10, 10a, from Java, where it "feeds on a species of *Ficus*,—December."—(Horsfield, MS.)

Genus IDEOPSIS, Horsfield.

DANAIIS (*Sect. IV.*), *E. Doubleday, Diurnal Lep.* p. 93 (1847).

"Closely resembles *Hestia* in form, texture, and colouring of the wings, and to which it has another resemblance in the absence of the sexual spot on the posterior wings. Like some species of the third section of *Danaïs*, and like the genus *Hestia*, it has the first subcostal veinlet anastomosing with the costal vein. Like most species of *Hestia*, the genus *Ideopsis* has the wings somewhat diaphanous, white; the outer margin, veins, veinlets, two or more vittæ in the cell,

and a series of dots between the veinlets sometimes coalescing, all fuscous: but, notwithstanding these points of resemblance, it may always be known from *Hestia* at first sight by its distinctly clavate antennæ, and on closer examination, by its claws devoid of paronychia or pulvilli."—(Doubleday.)

266. IDEOPSIS GAURA, Horsfield.

Idea ? Gaura, *Horsfield, Cat. Lep. Mus. E.I.C. t. 6, f. 1* (1829). *Boisduval, Spéc. Gén. Lép. t. 11, f. 11.*

Hestia Gaura, *E. Doubleday, List Lep. Brit. Mus. pt. I. p. 52.*

Danais Gaura, *E. Doubleday, Diurnal Lep. p. 93, n. 38.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

267. IDEOPSIS DAOS, Boisduval Sp.

Idea Daos, *Boisduval, Spéc. Gén. Lép. I. t. 24, f. 3* (1836).

Hestia Daos, *E. Doubleday, List Lep. Brit. Mus. pt. I. p. 52.*

Danais Daos, *E. Doubleday, Diurnal Lep. p. 92, n. 39.*

Hestia Eudora, *G. R. Gray, in Cat. Lep. Ins. of Nepal, p. 10, t. 9, f. 3* (1846).

a. ♂. Pinang. Presented by Dr. Cantor.

The larva and pupa of *Ideopsis Daos* are figured on Plate IV., fig. 11, 11a, for which we are indebted to J. O. Westwood, Esq., and which were discovered by Captain Hamilton on the Tenasserim coast.—(Vide Trans. Ent. Soc. I. n. s. pp. 35, 36.)

Genus HESTIA, Hübner.

HESTIA, *Hübner, Verz. bek. Schmett. p. 15* (1816). *E. Doubleday, Diurnal Lep. p. 94.*

IDEA, *Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 283, 1808).* *Latreille. Godart. Boisduval.*

268. HESTIA LYNCEUS, Drury Sp.

Papilio Lynceus, *Drury, Ins. II. t. 7, f. 1* (1773).

Hestia Lynceus, *E. Doubleday, Diurnal Lep. p. 95, n. 1, t. 13, f. 1* (*H. Idea, var. on plate*).

Idea Lynceus, *Godart, Enc. Méth.* IX. p. 195.

Papilio Idea, *Stoll, Suppl. à Cramer, Pap. Exot.* V. t. 42, f. 1 (1791).

a. ♂. Malacca. Presented by Colonel Buckley.

269. *HESTIA BELIA*, *Westwood*.

Hestia Belia, *Westwood, Cab. Orient. Ent.* p. 75, t. 37, f. 2 (1848).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The pupa of *Hestia Belia* is figured on Plate IV., fig. 12, drawn from a specimen taken in Java (from which the imago emerged) by Dr. Horsfield.

270. *HESTIA JASONIA*, *Westwood*.

Hestia Jasonia, *Westwood, Cab. Orient. Ent.* p. 87, t. 42, f. 1 (1848).

a. ♀. Canara (South India). Presented by S. N. Ward, Esq.

271. *HESTIA LEUCONOË*, *Erichson*.

Hestia Leuconoë, *Erichson, Nova Acta Acad. Nat. Cur. XVI.* p. 283 (1834). *E. Doubleday, Diurnal Lep.* p. 95, n. 4, t. 13, f. 2.

a. b. ♀. Borneo. Presented by Hugh Low, Esq.

Genus *TELCHINIA*, *Hübner*.

TELCHINIA, *Hübner, Verz. bek. Schmett.* p. 27 (1816).

ACRÆA (*TELCHINIA*), *E. Doubleday, Diurnal Lep.* p. 141 (1848).

272. *TELCHINIA VIOLÆ*, *Fabricius Sp.*

Papilio Violæ, *Fabricius, Syst. Ent.* p. 460 (1775); *Ent. Syst. III. I.* p. 164 (1793).

Acroæa Violæ, *Godart, Enc. Méth.* IX. p. 231. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 62.

Acroæa (*Telchinia*) *Violæ*, *E. Doubleday, Diurnal Lep.* p. 142.

Papilio Cephea, *Cramer, Pap. Exot. III. t. 298, f. D.E.* (1782).

Telchinia Cephea, *Hübner*, *Verz. bek. Schmett.* p. 27.

a. b. c. N. India. From the Asiatic Society of Bengal.

The larva and pupa of *Telchinia Violæ* are figured on Plate V., fig. 1, 1 α , copied from General Hardwicke's drawings in the British Museum.

Genus PAREBA, *E. Doubleday*.

ACRÆA (PAREBA), *E. Doubleday*, *Diurnal Lep.* p. 142 (1848).

273. *PAREBA VESTA*, *Fabricius Sp.*

Papilio Vesta, *Fabricius*, *Man. Ins.* II. p. 14 (1787); *Ent. Syst.* III. I. p. 163 (1793).

Aeræa Vesta, *Godart*, *Enc. Méth.* IX. p. 233. *E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 63.

Aeræa (Pareba) Vesta, *E. Doubleday*, *Diurnal Lep.* p. 142. *Papilio Terpsichore*, *Cramer*, *Pap. Exot.* III. t. 228, f. A.B.C. (1782).

Telchinia Issoria, *Hübner*, *Verz. bek. Schmett.* p. 27 (1816).

Var. ♀ *Aeræa anomala*, *Kollar*, *in Hügel's Kaschmir*, IV. pt. II. p. 425, t. 3, f. 3, 4 (1845).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. Darjeeling. From Pearson's Collection.

g. ♂. Bootan. From Pemberton's Collection.

h. ♀. N. India. From the Asiatic Society of Bengal.

i. ♀. (*A. anomala*, *Kollar*). N. India. Presented by Colonel Buckley.

The larva and pupa of *Pareba Vesta* are figured on Plate V., fig. 2, 2 α , from Java, "where it feeds on a species of *Urtica*, bearing the native name of *Latungan*,—December."—(Horsfield, MS.)

Genus VANESSA, *Fabricius*.

VANESSA, *Fabricius* (*Illiger's Mag.* VI. 1808). *E. Doubleday*, *Diurnal Lep.* p. 198.

EUGONIA; INACHIS, *Hübner*, *Verz. bek. Schmett.* (1816).

274. *VANESSA ANTIOPA*, *Linnæus Sp.*

Papilio Antiopa, *Linnæus*, *Syst. Nat.* II. p. 766 (1767).

Vanessa Antiopa, *Ochsenheimer*, *Schmett. von Eur.* IV.
p. 16. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.*
p. 80; *Diurnal Lep. p. 200, n. 10.*

Eugonia Antiopa, *Hübner*, *Verz. bek. Schmett.* p. 37.

a. ♀. Bootan. From Pemberton's Collection.

275. *VANESSA XANTHOMELAS*, *Denis et Schiff. Sp.*

Papilio xanthomelas, *Denis et Schiffermüller*, *Wien. Verz.* p. 175 (1776). *Hübner*, *Samml. Europ. Schmett. Pap. f. 85, 86.*

Vanessa xanthomelas, *Ochsenheimer*, *Schmett. von Eur.* IV. p. 17. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 81; *Diurnal Lep. p. 200, n. 4.*

a. ♀. Darjeeling. Presented by Colonel Buckley.

276. *VANESSA CASHMIRENSIS*, *Kollar.*

Vanessa Cashmirensis, *Kollar*, in *Hügel's Kaschmir*, IV.
pt. II. p. 442, t. 9, f. 3, 4 (1844). *E. Doubleday*,
Diurnal Lep. p. 200, n. 5.

Vanessa Flacilla, *E. Doubleday*. *G. R. Gray*, *Lep. Ins. of Nepal*, p. 11 (1846).

a. b. c. ♂ ♀. Bootan. From Pemberton's Collection.

d. ♂. N. India. Presented by Colonel Buckley.

e. ♀. Ladakh. Presented by Captain Strachey.

277. *VANESSA CHARONIA*, *Drury Sp.*

Papilio Charonia, *Drury*, *Ins. I. t. 15, f. 1, 2* (1770).
Cramer, *Pap. Exot. I. t. 47, f. A.B.C.* *Fabricius*,
Ent. Syst. III. I. p. 119.

Vanessa Charonia, *Godart*, *Enc. Méth. IX. p. 308.* *Kollar*,
in Hügel's Kaschmir, IV. pt. II. p. 442. *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 80;* *Diurnal Lep. p. 201, n. 12.*

a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.

d. ♀. N. India. From Mr. Argent's Collection.

Genus GRAPTA, *Kirby.*

GRAPTA, *Kirby*, *Fauna Bor. Amer.* p. 292 (1837).

POLYGONIA, *p.*; *EUGONIA*, *p.* *Hübner.*

278. *GRAPTA C. AUREUM*, Linnæus Sp.

Papilio C. aureum, *Linnæus*, *Syst. Nat.* II. p. 778 (1767).

Grapta C. aureum, *E. Doubleday*, *Diurnal Lep.* p. 197, n. 1.

Vanessa C. aureum, *Godart*, *Enc. Méth.* IX. p. 304. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 81.

Papilio Angelica, *Cramer*, *Pap. Exot.* IV. t. 388, f. G. H. (1782).

Eugonia Angelica, *Hübner*, *Verz. bek. Schmett.* p. 26.

a. ♂. Pinang. Presented by Dr. Cantor.

Genus PYRAMEIS, *Hübner*.

PYRAMEIS, *Hübner*, *Verz. bek. Schmett.* 1816. *E. Doubleday*, *Diurnal Lep.* p. 202 (1849).

VANESSA, p. *Latreille*. *Godart*. *Hübner*.

CYNTHIA, p. *Fabricius*.

279. *PYRAMEIS CALLIRHOË*, *Hübner*.

Hamadryas Callirhoë, *Hübner*, *Samml. Exot. Schmett.* (1806–16).

Pyrameis Callirhoë, *Hübner*, *Verz. bek. Schmett.* p. 33 (1816). *E. Doubleday*, *Diurnal Lep.* p. 204.

Vanessa Callirhoë, *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 78. *G. R. Gray*, *Lep. Ins. of Nepal*, p. 11.

Vanessa Vulcania, *Godart*, *Enc. Méth.* IX. p. 320 (1819). *Kollar*, *in Hügel's Kaschmir*, IV. pt. II. p. 442.

Papilio Atalanta, *Cramer*, *Pap. Exot.* I. t. 84, f. E. F.

a. b. c. ♂ ♀. Darjeeling. From Pearson's Collection.

d. ♀. N. India. Presented by Colonel Hearsey.

e. ♂. Ladakh. Presented by Captain Strachey.

f. ♀. Pinang. Presented by Dr. Cantor.

280. *PYRAMEIS CARDUI*, Linnæus Sp.

Papilio Cardui, *Linnæus*, *Syst. Nat.* II. p. 774 (1767).

Pyrameis Cardui, *E. Doubleday*, *Diurnal Lep.* p. 205, n. 7.

Vanessa Cardui, *Hübner*, *Verz. bek. Schmett.* p. 73.
Godart, *Enc. Méth.* IX. p. 323. *E. Doubleday*,
List Lep. Brit. Mus. pt. I. p. 79.

Cynthia Cardui, *Stephens*, *Ill. Haust.* I. p. 47.
Papilio Carduelis, *Cramer*, *Pap.* I. t. 26, f. E.F. (1775).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Madras. From Captain J. M. Jones's Collection, 1856.

f. ♂. Darjeeling. From Pearson's Collection.

g. h. ♂ ♀. Bootan. From Pemberton's Collection.

The larva and pupa of *Pyrameis Cardui* are figured on Plate V., fig. 3, 3a, from Java, "where it feeds on a species of *Artemisia* the native name of which is *Godomollo*,—December."—(Horsfield, MS.)

Genus JUNONIA, *Hübner*.

JUNONIA, *Hübner*, *Verz. bek. Schmett.* p. 34 (1816). *E. Doubleday*, *Diurnal Lep.* p. 206.

VANESSA, *p. Fabricius*. *Godart*. &c.

281. *JUNONIA LEMONIAS*, *Linnaeus* Sp.

Papilio Lemonias, *Linnaeus*, *Syst. Nat.* II. p. 770 (1767).
Fabricius, *Ent. Syst.* III. I. p. 90.

Junonia Lemonias, *E. Doubleday*, *Diurnal Lep.* p. 208, n. 1.

Vanessa Lemonias, *Godart*, *Enc. Méth.* IX. p. 310. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 83.

Papilio Aonis, *Cramer*, *Pap. Exot.* I. t. 35, f. D.E.F. (1775).

a. ♂. N. India. From the Asiatic Society of Bengal.

b. ♂. Assam. From McClelland's Collection.

c. ♀. N. India. Presented by Colonel Buckley.

d. ♂. Pinang. Presented by Dr. Cantor.

282. *JUNONIA ERIGONE*, *Cramer* Sp.

Papilio Erigone, *Cramer*, *Pap. Exot.* I. t. 62, f. E.F. (1775).

Junonia Erigone, *E. Doubleday, Diurnal Lep.* p. 209, n. 3.
Temenis Erigone, *Hübner, Verz. bek. Schmett.* p. 34.

Vanessa Erigone, *Godart, Enc. Méth.* IX. p. 311. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 83.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

283. *JUNONIA LAOMEDIA*, *Linnæus Sp.*

Papilio Laomedia, *Linnæus, Syst. Nat.* II. p. 772 (1767).

Drury, Ins. I. t. 5, f. 3. *Cramer, Pap. Exot.* I. t. 8, f. F.G. *Fabricius, Ent. Syst.* III. I. p. 98.

Junonia Laomedia, *E. Doubleday, Diurnal Lep.* p. 209, n. 4.

Temenis Laomedia, *Hübner, Verz. bek. Schmett.* p. 34.

Vanessa Laomedia, *Godart, Enc. Méth.* IX. p. 322. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 83.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Dukhun. Presented by Colonel Sykes.

f. ♀. Calcutta. Presented by Colonel Buckley.

g. ♀. China. Presented by Dr. Cantor.

The larva and pupa of *Junonia Laomedia* are figured on Plate V., fig. 4, 4a, from Java, "where it feeds on a species of *Achyranthes*, bearing the native name of *Krema*,—January and February."—(Horsfield, MS.)

284. *JUNONIA CÆNONE*, *Linnæus Sp.*

Papilio Cænone, *Linnæus, Syst. Nat.* II. p. 770 (1767).

Cramer, Pap. Exot. I. t. 35, f. A.B.C. *Fabricius, Ent. Syst.* III. I. p. 90.

Junonia Cænone, *Hübner, Verz. bek. Schmett.* p. 34. *E. Doubleday, Diurnal Lep.* p. 209, n. 7.

Vanessa Cænone, *Godart, Enc. Méth.* IX. p. 318. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 78.

Var. *Papilio Hierta*, *Fabricius, Ent. Syst.* V. p. 424 (1798).

Vanessa Hierta, *Godart, Enc. Méth.* IX. p. 318.

a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.

d. e. ♂ ♀. Dukhun. Presented by Colonel Sykes.

285. JUNONIA ORITHYIA, Linnæus Sp.

Papilio Orithyia, *Linnæus, Syst. Nat.* II. p. 770 (1767).

Fabricius, Ent. Syst. III. I. p. 91. *Cramer, Pap. Exot.* I. t. 19, f. C.D. ♀; id. t. 32, f. E.F. ♂; id. III. t. 290, f. A.B. ♂; t. 281, f. E.F.; t. 290, f. C.D.

Junonia Orithyia, *Hübner, Verz. bek. Schmett.* p. 34. *E. Doubleday, Diurnal Lep.* p. 209, n. 8.

Vanessa Orithyia, *Godart, Enc. Méth.* IX. p. 317; id. *Enc. Méth.* IX. *Suppl.* p. 821. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 77.

? Junonia Ocyale, *Hübner, Verz. bek. Schmett.* p. 34 (1816). *E. Doubleday, Diurnal Lep.* p. 209, n. 9.

a. b. c. d. e. f. g. h. Java. From Dr. Horsfield's Collection.

i. ♀. Sumatra. From Raffles's Collection.

j. k. ♂ ♀. Pinang. Presented by Dr. Cantor.

l. m. ♂ ♀. N. India. Presented by Colonel Buckley.

n. ♀. Bootan. From Pemberton's Collection.

o. ♂. Canara. Presented by S. N. Ward, Esq.

The larva and pupa of *Junonia Orithyia* are figured on Plate V., fig. 5, 5a, from Java, "and feeds on a species of *Vitex*, bearing the native name of *Laban*,—December."—(Horsfield, MS.)

286. JUNONIA VELLIDA, Fabricius Sp.

Papilio Vellida, *Fabricius, Ent. Syst.* III. I. p. 91 (1793).

Donovan, Ins. of New Holland, t. .

Junonia Vellida, *E. Doubleday, Diurnal Lep.* p. 209, n. 14.

Vanessa Vellida, *Godart, Enc. Méth.* IX. *Suppl.* p. 807. *Boisduval, Voy. de l'Astrolabe*, p. 120. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 77.

Vanessa Calybe, *Godart, Enc. Méth.* IX. p. 316 (1819).

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♂. Sumatra. From Raffles's Collection.

Remark.—These specimens of *Vanessa Vellida* have been compared at the British Museum with examples from Australia, with which they are perfectly identical.

287. *JUNONIA ASTERIE*, Linnæus Sp.

Papilio Asterie, Linnæus, *Syst. Nat.* II. p. 769 (1767).

Cramer, Pap. Exot. I. t. 58, f. D.E. *Fabricius, Ent. Syst.* III. I. p. 89.

Junonia Asterie, E. *Doubleday, Diurnal Lep.* p. 209, n. 15.

Aleyoneis Asterie, Hübner, *Verz. bek. Schmett.* p. 35.

Vanessa Asterie, Godart, *Enc. Méth.* IX. p. 321. E. *Doubleday, List Lep. Brit. Mus. pt. I.* p. 76.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. e. ♂ ♀. N. India. Presented by Colonel Buckley.

The larva and pupa of *Junonia Asterie* are figured on Plate V., fig. 6, 6a, from Java, "and feeds on a species of *Justicia*, bearing the native name of *Keji-Wangu*,—March."—(Horsfield, MS.)

288. *JUNONIA ALMANA*, Linnæus Sp.

Papilio Almana, Linnæus, *Syst. Nat.* II. p. 769 (1767).

Cramer, Pap. Exot. I. t. 58, f. F.G. *Fabricius, Ent. Syst.* III. I. p. 89.

Junonia Almana, E. *Doubleday, Diurnal Lep.* p. 209, n. 16.

Aleyoneis Almana, Hübner, *Verz. bek. Schmett.* p. 35.

Vanessa Almana, Godart, *Enc. Méth.* IX. p. 313. E. *Doubleday, List Lep. Brit. Mus. pt. I.* p. 77.

a. b. ♂. N. India. From the Asiatic Society of Bengal.

c. ♀. Assam. From Dr. McClelland's Collection.

d. e. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus PRECIS, Hübner.

PRECIS, Hübner, *Verz. bek. Schmett.* (1816).

JUNONIA, Sect. II. (*Precis*), E. *Doubleday, Diurnal Lep.* p. 209 (1849).

SALAMIS, p. Boisduval.

289. *PRECIS IDA*, Cramer Sp.

Papilio Ida, Cramer, *Pap. Exot.* I. t. 42, f. C.D. (1775);

id. IV. t. 374, f. C.D.

Vanessa Idamene, Godart, Enc. Méth. IX. p. 315 (1819).

Junonia (Precis) Iphita, ♀, apud E. Doubleday, Diurnal Lep. p. 210, n. 32.

? *Papilio Zelima, Donovan, Ins. of India (1800-3).*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Pinang. Presented by Dr. Cantor.

290. *PRECIS IPHITA, Cramer Sp.*

Papilio Iphita, Cramer, Pap. Exot. III. t. 209, f. C.D. (1780). Fabricius, Ent. Syst. III. I. p. 109.

Vanessa Iphita, Godart, Enc. Méth. IX. p. 314.

Salamis Iphita, Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 85.

Junonia (Precis) Iphita, E. Doubleday, Diurnal Lep. p. 210, n. 32.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. Darjeeling. From Pearson's Collection.

g. h. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

291. *PRECIS HARA, Moore (Plate IIIa., fig. 1).*

Precis Hara, n. sp.—UPPER-SIDE golden-brown, darker on outer margins : *fore* and *hind-wings* each having three narrow transverse black bands on the anterior half of the wings, the outer band on hind-wing being zig-zagged, with a parallel row of small black spots between the outer and middle band ; within each discoidal cell are two narrow and rather square-shaped spaces, those on the hind-wing being the narrowest. *UNDER-SIDE* dull brown, marked as above, with deep brown, but the two inner bands broad, the outer very narrow ; on the *hind-wing*, above the marks, within discoidal cell, is a narrow brown mark ; the outer margins are also deep brown.

a. ♂. Silhet. Presented by the Trustees of the British Museum.

b. c. d. ♂. N. India. Presented by Lieut. Hugo James.

c. ♀. N. India. Presented by Colonel Buckley.

292. *PRECIS POLYNICE*, Cramer Sp.

Papilio Polynice, *Cramer, Pap. Exot.* II. t. 195, f. D.E.
(1780). *Fabricius, Ent. Syst.* III. I. p. 89.

Junonia (Precis) Polynice, *E. Doubleday, Diurnal Lep.*
p. 210, n. 35.

Vanessa Polynice, *E. Doubleday, List Lep. Brit. Mus.*
pt. I. p. 80.

Vanessa Polynissa, *Godart, Enc. Méth.* IX. p. 308 (1819).

a. ♀. Java. From Dr. Horsfield's Collection.

Genus *ERGOLIS*, Boisduval.

ERGOLIS, *Boisduval, Spéc. Gén. Lép.* I. p. 2, t. 4, f. 4 (1836).
Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 409
(1851).

ARIADNE, *Horsfield, Cat. Lep. Mus. E.I.C.* t. 6, f. 2 (1829).

ELYMNIAS, *p. Hübner.*

BIBLIS, *p. Godart.*

293. *ERGOLIS ARIADNE*, Linnæus Sp.

Papilio Ariadne, *Linnæus, Syst. Nat.* II. p. 778 (1767).
Fabricius, Syst. Ent. p. 507.

Ergolis Ariadne, *Boisduval. E. Doubleday, List Lep.*
Brit. Mus. pt. I. p. 145. *Westwood, in Doubleday*
and Hewitson's Diurnal Lep. p. 410, n. 1.

Elymnias Ariadne, *Hübner, Verz. bek. Schmett.* p. 328.

Biblis Ariadne, *Godart, Enc. Méth.* IX. p. 327.

Papilio Merione, *Cramer, Pap. Exot.* II. t. 144, f. G.H.
(1779).

a. b. ♂ ♀. N. India. Presented by Colonel Sykes.

c. d. ♂ ♀. Darjeeling. From Pearson's Collection.

294. *ERGOLIS CORYTA*, Cramer Sp.

Papilio Coryta, *Cramer, Pap. Exot.* I. t. 86, f. E.F.
(1779).

Ergolis Coryta, *Boisduval, Spéc. Gén. Lép.* I. p. 2, t. 4,
f. 4. *E. Doubleday, List Lep. Brit. Mus. pt. I.*
p. 145. *Westwood, in Doubleday and Hewitson's*
Diurnal Lep. p. 410, n. 2.

Ariadne Coryta, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 6, f. 2.

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
- e. f. ♂ ♀. N. India. From the Asiatic Society of Bengal.
- g. ♂. Assam. From McClelland's Collection.
- h. ♂. Madras.

The larva and pupa of *Ergolis Coryta* are figured on Plate VI., fig. 6, 6a, from Java, and "feeds on the *Jarak* (*Ricinus communis*)—December. Not very common."—(Horsfield, MS.)

The transformations of a species of this genus were also discovered in Ceylon by E. L. Layard, Esq., and are figured among his original drawings.

Genus CYNTHIA, *Fabricius*.

CYNTHIA, *Fabricius*, *Syst. Gloss.* (*Illiger's Mag.* VI. p. 281, 1808).

ANARTIA, *p.* *Hübner*.

PAPHIA, *p.* *Horsfield*.

295. *CYNTHIA ARSINOË*, *Cramer* Sp.

♂ *Papilio Arsinoë*, *Cramer*, *Pap. Exot.* II. t. 160, f. A.B. (1777). *Fabricius*, *Ent. Syst.* III. I. p. 74.

Cynthia Arsinoë, *Fabricius*, *Syst. Gloss.* (*Illiger's Mag.* VI. p. 281, 1808). *E. Doubleday*, *Diurnal Lep.* p. 213, t. 24, f. 5.

Vanessa Arsinoë, *Godart*, *Enc. Méth.* IX. p. 297. *Boisduval*, *Voy. de l'Astrolabe*, *Ins.* p. 123.

Anartia Arsinoë, *Hübner*, *Exot. Samml.* *E. Doubleday*, *List Lep. Brit. Mus. pt.* I. p. 84.

♀ *Papilio Juliana*, *Cramer*, *Pap. Exot.* III. t. 280, f. A.B. (1782). *Fabricius*, *Ent. Syst.* III. I. p. 108.

♀ *Cynthia Deione*, *Erichson*, *Nova Acta Acad. Nat. Cur.* XVI. II. t. 40, f. 2, 2a (1833).

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♂. Darjeeling. From Pearson's Collection.

d. e. ♂ ♀. Darjeeling. Presented by Colonel Buckley.

Genus AMNOSIA, Boisduval.

AMNOSIA, Boisduval. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 259 (1850).

296. AMNOSIA DECORA, Boisduval.

Amnobia decora, Boisduval, MS. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 88. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 260, t. 51, f. 4, ♂.

- a. ♂. Java. From Mr. Argent's Collection.
- b. ♀. Java. From T. Tatum, Esq.

Genus CYRESTIS, Boisduval.

CYRESTIS, Boisduval, *Voy. de l'Astrolabe*, p. 117 (1832).

297. CYRESTIS LUTEA, Zinken-Sommer Sp.

Amathusia lutea, ♂, et A. nivea, ♀, Zinken-Sommer, in *Nova Acta Acad. Nat. Cur.* XVI. p. . t. 14, f. 1, 2 (1831).

Marpesia Hylas, Hübner, *Samml. Exot. Schmett. Band III.* f. , ♂ ♀, nec Clerck.

Nymphalis Hylas, Godart, *Enc. Méth. IX.* p. 361.

Cyrestis Hylas, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 88.

Cyrestis Recaranus, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 261, n. 3 (1850).

- a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.
- c. d. e. Java. From T. Tatum, Esq.

298. CYRESTIS THYODAMAS, Boisduval.

Cyrestis Thyodamas, Boisduval, in *Cuv. Rég. An. edit. Crochard, Ins.* t. 138, f. 4 (). E. Doubleday, List Lep. Brit. Mus. pt. I. p. 88. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 261, n. 4, t. 32, f. 3.

Amathusia Ganescha, Kollar, in Hügel's *Kaschmir*, IV. pt. II. p. 430, t. 7, f. 3, 4 (1844).

- a. b. ♂ ♀. Darjeeling. From Pearson's Collection.
- c. d. ♂. N. India. Presented by Colonel Buckley.

299. CYRESTIS PERIANDER, *Fabricius Sp.*

Papilio Periander, *Fabricius*, *Ent. Syst.* III. I. p. 67 (1793). *Donovan, Ins. of India*, t. 37, f. 1.

Cyrestis Periander, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 261, n. 8.

Nymphalis Periander, *Godart, Enc. Méth.* IX. p. 362.

Paphia Periander, *Horsfield, Descr. Cat. Lep. Mus. E.I.C.* t. 5, f. 3, 3a.

a. b. Java. From Dr. Horsfield's Collection.

300. CYRESTIS RISA, *E. Doubleday*.

Cyrestis Risa, *E. Doubleday, MS. Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 262, n. 10, t. 32, f. 4.

a. b. Assam. From Griffith's Collection.

c. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

301. CYRESTIS RAHRIA, *Westwood* (*Plate IIIa, fig. 2*).

Cyrestis Rahria, *Westwood, MS. in Doubleday and Hewitson's Diurnal Lep.* p. 262, n. 11 (1850).

? Cyrestis Neduna, *De Haan, MS.*

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

Genus PARTHENOS, *Hübner*.

PARTHENOS, *Hübner, Verz. bek. Schmett.* p. 38 (1816).

MINETRA, *Boisduval, Voy. de l'Astrolabe*, p. 126 (1832). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 265.

302. PARTHENOS SYLVIA, *Cramer Sp.*

Papilio Sylvia, *Cramer, Pap. Exot.* I. t. 43, f. F.G. (1779).

Parthenos Sylvia, *Hübner, Verz. bek. Schmett.* p. 38.

Nymphalis Sylvia, *Godart, Enc. Méth.* IX. p. 381.

Phyllophasis Sylvia, *Blanchard, in Lap. Hist. Nat. An.* Art. III. p. 448.

Minetra Sylvia, *Boisduval, Voy. de l'Astrolabe*, p. 127.

*E. Doubleday, List Lep. Brit. Mus. pt. I. p. 86.
Westwood, in Doubleday and Hewitson's Diurnal
Lep. p. 266.*

- a. b. c. d. ♂ ♀.* Java. From Dr. Horsfield's Collection.
- e.* Pinang. Presented by Dr. Cantor.

303. *PARTHENOS GAMBRISIUS*, Fabricius Sp.

*Papilio Gambrisius, Fabricius, Ent. Syst. III. I. p. 85
(1793). Donovan, Ins. of China, t. 38.*

*Minetra Gambrisius, E. Doubleday, List Lep. Brit. Mus.
pt. I. p. 86. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 266, t. 51, f. 2.*

- a.* N. India. Presented by Colonel Sykes.
- b.* Silhet. Presented by Colonel Buckley.
- c.* Pinang. Presented by Dr. Cantor.

Captain Mortimer J. Slater, in his MS. Notes, p. 244, remarks:—
“I met with this insect (*Pap. Gambrisius*) at Dacca, in the deepest jungle. Its flight is rapid and dodgy, and in consequence is very difficult to capture on the wing.”

Genus PROTHOË, Hübner.

*PROTHOË, Hübner, Samml. Exot. Schmett. (1806–27). Westwood, in
Doubleday and Hewitson's Diurnal Lep. p. 267.*

AUTONEMA, Boisduval, MSS.

304. *PROTHOË FRANCKI*, Godart Sp.

*Nymphalis Franck, Godart, Enc. Méth. IX. p. 825
(1819).*

*Prothoë Franckii, Hübner, Samml. Exot. Schmett. Band
II. t. . E. Doubleday, List Lep. Brit. Mus.
pt. I. p. 88. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 287, t. 51, f. 3.*

*Paphia Franck, Horsfield, Cat. Lep. Mus. E.I.C. t. 5,
f. 4, 4a.*

Autonema Franckii, Boisduval, MSS.

- a. b. c. ♂.* Java. From Dr. Horsfield's Collection.

Genus TERINOS, *Boisduval*.

TERINOS, *Boisduval*, *Spéc. Gén. Lép.* I. t. 9, f. 4 (1836). *E. Doubleday, Diurnal Lep.* p. 159.

305. TERINOS CLARISSA, *Boisduval*.

Terinos Clarissa, *Boisduval*, *Spéc. Gén. Lép.* I. t. 9, f. 4 (1836). *E. Doubleday, Diurnal Lep.* p. 160, t. 21, f. 3.

a. ♀. Java. From Mr. Argent's Collection.

b. ♂. Borneo. Presented by Edwin Shepherd, Esq., 1857.

Genus CIRROCHROA, *E. Doubleday*.

CIRROCHROA, *E. Doubleday, Diurnal Lep.* p. 157 (1848).

306. CIRROCHROA THAIS, *Fabricius Sp.*

Papilio Thais, *Fabricius, Ent. Syst.* III. I. p. 149 (1793).

Cirrochroa Thais, *E. Doubleday, Diurnal Lep.* p. 158, n. 2.

Argynnis Thais, *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 66.

Argynnis Thea, *Godart, Enc. Méth.* IX. p. 260 (1819).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

307. CIRROCHROA AORIS, *E. Doubleday*.

♂ Cirrochroa Aoris, *E. Doubleday, MS. Westwood and Hewitson's Diurnal Lep.* p. 158, t. 21, f. 1 (1847).

a. b. c. ♂. Darjeeling. From Pearson's Collection.

e. f. g. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

308. CIRROCHROA CLAGIA, *Godart Sp.*

Argynnis Clagia, *Godart, Enc. Méth.* IX. *Suppl.* p. 816 (1823). *Boisduval, Spéc. Gén. Lép.* I. t. 10, f. 6. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 66.

Cirrochroa Clagia, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 158, n. 3.

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

309. *CIRROCHROA BAJADETA*, Moore (Plate IIIa, fig. 3).

Cirrochroa Bajadeta, n. sp.—Male: UPPER-SIDE ferruginous, with a broadish blackish border from the anterior angle along the exterior margin of both wings, and having narrow blackish lunular marks disposed along the inner margin of the border, also three black spots on each posterior wing from the anal angle. UNDER-SIDE paler, with a tapering fascia crossing both wings, this being white on the posterior wing; also two ranges of lunular markings along the exterior margin, and six spots disposed across the posterior wing, near the white fascia.

The female differs above, in being of a brighter colour, and in having the border blacker, the anterior wings having a narrow blackish disco-cellular mark; the lunular marks of the posterior wings being darker and more suffused, and having six spots disposed from anal to anterior angle. Beneath, the colours and markings are stronger. *var. Bornco.*

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

310. *CIRROCHROA RAVANA*, Moore.

Cirrochroa Ravana, n. sp.—Female: UPPER-SIDE, with base of both wings, to irregular transverse black line from middle of costal margin to a little above the anal angle, dull ferruginous, the rest being paler and brighter, and almost yellow on the fore-wing; exterior margin of *fore-wing* broadly from the apex to posterior angle, and narrowly along exterior margin of *hind-wing* black, the former with an indistinct pale spot on the apex, the latter with marginal and sub-marginal lunulated line and some spots across the disc black. UNDER-SIDE much paler, faintly glossed with pearly-white; a tapering pearly-white band from middle of costal margin to near anal angle, a marginal and a sub-marginal lunulated line, and a row of black spots encircled with ferruginous across disc of hind-wing; markings within discoidal cell ferruginous; two small white spots at the apex of fore-wing.

a. ♀. Bornco. Presented by Edwin Shepherd, Esq.,
1857.

Genus MESSARAS, E. Doubleday.

MESSARAS, E. Doubleday, MS. Westwood and Hewitson's Diurnal Lep. p. 163 (1848).

311. *MESSARAS ERYMANTHIS*, Drury Sp.

Papilio Erymanthis, Drury, Ins. I. t. 15, f. 3, 4 (1770).

- Cramer, Pap. Exot.* III. t. 238, f. F.G. *Fabricius,*
Ent. Syst. III. I. p. 139.
Messaras Erymanthis, E. Doubleday, MS. Westwood and
Hewitson's Diurnal Lep. p. 164.
Argynnus Erymanthis, Godart, Enc. Méth. IX. p. 257.
Boisduval, Voy. de l'Astrolabe, Ins. p. 115. *E.*
Doubleday, List Lep. Brit. Mus. pt. I. p. 67.
a. b. c. d. Java. From Dr. Horsfield's Collection.
e. ♀. N. India. Presented by Colonel Sykes.
f. ♂. Canara. Presented by S. N. Ward, Esq.
g. ♀. Pinang. Presented by Dr. Cantor.
h. Borneo. Presented by Edwin Shepherd, Esq., 1857.

"Tolerably plentiful at Dacca, 1844. Flies quietly, and is easily captured. I used to find one or two almost invariably near a deep pool of stagnant water overhung with bamboo jungle, where *Athyma Leucothoë* was always to be found, together with two of the species resembling it in flight and general colour."—(Captain Mortimer J. Slater's MS. Notes, p. 148.)

312. MESSARAS ALCIPPE, Cramer Sp.

Papilio Alcippe, Cramer, Pap. Exot. IV. t. 389, f. G.H.
 (1782).

Messaras Alcippe, E. Doubleday, MS. Westwood and
Hewitson's Diurnal Lep. p. 165.

Argynnus Alciope, Godart, Enc. Méth. IX. p. 259. *E.*
Doubleday, List Lep. Brit. Mus. pt. I. p. 67.

- a. ♀.* Silhet. Presented by the Trustees of the
 British Museum.
b. ♂. ? Borneo. Presented by Edwin Shepherd, Esq.,
 1857.

Genus ATELLA, E. Doubleday.

ATELLA, E. Doubleday, Diurnal Lep. p. 165 (1848).

PHALANTA, Horsfield, Desc. Cat. Lep. Mus. E.I.C. t. 7, f. 5
 (1829).

313. ATELLA PHALANTA, Drury Sp.

Papilio Phalanta, Drury, Ins. I. t. 21, f. 1, 2 (1770).
Fabricius, Ent. Syst. III. I. p. 149.

- Atella Phalanta*, *E. Doubleday, MS.* *Westwood and Hewitson's Diurnal Lep.* p. 167.
Argynnis Phalanta, *Godart, Enc. Méth.* IX. p. 259. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 67.
Papilio Columbina, *Cramer, Pap. Exot.* III. t. 238, f. A.B. (1780); t. 337, f. D.E. var.
Argyronome Columbina, *Hübner, Verz. bek. Schmett.* p. 32.
Phalanta Columbina, *Horsfield, Desc. Cat. Lep. Mus. E.I.C.* t. 7, f. 5.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. f. g. ♂ ♀. N. India. From the Asiatic Society of Bengal.
h. ♀. N. India. Presented by Colonel Buckley.

The larva and pupa of *Atella Phalanta* are figured on Plate V., fig. 7, 7a, from Java, and "feeds on a species of *Ixora*,—November. Seearce."—(Horsfield, MS.)

314. *ATELLA EGISTA*, Cramer Sp.

- Papilio Egista*, *Cramer, Pap. Exot.* III. t. 281, f. C.D. (1780).
Atella Egista, *E. Doubleday, MS.* *Westwood and Hewitson's Diurnal Lep.* p. 167, n. 6.
Issoria Egista, *Hübner, Verz. bek. Schmett.* p. 31.
Argynnis Egista, *Godart, Enc. Méth.* IX. p. 261. *Boisduval, Voy. de l'Astrolabe, Ins.* p. 113. *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 66.
Terinos Sinha, *Kollar, in Hügel's Kaschmir*, IV. pt. II. p. 438 (1844).
a. b. ♂ ♀. Java. Presented by Dr. Horsfield.
c. d. e. ♂ ♀. N. India. Presented by Colonel Buckley.

Genus LAOGONA, Boisduval.

- LAOGONA*, *Boisduval, Spéc. Gén. Lép.* I. t. 10, f. 3 (1836).
SYMBRENTHIA, *Hübner, Verz. bek. Schmett.* (1816).
HYPANARTIA, *Hübner-Geyer, Samml. Exot. Schmett.* (1841).

315. *LAOGONA HYPOPOCLA*, Cramer Sp.

Papilio Hypopoca, Cramer, Pap. Exot. III. t. 220, f. C.D. (1780).

Laogona Hypopoca, Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 83. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 191, n. 1.

Hypanartia Hypopoca, Hübner and Geyer, Samml. Exot. Schmett. III. (1841).

Vanessa Hypopoca, Godart, Enc. Méth. IX. p. 298.

♀ *Papilio Lucina*, Cramer, Pap. Exot. IV. t. 330, f. E.F. (1782).

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. Darjeeling. From Pearson's Collection.

h. Bootan. From Pemberton's Collection.

i. j. N. India. Presented by Colonel Buckley.

316. *LAOGONA HYPSELIS*, Godart Sp.

Vanessa Hypselis, Godart, Enc. Méth. IX. Suppl. p. 818 (1823).

Laogona Hypselis, Boisduval, Spéc. Gén. Lép. I. t. 10, f. 3. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 83. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 191, n. 2, t. 25, f. 1.

a. b. c. ♂. N. India. Presented by Colonel Buckley.

d. Var. From T. Tatum, Esq.

Genus *CETHOSIA*, Fabricius.

CETHOSIA, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 280, 1807.)
Godart. Latreille. Doubleday.

ALAZONIA, Hübner, Verz. bek. Schmett. p. 46 (1816).

317. *CETHOSIA BIBLIS*, Drury Sp.

Papilio Biblis, Drury, Ins. I. t. 4, f. 2 (1780). Cramer, Pap. Exot. II. t. 175, f. A.B.

Cethosia Biblis, Westwood, in Drury's Ins. 2nd edit. I. p. 9. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 65; Diurnal Lep. p. 151, n. 1.

- Alazonia Symbiblis (part), Hübner, Verz. bek. Schmett.*
p. 46 (1816).
- Cethosia Biblina, Godart, Enc. Méth. IX. p. 248.*
- Papilio Penthesilea, apud Fabricius, Spec. Ins. II. p. 88.*
- a. b. ♂. N. India. Presented by Lieut. Hugo James.*
- c. d. ♂ ♀. Darjeeling. From Pearson's Collection.*
- e. f. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.*

318. *CETHOSIA PENTHESILEA*, Cramer Sp.

Papilio Penthesilea, Cramer, Pap. Exot. II. t. 145,
f. B.C. (1776).

Cethosia Penthesilea, Godart, Enc. Méth. IX. p. 248. Lucas, Lep. Exot. t. 55, f. 1. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 65; Diurnal Lep. p. 151, n. 2.

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

319. *CETHOSIA ÆOLE*, De Haan MS.

Cethosia Æole, De Haan.—UPPER-SIDE: male, *fore-wing* having the posterior basal third to near posterior angle rufous, the rest black, also some transverse markings in discoidal cell: the basal half of the wing with a bluish-purple tinge, being most vivid at the unison of the two colours; also a white transverse patch near middle of wing: *hind-wing* with a broad black exterior border, the rest rufous, with bluish-purple tinge; three small black spots disposed linearly from near anal angle; also an indistinct spot between discoidal and median veinlets; wings deeply scalloped, and with white zigzag marginal markings. Female, with the upper-side of the *fore-wing* nearly all black, excepting a narrow yellow space along the middle of posterior margin: *hind-wing* with a broader marginal border than in male, the abdominal margin yellowish, the rest rufous; discal spot larger, and tooth-shaped; white patch also larger. Underside of both sexes nearly as in *Ceth. Penthesilea*, but the white central irregular fascia of the *hind-wing* is very broad in *Ceth. Æole*.

- a. b. ♂ ♀. Sumatra. From Dr. Horsfield's Collection.*
- c. ♂. Borneo. Presented by Edwin Shepherd, Esq., 1857.*

320. *CETHOSIA CYANE*, Fabricius Sp.

Papilio Cyane, *Fabricius*, *Syst. Ent.* p. 503 (1775); *Ent. Syst.* III. I. p. 115. Drury, *Ins.* I. t. 4, f. 1. Cramer, *Pap. Exot.* IV. t. 295, f. C.D.

Cethosia Cyane, Godart, *Enc. Méth.* IX. p. 247. Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 443. E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 65; *Diurnal Lep.* p. 152, n. 6.

Alazonia Symbiblis (*part*), Hübner, *Verz. bek. Schmett.* p. 46.

a. b. ♂ ♀. Canara. Presented by S. N. Ward, Esq.
d. e. f. g. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

The larva and pupa of *Cethosia Cyane* are figured on Plate V. fig. 8, 8a, from Canara, Southern India, and were copied from the original drawings made by S. N. Ward, Esq.

Captain Mortimer J. Slater also observed the transformations of "a species of *Cethosia*, from a larva feeding on the Passion-flower. The larva was yellow, black, and crimson in alternate bars, and covered with stinging spines. Pupa hung by tail and band, and in leaf-like appendages resembled that of *Papilio Pammon*."—(MS. Notes, p. 302.)

Genus ARGYNNIS, *Fabricius*.

ARGYNNIS, *Fabricius*. (*Illiger's Mag.* VI. 1808.) Ochsenheimer, *Schmett. von Eur.* IV. p. 16 (1816). Doubleday.

ARGYNNIS, BRENTHIS, ISSORIA, ACIDALIA, ARGYRONOME, Hübner, *Verz. bek. Schmett.* (1816).

321. *ARGYNNIS NIPHE*, *Linnæus* Sp.

♀ Papilio Niphe, *Linnæus*, *Syst. Nat.* II. p. 785 (1767). Cramer, *Pap. Exot.* I. t. 14, f. B.C.D.E. *Fabricius*, *Ent. Syst.* III. I. p. 142.

Argynnis Niphe, Godart, *Enc. Méth.* IX. pp. 261, 806. Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 440, t. 13, f. 12. E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 67; *Diurnal Lep.* p. 174, n. 1.

Acidalia Niphe, Hübner, *Verz. bek. Schmett.* p. 31.

♀ Papilio Hyperbius, *Linnæus*, *Amoen. Acad.* VI. p. 408.

- δ Papilio Argyrius, Sparmann, *Amœn. Acad.* VII. p. 502 (1768).
- δ Papilio Argynnis, Drury, *Ins. I. t. 6, f. 2* (1770).
- Argynnis Tephnia, Godart, *Enc. Méth.* IX. p. 262 (1819).
- a. b. c. d. δ ♀. Java. From Dr. Horsfield's Collection.
- e. ♀. Pinang. Presented by Dr. Cantor.
- f. ♀. Darjeeling. From Pearson's Collection.
- g. h. i. δ ♀. N. India. Presented by Colonel Buckley.

322. *ARGYNNIS ARUNA*, Moore (Plate IIIa, fig. 4).

A. Aruna, n. sp.—UPPER-SIDE, wings fulvous; anterior wings with a broad black patch in discoidal cell and between the nervules, and duller markings on the exterior margin: posterior wings having the whole space between the nervules posteriorly occupied with black, and leaving only small lunular markings on the exterior margin, the latter becoming bluish at the anal angle. UNDER-SIDE with the anterior wings broadly suffused with black; posterior wings with the basal half silvery, and with black markings across the wing from outer to anal angle. Shape of wings as in *Arg. Niphe*; expanse $3\frac{1}{8}$ in.

a. δ . N. India. Presented by Colonel Buckley.

323. *ARGYNNIS ISSÆA*, G. R. Gray.

Argynnis Issæa, G. R. Gray, *Lep. Ins. of Nepal*, p. 11 (1846). E. Doubleday, *Diurnal Lep.* p. 174, n. 3.
Argynnis Lathonia, apud Kollar, in Hügel's *Kaschmir*, IV. pt. II. p. 440.

a. b. δ ♀. Bootan. From Pemberton's Collection.

c. d. δ . N. India. Presented by Colonel Hearsey.

324. *ARGYNNIS KAMALA*, Moore.

Argynnus Kamala, n. sp.—UPPER-SIDE golden-yellow; fore-wings somewhat dusky at the base; hind-wings tinged in parts with yellowish-green: fore-wing with all the veins and veinlets deeply defined with black; the black markings disposed exactly as in *Argyn. Paphia*, but larger and duller, and in *A. Kamala* the submarginal row of spots are less defined and continuous, rather forming a stripe than a linear series of spots; the marginal row is connected between the

veinlets by a narrow line: *hind-wing* with an irregular zigzag inner black band, then a row of five round black spots, and a submarginal row of seven linear-shaped spots (the anterior and third being smallest), also a marginal black line; the angles and end of veinlets also black. *UNDER-SIDE*, *fore-wing* yellow, the markings within discoidal cell and about the disc as above, but submarginal row paler; none on margin; near the apex a green patch, containing three silvery-white spots: *hind-wing* beautiful metallic green, with a broad submarginal and inner band, and a series of marks about the basal half silvery-white, all margined more or less on one side with black; between the submarginal and inner band some small spaces of yellow, and an indistinct row of reddish spots. Wings shaped as in *Argynnis Paphia*; expanse $2\frac{7}{8}$ in.

a. b. c. ♂. N. India. Presented by Colonel Buckley.

325. ARGYNNIS RUDRA, Moore.

Argynnus Rudra, n. sp.—*UPPER-SIDE* bright fulvous, duller and slightly tinged with green at the base: *fore-wing* with the markings disposed as in *Arg. Childreni*, but somewhat smaller, but with a marginal row of small spots: *hind-wing* with the spots also disposed as in *Arg. Childreni*, but with a marginal row of spots. *UNDER-SIDE*, *fore-wing* fulvous-yellow, greenish at the apex; the spot on the exterior half indistinct: *hind-wing* light green, palest on abdominal margin; a ferruginous irregular band crossing the basal half; an irregular more or less white line crossing from anterior margin near the angle to abdominal angle; also some white disposed from anal angle and inclosing the green colour, and thus forming a submarginal row; along the outside of the white line, some ferruginous spots, centred with white. Wings shaped as in *Arg. Laodice*, which appears to be a near ally; expanse of wings 3 in.

a. ♂. N. India. Donor unknown.

326. ARGYNNIS CHILDRENI, G. R. Gray.

Argynnus Childreni, G. R. Gray, *Gray's Zool. Misc.* p. 33 (1832); *Lep. Ins. of Nepal*, p. 11, t. 11. E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 68; *Diurnal Lep.* p. 174, n. 6.

Argynnus Sakontala, Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 439, t. XIII. (1844).

a. b. ♀. N. India. Presented by Colonel Buckley.

Genus DIADEMA, Boisduval.

DIADEMA, Boisduval, *Voy. de l'Astrolabe, Ins. p. 135* (1832). Westwood, in Doubleday and Hewitson's *Diurnal Lep. p. 279*.

HYPOLIMNAS, ESOPTRIA, et PANOPEA, p. Hübner, Verz. bek. Schmett. (1816).

APATURA (*part*), Fabricius. Horsfield.

NYMPHALIS (*part*), Godart.

327. DIADEMA BOLINA, Linnæus Sp.

♂ Papilio Bolina, Linnæus, *Syst. Nat. II. p. 781* (1767).

Clerck, *Icon. t. 21. Drury, Ins. I. t. 14, f. 1, 2.*

Cramer, *Pap. Exot. I. t. 65, f. E.F. Fabricius, Ent. Syst. III. pt. I. p. 126. Herbst, Pap. t. 244, f. 3, 4.*

Diadema Bolina, Boisduval, *Faune Ent. de Madag. p. 39.*

E. Doubleday, *List Lep. Brit. Mus. pt. I. p. 98.*

Westwood, in Doubleday and Hewitson's *Diurnal Lep. p. 280, n. 1.*

Nymphalis Bolina, Godart, *Enc. Méth. IX. p. 396.*

Apatura Bolina, Zinken-Sommer, *Nova Acta Acad. Nat. Cur. XV. p. 194.*

♀ Papilio Misippus, Linnæus, *Syst. Nat. II. p. 767.*

Danaïs Misippus, Godart, *Enc. Méth. IX. p. 394.*

♀ Papilio Dioceippus, Cramer, *Pap. Exot. I. t. 28, f. B.C. (1779). Herbst, Pap. t. 155, f. 3, 4. Fabricius, Ent. Syst. III. pt. I. p. 51.*

Var. ♀ Papilio Inaria, Cramer, *Pap. Exot. III. t. 214, f. A.B. Herbst, Pap. t. 157, f. 5, 6.*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. g. h. ♂ ♀. N. India. Presented by Colonel Buckley.

i. (Hermaphrodite). Canara. Presented by S. N. Ward, Esq.

j. ♂. Borneo. Presented by H. Low, Esq.

k. ♀ var. (*P. Inaria*, Cram.) Madras. From Captain J. M. Jones's Collection.

328. DIADEMA AUGE, Cramer Sp.

♂ Papilio Auge, Cramer, *Pap. Exot. II. t. 190, f. A.B. (1779).*

Diadema Auge, *E. Doubleday, List Lep. Brit. Mus. pt. I.*
p. 98.

♂ Papilio Lasinassa, *Cramer, Pap. Exot. III. t. 205, f. A.B.*
(1782). *Fabricius, Ent. Syst. III. pt. I. p. 127.*
Lucas, Hist. Nat. Lep. Exot. t. 70, f. 2.

Nymphalis Lasinassa, *Godart, Enc. Méth. IX. p. 395.*

Apatura Lasinassa, *Horsfield, Cat. Lep. Mus. E.I.C. t. 8.*

Var. ♂ Papilio Iphigenia, *Cramer, Pap. Exot. I. t. 67, f. D.E.*
(1779).

♂ Papilio Alcithoë, *Cramer, Pap. Exot. I. t. 80, f. A.B.*
(1779). *Hübner, Samml. Exot. Schmett. Bd. I.*
t. —.

Var. ♀ Papilio Perimele, *Cramer, Pap. I. t. 65, f. C.D.; id. t. 67,*
f. B. (1779).

Var. ♀ Papilio Melita, *Cramer, Pap. I. t. 28, f. D.E. (1779).*

Var. ♀ Papilio Alcmena, *Cramer, Pap. I. t. 67, f. A. (1779).*

Var. ♀ Papilio Antigone, *Cramer, Pap. I. t. 67, f. C. (1779).*

Var. ♀ Papilio Proserpina, *Cramer, Pap. III. t. 218, f. C.D.*
(1782).

Var. ♀ Papilio Manilia, *Cramer, Pap. Exot. t. 255, f. A.B.*
(1782).

Var. ♀ Papilio Porphyria, *Cramer, Pap. t. 255, f. E.F.*

Var. ♀ Papilio Velleda, *Cramer, Pap. t. 349, f. C.D. (1782).*

Var. ♀ Papilio Eriphile, *Cramer, Pap. t. 376, f. A.B. (1782).*

♂ Papilio Liria, *Fabricius, Ent. Syst. III. pt. I. p. 126.*

Nymphalis Liria, *Godart, Enc. Méth. IX. p. 395.*

♀ Papilio Nerina, *Fabricius, Ent. Syst. III. pt. I. p. 133.*
Donovan, Ins. of New Holl. t. 27, f. 1.

Var. ♀ Papilio Jacintha, *Fabricius, Ent. Syst. III. pt. I. p. 60.*
Jones, Icon. IV. t. 51. Donovan, Ins. of China,
t. 37, f. 1. Drury, Ill. App. vol. II. t. 21, f. 1, 2.

♀ Papilio Avia, *Fabricius, Ent. Syst. III. pt. I. p. 111.*

a. b. c. d. e. f. g. h. ♂ ♀. Java. From Dr. Horsfield's
Collection.

i. j. ♂ ♀. N. India. Presented by Colonel Sykes.

k. l. m. n. o. ♂ ♀. N. India. Presented by Colonel
Buckley.

p. ♂. Borneo. Presented by H. Low, Esq.

The larva and pupa of *Diadema Auge* are figured on Plate V. fig. 9, 9a, taken from the original drawing made by E. L. Layard, Esq., in Ceylon.

329. DIADEMA ANTILOPE, Cramer.

- ♀ Papilio Antilope, *Cramer, Pap. Exot. II. t. 183, f. E.F.*
- Diadema Antilope, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 281, n. 6.*
- Nymphalis Antilope, *Godart, Enc. Méth. IX. p. 397.*
- a. ♂. Java. Presented by the Trustees of the British Museum.

Genus PENTHEMA, E. Doubleday.

PENTHEMA, *E. Doubleday, MS.; Diurnal Lep. t. 39 (1847).*

DIADEMA (PENTHEMA), *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 281 (1850).*

330. PENTHEMA LISARDA, E. Doubleday.

Diadema Lisarda, *E. Doubleday, Ann. Nat. Hist. XVI. p. 233 (1845).*

Penthemis Lisarda, *E. Doubleday, Diurnal Lep. t. 39, f. 3.*

Diadema (Penthema) Lisarda, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 281, n. 9.*

Nymphalis Euphrone, *Westwood, Cabinet Orient. Entomol. p. 55, t. 27, f. 1 (1847).*

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.

c. d. ♂ ♀. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus HESTINA, Westwood.

DIADEMA (HESTINA), *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 281 (1850).*

331. HESTINA ASSIMILIS, Linnaeus Sp.

Papilio assimilis, *Linnaeus, Syst. Nat. II. p. 782 (1767).*

Cramer, Pap. Exot. II. t. 154, f. A. Esper, Pap. Exot. t. 57, f. 1. Fabricius, Ent. Syst. III. pt. I.

*p. 39. Clerck, Icon. t. 16, f. 1, 2. Drury, Ill. I.
t. 17, f. 3, 4.*

Diadema (Hestina) assimilis, *Westwood, in Doubleday
and Hewitson's Diurnal Lep.* p. 281, n. 18.

Hestia assimilis, *Hübner, Verz. bek. Schmett.* p. 15.

Nymphalis assimilis, *Godart, Enc. Méth. IX.* p. 393.

Diadema assimilis, *E. Doubleday, List Lep. Brit. Mus.
pt. I. p. 97.*

a. b. ♂. N. India. Donor unknown.

c. ♀. N. India. Presented by the Trustees of the
British Museum.

332. *HESTINA CONSIMILIS*, *Westwood.*

♀ Diadema (Hestina) consimilis, *Westwood, in Doubleday
and Hewitson's Diurnal Lep.* p. 281, n. 20 (1850).

a. b. ♀ (type). N. India. Presented by Colonel
Buckley.

333. *HESTINA ISA*, *Moore.*

Hestina Isa, n. sp.—Female: UPPER-SIDE, fore-wing dusky-brown, darker and somewhat bluish on apical half; a broad oblique short white patch from middle of costal margin, intersected by three of the veinlets; a whitish mark in middle of discoidal cell, and some narrow longitudinal white marks at the apex; also an indistinct marginal row of small whitish spots: hind-wing dusky-brown, with the middle of the wing to abdominal margin white, intersected by the veins; also a marginal and submarginal row of small bluish-white spots from anal angle. UNDER-SIDE paler brown, and marked as above. Wings shaped as in *Hest. consimilis*; expanse $3\frac{1}{2}$ inches.

a. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

334. *HESTINA PERSIMILIS*, *Westwood.*

Diadema (Hestina) persimilis, *Westwood, in Doubleday
and Hewitson's Diurnal Lep.* p. 281, n. 19 (1850).

a. ♂ (type). N. India. Presented by Colonel
Buckley.

b. c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

335. *HESTINA NAMA*, Boisduval Sp.

Diadema Nama, *Boisduval*, MS. E. *Doubleday, Ann. Nat. Hist.* XVI. p. 232 (1845); id. *Diurnal Lep. t. 39, f. 2*; *List Lep. Brit. Mus. pt. I. p. 97.*

Diadema (Hestina) Nama, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 281, n. 21.

a. b. c. d. ♂ ♀. N. India. Presented by Lieutenant Hugo James.

e. f. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus Nov. *CALINAGA*, Moore.

Head small, with a somewhat long frontal tuft.

Thorax large, hairy; neck covered with long velvety hairs.

Abdomen long, slender.

Labial palpi porrected, ascending, elevated to the level of the middle of the eyes, slender, covered with fine hairs, those in front very long; apical joint ovate-cylindric.

Antennæ (broken off).

Fore-legs short, pectoral; femur clothed with fine long hairs; tibia and tarsus hairy; tarsus two-thirds the length of tibia, armed with a single claw.

Hind-legs rather stout; femur clothed with hairs; tibia the same length as the femur, and partially covered with short spiny hairs, with two short tibial spurs; tarsus with several rows of shorter spines; of five joints, the first as long as the other four together; second, third, and fourth short; fifth rather long, half the length of the first; claws sharp.

Fore-wings elongate-trigonal: anterior margin slightly arched, apical angle rounded; exterior margin about half the length of the anterior; posterior margin slightly convex, two-thirds the length of anterior. Costal vein strong, and extending to two-thirds of the anterior margin: subcostal vein separated from the costal, its first branch arising at two-fifths from the base; second branch at a little distance off, and at equal distance from the first as from the extremity of discoidal cell, and both running parallel with the costal: third and fourth branches arising at equal distances from each other and from the extremity of the discoidal cell and the tip of the wing. Upper disco-cellular vein very short, almost obsolete,

arising from the subcostal at about the middle of the wing: middle disco-cellular vein four times the length of first, extending downward, and slightly convex; lower disco-cellular vein long, curving obliquely downward and outward, and uniting with the third branch of the median vein at its origin, closing the discoidal cell.

Hind-wings elongate-trigonal: anterior margin convex at its base, then quite straight; outer margin rounded; abdominal margin nearly straight, slightly rounded: precostal vein rather indistinct, curved inward. Costal vein arched near the base, then straight, extending to costal margin on three-fourths of the wing: subcostal vein straight to two-fifths of the wing, where it branches slightly upward, and extends to the apex. Upper disco-cellular vein short, straight, arising at the subcostal branch; middle disco-cellular vein slightly oblique, concave; lower disco-cellular vein oblique, convex, uniting with the third branch of median vein a little beyond its origin, closing the discoidal cell.

336. *CALINAGA BUDDHA*, Moore (Plate IIIa, f. 5).

Calinaga Buddha, n. sp.—*UPPER-SIDE* smoky-brown: *fore-wing* with a submarginal row of rounded creamy-white spots; also a parallel row of irregular-shaped longitudinal whitish streaks from costal margin, widening to posterior margin; space within discoidal cell whitish, divided transversely in the middle by a blackish bar: *hind-wing* with a submarginal row of white oval spots; another row from anterior margin narrowing towards anal angle; interior of discoidal cell and abdominal margin broadly whitish, the latter tinged at the anal angle with ochreous; thorax ferruginous; abdomen black. *UNDER-SIDE*: *fore-wings* as in the upper-side, but the markings less defined; *hind-wings* dull ochreous, paler at the base and on abdominal margin; whitish markings rather indistinct; sides of thorax ferruginous. Expanse of wings $3\frac{1}{2}$ inches.

a. ♂. N. India. Presented by Colonel Buckley.

Genus NEPTIS, Fabricius.

NEPTIS, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 282, 1808.)
Horsfield. Westwood, in Doubleday and Hewitson's Diurnal
Lep. p. 270.

ACCA, p. Hübner, Verz. bek. Schmett. (1816).

LIMENITIS, p. Boisduval, Ind. Méth. Eur. Lép. p. 16 (1840). E.
Doubleday, List Lep. Brit. Mus.

337. *NEPTIS HORDONIA*, Stoll Sp.

Papilio Hordonia, Stoll, *Suppl. Cramer Pap. Exot.* t. 33, f. 4, 4D (1791).

Neptis Hordonia, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 271, n. 3.

Nymphalis Hordonia, Godart, *Enc. Méth.* IX. p. 429.

Limenitis Hordonia, E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 93.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. d. ♂ ♀. N. India. Presented by Colonel Sykes.

e. f. ♂ ♀. Darjeeling. From Indian Collection, *Exposition Universelle at Paris*, 1855.

Remark.—*Neptis Hordonia* may be known by the mottled markings of the under-side.

338. *NEPTIS HELIODORE*, Fabricius Sp.

Papilio Heliodore, Fabricius, *Ent. Syst.* III. pt. I. p. 130 (1793). Jones, *Icon.* IV. t. 76, f. 2, nec. *Cramer*.

Neptis Heliodore, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 271, n. 4.

Nymphalis Heliodore, Godart, *Enc. Méth.* IX. p. 429.

Limenitis Heliodore, E. Doubleday, *List Lep. Brit. Mus.* pt. I. p. 93.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

Remark.—*Neptis Heliodore* may be distinguished from the preceding by the well-defined markings of the under-side.

339. *NEPTIS MIAH*, Moore (*Plate IVa, fig. 1*).

Neptis Miah, n. sp.—UPPER-SIDE brown-black: *fore-wing* with a longitudinal streak from base of wing, an oblique transverse short apical fascia, and which nearly meets a reversely-oblique fascia on posterior margin, rufous: *hind-wing* with a nearly straight broad inner band, and a narrow submarginal band, rufous. UNDER-SIDE dark ferruginous: *fore-wing* with the longitudinal and oblique marks pinky-white; two narrow submarginal lines purple: *hind-wing* with inner band pinky-white; two submarginal and a less distinct middle

line purple; costal margin at the base whitish. Expanse of wings $2\frac{2}{3}$ inches.

a. b. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—Allied to the two preceding species, but may easily be distinguished by the colour and markings of the under-side.

340. *NEPTIS VIKASI*, Horsfield.

Neptis Vikasi, Horsfield, Cat. Lep. Mus. E.I.C. t. 5, f. 2, 2a (1829).

Athyra Vikasi, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 274, n. 10.

Limenitis Vikasi, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 94.

a. b. c. d. Java (type). From Dr. Horsfield's Collection.

e. f. N. India. Presented by Colonel Buckley.

g. h. Darjeeling. Reserved from Indian Collection, Exposition Universelle at Paris, 1855.

341. *NEPTIS MANASA*, Moore (Plate IVa, fig. 2).

Neptis Manasa, n. sp.—UPPER-SIDE dull dusky-brown: markings white, tinged with very pale ferruginous: *fore-wing* with the discoidal streak long, and continued in a curve to near the posterior angle; an apical oblique streak; two small spots on costal margin, and one below the margin; a spot on middle of posterior margin; a marginal row of small indistinct spots: *hind-wing* with broad inner and narrower outer band, both extending across the abdominal margin; an indistinct marginal line and line between the two bands; narrow ciliae white. UNDER-SIDE pale ochreous, with indistinctly-defined white markings; the discoidal streak is continued uninterrupted to the middle of posterior margin, the space within being blackish; on the hind-wing are some small white markings between the inner band and costal vein. Expanse $2\frac{1}{2}$ inches.

a. ♂. N. India. Presented by Colonel Buckley.

Remark.—*Neptis Manasa* may be distinguished above by the discoidal streak being nearly confluent with the spot on the middle of posterior margin, it being quite confluent on the under-side.

342. *NEPTIS ANANTA*, Moore (Plate IVa, fig. 3).

Neptis Ananta, n. sp.—UPPER-SIDE brownish-black; markings ferruginous: *fore-wing* with discoidal streak straight; a curved twice-interrupted band from anterior margin near apex to middle of posterior margin; a pale marginal line: *hind-wing* with rather broad inner and narrow outer band, both extending across the abdominal margin; also a pale marginal line and line between the bands. UNDER-SIDE very deep ferruginous: *fore-wing* with discoidal streak, spots near apex, and from posterior margin, ferruginous-white; posterior margin broadly patched with black; a marginal and submarginal bluish-ashy line: *hind-wing* with inner band white, outer band grey and indistinct; a marginal line and zigzag line between the two bands bluish-ashy. Expanse $2\frac{1}{2}$ inches.

a. ♀. N. India. Presented by Colonel Buckley.

Remark.—*Neptis Ananta* may be known from *N. Manasa* by its ferruginous markings and straight discoidal streak.

343. *NEPTIS RADHA*, Moore (Plate IVa, fig. 4).

Neptis Radha, n. sp.—UPPER-SIDE brownish-black: markings ferruginous: *fore-wing* with a long discoidal streak, narrow at base, and extending to a short distance between the third and second median veinlets; two small spots on costal margin, and two larger oblique spots near the apex; a large spot on disc, and an elongate spot to middle of posterior margin; marginal line pale: *hind-wing* with rather broad inner and narrower outer band, both extending across the abdominal margin; a marginal line and line between the two bands pale brown. UNDER-SIDE ferruginous-brown; markings as above, but indistinct, owing to the surface being mottled over with ashy-blue. Expanse $2\frac{7}{8}$ inches.

a. ♀. Bootan. From Pemberton's Collection.

b. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

344. *NEPTIS COLUMELLA*, Cramer Sp.

Papilio Columella, Cramer, Pap. Exot. IV. t. 296, f. A.B. (1782).

Neptis Columella, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 272, n. 7.

Limenitis Columella, *E. Doubleday, List Lep. Brit. Mus.*

pt. I. p. 95.

Aeca Columena, *Hübner, Verz. bek. Schmett. p. 44.*

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

b. c. ♀. N. India. Presented by Colonel Sykes.

Inhabits also China, Ceylon; and specimens from the island of Lombok which I have examined, are also identical.

345. *NEPTIS JUMBAH*, Moore (Plate IVa, fig. 5).

Neptis Jumbah, n. sp.—UPPER-SIDE smoky-black: *fore-wing* with a narrow longitudinal discoidal streak, and a parallel triangular spot, white; a transverse curved row of interrupted white spots from apical third of costal margin to near middle of posterior margin, composed of two very narrow longitudinal spots on the costal margin, beneath which are two large spots, then a very small streak, then two large spots, and lastly two narrower spots terminating on posterior margin; between this band of spots and exterior margin are two rows of more or less-defined lunular-shaped white spots, these being bounded exteriorly by a shade of deep black: *hind-wing* with a broad white band from costal to abdominal margin, being intersected by the veins; near outer margin a row of ill-defined white lunated marks, bounded exteriorly by deep black, then a submarginal row of black lines; between the white band and row of white lunated marks is a broad band of black. UNDER-SIDE ferruginous: *fore-wing* with markings as above, but the four rows of markings along exterior margin white, and suffused at the apex and near the middle with ferruginous: *hind-wing* with the broad white band; abdominal margin, base of costal margin, and broadly across parallel with the band, suffused with white; a marginal and two submarginal rows of whitish marks, between which and the broad band a series of fine dark ferruginous spots from abdominal margin, and terminating in white marks on costal margin. Body above, black; beneath, white. Sexes alike. Expanse $2\frac{3}{8}$ to $2\frac{1}{2}$ inches.

a. b. ♂. N. India. Presented by Colonel Sykes.

c. ♀. N. India. Presented by Colonel Buckley.

d. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

e. Ceylon. Donor unknown.

346. *NEPTIS NATA*, Moore (Plate IVa, fig. 6).

Neptis Nata, n. sp.—UPPER-SIDE deep black; markings very white: *fore-wing* with the discoidal streak long, in two portions; curved series of spots small: *hind-wing* with the bands narrow. UNDER-SIDE dusky-brown; inner band of hind-wing not extending to anterior margin; a marginal line *only* between outer band and exterior margin, the submarginal line being obsolete. Sexes alike. Expanse $2\frac{2}{5}$ inches.

a. ♂. Borneo. Presented by Edwin Shepherd, Esq.,
1857.

347. *NEPTIS NANDINA*, Moore (Plate IVa, fig. 7).

Neptis Nandina, n. sp.—UPPER-SIDE black; markings white: *fore-wing* with discoidal streak in two portions; the curved row of seven spots broadly interrupted; and submarginal row of small less-defined spots: *hind-wing* with inner band, and narrow outer row of less-defined square spots; indistinct marginal line and line between the bands, brown. UNDER-SIDE deep chocolate-brown, with markings as above, but more defined and of a brighter colour; those of the hind-wing gently curved. Sexes alike. Expanse $2\frac{1}{4}$ to $2\frac{5}{8}$ inches.

a. ♂. Java. From Dr. Horsfield's Collection.

b. c. d. ♂ ♀. Darjeeling. From Indian Collection,
Exposition Universelle at Paris, 1855.

Remark.—This species may be known from *Neptis Aceris* by the deep chocolate-colour of the under-side, by the anterior margin of hind-wing being evenly arched to beyond the middle; all the markings of this wing being nicely curved.

348. *NEPTIS ACERIS*, Esper Sp.

Papilio Aceris, Esper, Ausl. Schmett. t. 81, cont. 31,
f. 3, 4, t. 82, f. 1 (1785). Fabricius, Mant. Ins. II.
p. 52; Ent. Syst. III. pt. I. p. 245.

Neptis Aceris, Horsfield, Cat. Lep. Mus. E.I.C. t. 7, f. 9.
Westwood, in Doubleday and Hewitson's Diurnal
Lep. p. 271, n. 5.

Nymphalis Aceris, Godart, Enc. Méth. IX. p. 430.

Acca Aceris, Hübner, Verz. bek. Schmett. p. 44.

Limenitis Aceris, Boisduval, Icon. Hist. t. 18, f. 2; id.

Ind. Méth. p. 16. *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 95. *Kollar, in Hügel's Kaschmir, IV. pt. II.* p. 428.

Papilio Plautilla, *Hübner, Europ. Schmett. Pap. f.* 99, 100 (1805).

Acca Matuta, *Hübner, Verz. bek. Schmett. p. 44* (1816).

Neptis Matuta, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 272, n. 6.*

Papilio Leucothoë, *Cramer, Pap. Exot. IV. t. 296, f. E.F. (nec. Linn.)*

Papilio Leucothoë, *apud Donovan, Ins. of China, 1st edit. t. 35, f. 3.*

Limenitis Eurynome, *Westwood, in 2nd edit. Donovan's Ins. of China, p. 66, t. 35, f. 4 ().*

Athyma Eurynome, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 274, n. 2.*

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. h. i. j. ♂ ♀. Europe. From M. Becker's Collection.

k. ♂. N. W. India. Presented by Colonel Hearsey.

l. m. ♂ ♀. Canara. Presented by S. N. Ward, Esq.

n. o. ♂. Darjeeling. From Pearson's Collection.

p. q. r. s. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

t. u. Borneo. Presented by H. Low, Esq.

v. w. ♂ ♀. China. From the Asiatic Society of Bengal.

x. ♂. Pinang. Presented by Dr. Cantor.

The larva and pupa of *Neptis Aceris* are figured on Plate V., fig. 10, 10a, from Java, where it "feeds on a species of *Hedysarum*, bearing the native name of *Kajangan*. May. Rather scarce."—(Horsfield, MS. Notes.)

Genus ATHYMA, Westwood.

ATHYMA, *Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 272 (1850).*

BIBLIS, *p.* *Fabricius*, *Syst. Gloss.* *Horsfield*, *Desc. Cat. Lep. Mus. E.I.C. t. 8, f. 3* (1829).

ACCA, *p.* *Hübner*, *Verz. bek. Schmett.*

LIMENITIS, *p.* *E. Doubleday*, *List Lep. Brit. Mus.*

349. ATHYMA LEUCOTHOË, *Linnæus* Sp.

Papilio Leucothoë, *Linnæus*, *Mus. Lud. Ulr.* 292; *Syst. Nat. ed. 12*, II. p. 780. *Fabricius*, *Ent. Syst. III. I.* p. 129. *Sulzer*, *Hist. Ins. t. 18, f. 2, 3.*

Athyma Leucothoë, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 273.

Limenitis Leucothoë, *Westwood*, in *Donovan's Ins. of China* (2nd edit.). *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 428. *E. Doubleday*, *List Lep. Brit. Mus. pt. I.* p. 94.

Acca Leucothoë, *Hübner*, *Verz. bek. Schmett.* p. 44.

Nymphalis Leucothoë, *Godart*, *Enc. Méth. IX.* p. 430.

Biblis Leucothoë, *Horsfield*, *Cat. Lep. Mus. E.I.C. t. 8, f. 3.*

Papilio Erosine, *Cramer*, *Pap. Exot. III. t. 203, f. E.F.* *Najas hilaris Erosine*, *Hübner*, *Samml. Exot. Schmett. Band I. tab.*

Papilio Polyzena, *Donovan*, *Ins. of China*, t. 35, f. 4 (1st edit.).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Assam. From McClelland's Collection.

f. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

g. ♂. N. India. Presented by Colonel Buckley.

The larva and pupa of *Athyma Leucothoë* are figured on Plate V., fig. 11, 11a, from Java, where it "feeds on a species of *Phyllanthus*, bearing the native name of *Dempul-lolot*. April."—(Horsfield, MS. Notes.)

Remark.—The specimens of *Athyma Leucothoë* from Java are smaller than those from India, and have the central band broader and the portions closer together. Sexes alike.

Captain Mortimer J. Slater remarks, in his MS. Notes, p. 152: "This insect and its allies hover and flit in the air, especially in little open places, as where two paths meet, and over pools of water, &c."

350. *ATHYMA ASURA*, Moore (Plate V., fig. 1).

Athyma Asura, n. sp.—UPPER-SIDE smoky-brown; markings creamy-white: *fore-wing* with a narrow discoidal streak, which is terminated at a short distance by an angular mark; a band of spots curving outward from anterior to middle of posterior margin, the first spot commencing as a very narrow line, second and third long and oval, fourth the smallest, fifth somewhat larger and rounded, sixth larger still and oval, seventh the largest, square, indented at the side, eighth narrow and on posterior margin; a submarginal row of well-defined lunular marks, terminated on the apex of the wing by an inner row of three small spots: *hind-wing* with a broad inner band; also a less broad band from abdominal to anterior angle, this being intersected by the veinlets, and having a single black spot in the middle between each veinlet; a marginal pale-brown line in both wings; a narrow bluish-white collar and band across the base of the abdomen. UNDER-SIDE bright ferruginous; markings as above, but the *fore-wing* has the submarginal row of marks broad, and having a black spot in the middle of each; also a marginal row of small spots; some black lines bordering the discoidal marks; also a small black circle near base of wing, and a patch of black on posterior margin near the angle: *hind-wing* with the inner, and spotted outer band the same as above; a bluish-green curved line across the base of wing, and a marginal row of lunular spots; body and upper part of abdominal margin bluish-green. Expanse 3 inches. Sexes alike.

a. ♀. N. India. Presented by Colonel Buckley.

Remark.—This species may be distinguished from all others by the outer band on the hind-wings having a central spot between each vein. Specimens of both sexes of *Ath. Asura* are in the Collection of W. W. Saunders, Esq.

351. *ATHYMA OPALINA*, Kollar Sp. (Plate V., fig. 2).

Limenitis Opalina, Kollar, in Hügel's Kaschmir, IV.
pt. II. p. 427 (1844).

Athyma Opalina, Westwood, in Doubleday and Hewitson's
Diurnal Lep. p. 274, n. 5.

a. ♂. N. India. From the Asiatic Society of Bengal.

b. ♂. N. India. Presented by Colonel Buckley.

c. ♂. N. India. Presented by Colonel Hearsey.

d. e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

f. g. ♀. N. India. Presented by Edwin Shepherd, Esq., 1857.

Athyma Opalina may be distinguished by the discoidal streak being divided into four portions, as in *Ath. Leucothoë*, and by the deep ferruginous colour of the under-side being suffused in parts with grey and greyish-brown. Expanse $2\frac{2}{5}$ to $2\frac{7}{8}$ inches. Sexes alike.

352. *ATHYMA LARYMNA*, E. Doubleday Sp.

Limenitis Larymna, E. Doubleday MS.

Athyma Larymna, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 274, n. 7, t. 35, f. 1 (1850).

a. ♂. Borneo. Presented by Edwin Shepherd, Esq., 1857.

353. *ATHYMA JINA*, Moore (Plate V., fig. 3).

Athyma Jina, n. sp.—Male: UPPER-SIDE smoky-brown; markings creamy-white: *fore-wing* with the discoidal streak entire, long, broad, and thickening to the extremity; a series of seven spots from subcostal vein curving outward to middle of posterior margin, the first being small, second larger, broader, third narrow, fourth small, fifth larger, oval, sixth the largest, somewhat square and indented at the side, seventh narrow, elongated, triangular; a submarginal row of rather indistinct spots, those on the apex broadest: *hind-wing* with inner band somewhat narrow; outer band composed of broad lunulated spots; band across base of abdomen whitish. UNDER-SIDE brilliant ferruginous, posterior margin of *fore-wing* blackish; markings the same as above, but the *fore-wing* with an additional spot on costal margin to the curved row; extreme posterior margin blackish, bounded inwardly by a submarginal row of white lines: *hind-wing* with the inner band extending across abdominal margin; space between base of wing and costal vein white; extreme exterior margin blackish, bounded inwardly by a marginal row of narrow lunular marks; body white. Expanse 3 inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—*Athyma Jina* may at once be distinguished by the discoidal streak being entire.

354. *ATHYMA PRAVARA*, Moore (Plate V., fig. 4).

Athyma Pravara, n. sp.—UPPER-SIDE smoky-brown; markings creamy-white: *fore-wing* with an *entire* club-shaped streak; a transverse row of spots curving outward from costal vein towards the apex to middle of posterior margin, the first and second spots being rather large and square, the third much smaller, the fourth a mere dot, the fifth the largest, and oval, the sixth large and indented at the sides, the last narrow; a row of submarginal linear-shaped spots, more or less distinct: *hind-wing* with a broad inner band, and a narrow band of spots curving outwardly from anal angle to anterior angle; a submarginal line and abdominal margin light brown; body with a narrow bluish-white band across the thorax, and another across the base of the abdomen. UNDER-SIDE paler brown, with markings as above, but with dark-brown patches between the markings; body and abdominal margin greyish. Sexes alike. Expanse $2\frac{1}{8}$ inches.

a. ♂. Borneo. Presented by Edwin Shepherd, Esq.,
1857.

b. ♀. Java. From Dr. Horsfield's Collection.

355. *ATHYMA NEFTE*, Cramer Sp.

♀ *Papilio Nefte*, Cramer, Pap. Exot. III. t. 256, f. E.F. (1782).

Athyma Nefte, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 274, n. 11.

Pantoporia Nefte, Hübner, Verz. bek. Schmett. p. 44.

Nymphalis Nefte, Godart, Enc. Méth. IX. p. 429.

Limenitis Nefte, Boisduval, Spéc. Gén. Lép. t. 8 (4B), f. 6. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 93.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. g. ♂ ♀. Borneo. Presented by Edwin Shepherd, Esq., 1857.

Remark.—The male of *Athyma Nefte* has all the markings of the upper-side white, these in the female being orange-coloured.

356. *ATHYMA INARA*, E. Doubleday, Sp.

♂ *Limenitis Inara*, E. Doubleday MS.

Athyma Inara, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 274, t. 34, f. 3 (1850).

- a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.
- d. ♀. N. India. Presented by Colonel Sykes.
- e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—The female of *Athyma Inara* has all the markings on the upper-side orange-colour.

357. ATHYMA CAMA, Moore (Plate V., fig. 5, ♂ ♀).

Athyma Cama, n. sp.—Male: UPPER-SIDE velvety-black; fore-wing with brown marginal and submarginal line; discoidal streak indistinct ferruginous, and dusted over with black; near the apex a ferruginous spot; two oblique spots from subcostal vein on one-third of the wing from the apex, and a band of three spots from middle of wing to posterior margin, joining a band across hind-wing, white, bordered with blue; on the hind-wing also a marginal and submarginal brown line; body black, with a broad white band across base of abdomen. UNDER-SIDE pale ferruginous; markings purplish-white; the curved band as above: fore-wing with a long broad irregular discoidal streak, submarginal and indistinct marginal line; a black patch near the base of wing and posterior angle: hind-wing with a curved line near base of wing, and a rather broad outer band; a marginal row of marks; some patches of brown between outer and inner band; body and abdominal margin broadly grey. Female: UPPER-SIDE black; markings ferruginous: fore-wing with a long discoidal streak; an oblique band from anterior to near middle of exterior margin, and nearly joining a band running to middle of posterior margin; a marginal and submarginal brown line, the latter ferruginous anteriorly and posteriorly: hind-wing with a broad inner and narrower outer band; also a marginal brown line; body black, with a white band across base of abdomen, and bound below with ferruginous; also a pale-white collar. UNDER-SIDE ferruginous; markings as in male, but pinky-white, except marginal and submarginal lines, which are purplish-white. Expanse of wings in male $2\frac{1}{2}$ to $2\frac{7}{8}$ inches; female, 3 inches.

- a. b. ♂. Darjeeling. From Pearson's Collection.
- c. ♂. Darjeeling. Presented by Colonel Buckley.
- d. e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—The male of *Athyra Cama* may at once be known from that sex of *Athyra Selenophora* in having on the upper-side a ferruginous spot close to the apex, and the band on the fore-wing being composed of three spots.

358. *ATHYMA SELENOPHORA*, Kollar Sp.

Limenitis Selenophora, Kollar, in *Hügel's Kaschmir*, IV.
pt. II. p. 426, t. 7, f. 1, 2 (1844). Westwood, in
"Doubleday and Hewitson's Diurnal Lep. p. 276,
n. 8.

- a. ♂. N. India. Presented by Colonel Buckley.
- b. c. d. ♂. Darjeeling. From Indian Collection,
Exposition Universelle at Paris, 1855.
- e. f. ♂ var. (with discoidal streak in two portions).
Darjeeling.

Athyra Selenophora.—The male may be distinguished from *Athyra Cama* on the upper-side by having the oblique sub-apical spots narrow, and the band on the fore-wing being composed of four spots; in the under-side being of a darker ferruginous colour, and the streak along discoidal cell being divided into four portions, and in having darker black blotches; also in having on the hind-wing four short black lines disposed between the inner band and discoidal vein. Expanse of male $2\frac{1}{2}$ inches. Female unknown, but would have the markings above ferruginous.

359. *ATHYMA RANGA*, Moore (Plate V., fig. 6).

Athyra Ranga, n. sp.—UPPER-SIDE smoky-black: *fore-wing* with a curved interrupted white band from anterior margin, one-third from the apex, to middle of posterior margin, being composed of eight spots, the first on anterior margin, very narrow, second, third, and fourth elongate-conical, the fourth being the shortest, fifth triangular and broadly divided from the fourth, sixth broad, largest, and nearly square, seventh narrower and broadly divided from the sixth, eighth long and narrow; a marginal and submarginal row of rather indistinct whitish spots; base of wing covered with indistinct white spots: *hind-wing* with broad inner band, intersected by the veins; an outer or submarginal row of broad conic-shaped indistinct white spots; a marginal row of very indistinct spots; also indistinct spots at the base of wing; abdominal margin whitish; body dark brown, abdomen with two rows of small white spots. UNDER-SIDE with the markings

the same, and very distinct; body and abdominal margin greenish-white. Expanse $2\frac{3}{8}$ inches. Sexes alike.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Specimens of both sexes are in the Collection of W. W. Saunders, Esq.

360. *ATHYMA MAHESA*, Moore (Plate V., fig. 7).

Athyma Mahesa, n. sp.—UPPER-SIDE smoky-black, *fore-wing* with a green gloss in some lights: *fore-wing* with three oblique white ovate spots from subcostal vein, one-third from the apex; two white spots in middle of the wing, the upper one very small, and two white spots on middle of posterior margin, the upper oval, the lower narrow; two indistinct white spots in discoidal cell, and some indistinct greenish spots at the base of the wing; a marginal and submarginal row of light-brown spots: *hind-wing* with an inner white band, divided by the veins, and an outer or submarginal row of indistinct brownish-white conic-shaped spots; also a very indistinct marginal row of small light-brown spots; body brown, abdomen with two rows of white spots. UNDER-SIDE paler, tinged with ferruginous about the disc; markings the same, but all very distinct, and more or less white; the spots within discoidal cell and base of *fore-wing* divided by black marks; base of costal margin yellowish-white: *hind-wing* with space between precostal and costal vein yellowish-white; a curved oval black mark, whitish within, between the costal vein and inner band; a row of black patches between inner and outer band; body, and broadly on abdominal margin, yellowish-grey. Expanse $2\frac{7}{8}$ inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—*Athyma Mahesa* may be known from the preceding by its larger size and much narrower band.

Genus Nov. *ABROTA*, Moore.

Head of moderate size, tufted in front.

Eyes prominent, naked.

Antennæ long, filiform at the base, gradually thickening to a slender cylindric club.

Labial palpi elevated above the level of the eyes, hairy, those at the base, side, and in front, long.

Thorax robust, clothed with long hairs.

Abdomen of moderate size.

Fore-wings elongate-trigonal; anterior margin boldly curved; apical angle slightly rounded; outer margin somewhat straight in the male, concave in the female, slightly waved, two-thirds the length of the anterior; posterior margin slightly concave, rather longer than the outer; costal vein thick at the base, extending to beyond the middle of the costa; subcostal vein with the first branch arising about one-fourth the length of the wing, second a little beyond it, and before the extremity of the discoidal cell, third branch arising at two-thirds the length of the wing, and extending in a slight curve to the tip, fourth branch straight, arising half-way between it and the apex, which it reaches just below the tip; the terminal portion of the vein deflexed; upper disco-cellular vein very short, outwardly oblique, arising at one-third from the base of the wing; middle disco-cellular vein curved outward; lower disco-cellular vein long, curved outward, and uniting with the third branch of the median vein at a little distance beyond its origin, closing the cell at a little more than one-third the length of the wing.

Hind-wings somewhat rounded; anterior margin arched at the base, from thence nearly straight to the apex, where it curves slightly downward; exterior margin rather longer than the anterior, rounded and waved; precostal vein curved outward; costal vein curved, extending to the apex; subcostal vein branching near its base; upper disco-cellular vein arising at a short distance from the base of the subcostal branch; lower disco-cellular vein obsolete, so that the cell is open.

Fore-legs of the male slender; femur scaly in front, beneath clothed with fine long silky hairs; tibia as long as the femur, clothed with downy hairs; tarsus two-thirds the length of tibia, conical, covered with uniform hairs.

Fore-legs of the female scaly; femur with a few silky hairs beneath; tarsus rather longer than the male, truncated at the tip, where it is finely spined beneath.

Four hind-legs moderately long, of equal length; tibia shorter than the femur; femur thickly scaled; tibia thickly clothed with short scaly hairs, with two rows of short thick spines; tarsus as long as the tibia, thickly clothed at the sides and beneath with rows of short spines; claws sharp, and much curved.

361. *ABROTA GANGA*, Moore (Plate VIa, fig. 1, ♂ ♀).

Abrota Ganga, n. sp.—Male: UPPER-SIDE ferruginous: fore-wing along upper part of discoidal cell, a spot, and disco-cellular mark within it, a broad oblique streak, tapering from middle of costal vein to middle of lower discoidal veinlet, then extending in a zigzag manner to posterior margin near the base, the space between the latter and median vein only smeared, exterior and apical margin, submarginal line, and smearable patches near the apex and lower part of disc, black: hind-wing with marginal, submarginal, third and fourth inner bands, black; ciliae with small white spots. UNDER-SIDE dull-yellow, with duller exterior margins; a distinct dark narrow streak from near apex of fore-wing to near abdominal margin; two indistinct zigzag lines across disc, and markings about the base of both wings dull-ferruginous; from apex of fore-wing some white spots. Female black-brown: fore-wing with streak from base along lower part of discoidal cell to middle of wing, with a dentate mark on disco-cellular vein; two outward oblique spots from near apex; a broad band from middle of disc, inward oblique, to posterior margin, and indistinct submarginal series of small spots, pale ferruginous; also two small outward oblique white spots at the apex: hind-wing with broad inner and narrower outer slightly-curved bands, pale ferruginous; ciliae with small white spots. UNDER-SIDE pale chocolate-brown, with patch on costal margin near the apex, space about the disc of fore-wing and across middle of hind-wing, yellowish; an ill-defined streak from near apex of fore-wing to middle of abdominal margin; zigzag lines across disc, and patches about base of both wings, whitish; some spots at apex of fore-wing white. Expanse of male 3, female 3 $\frac{5}{8}$ inches.

a. b. c. d. e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus LIMENITIS, Fabricius.

LIMENITIS, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 281, 1807).

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 274.

NYMPHALIS, Boisduval, Ind. Méth. Eur. Lép. p. 16 (1840). E.

Doubleday, List Lep. Brit. Mus. pt. I. p. 96.

ACCA, p. Hübner, Verz. bek. Schmett.

LIMENITIS, p. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 96.

NEPTIS, p. Horsfield, Cat. Lep. Mus. E.I.C. t. 7, f. 8.

BIBLIS, p. Horsfield; id. t. 8, f. 4.

362. *LIMENITIS PROCRIS*, Cramer Sp.

Papilio Procris, Cramer, Pap. Exot. II. t. 106, f. E.F. (1779). *Fabricius*, Ent. Syst. III. pt. I. p. 138.

Limenitis Procris, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 97. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 276, n. 7.

Nymphalis Procris, Godart, Enc. Méth. IX. p. 404.

Biblis Procris, Horsfield, Cat. Lep. Mus. E.I.C. t. 7, fig. 8.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection. *par. simil. & less colored than Cramer*

e. f. ♂ ♀. N. India. Presented by Colonel Sykes.

g. h. ♂. N. India. Presented by Colonel Buckley.

The larva and pupa of *Limenitis Procris* are figured on Plate V., fig. 12, 12a, from Java, where it "feeds on a species of *Nuuclea*, bearing the native name of *Kleppu*. June."—(Horsfield, MS. Notes.)

363. *LIMENITIS ZULEMA*, E. Doubleday.

Limenitis Zulema, E. Doubleday, MS. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 276, n. 2, t. 34, f. 1 (1850).

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

364. *LIMENITIS ALANKARA*, Horsfield Sp.

Aconthea Alankara, Horsfield, Cat. Lep. Mus. E.I.C. t. 5, f. 6 (1829).

Limenitis Alankara, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 276, n. 14.

a. ♂. Java. From Dr. Horsfield's Collection.

Remark.—This species also inhabits Malacca and Sumatra.

365. *LIMENITIS PADUKA*, Moore.

Limenitis Paduka, n. sp.—This is larger than *Lim. Alankara*, and has the upper-side darker: fore-wing with the white patch at the apex larger, the marginal zigzag markings thicker, and with blackish

markings between: on the *hind-wing* the fascia terminates very narrowly on the abdominal margin, where it is bluish; the marginal zigzag markings are less defined and wider apart, and with blackish markings between. In the female, the wings are more rounded than in the other sex. Expanse 2 $\frac{3}{8}$ inches.

a. b. ♂ ♀. Borneo. Presented by Hugh Low, Esq.

366. *LIMENITIS ISMENE*, E. Doubleday.

♂ *Limenitis Ismene*, E. Doubleday, MS. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 276, n. 10, t. 34, f. 2 (1850).

a. b. c. ♂. Silhet. Presented by Colonel Buckley.

367. *LIMENITIS DARAXA*, E. Doubleday.

Limenitis Daraxa, E. Doubleday, MS. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 276, n. 11, t. 34, f. 4 (1850).

a. b. c. d. ♂ ♀. Darjeeling. From Pearson's Collection.

e. f. ♂. N. India. Presented by Lieutenant Hugo James.

368. *LIMENITIS DUDU*, Westwood.

Limenitis Dudu, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 276, n. 13 (1850).

a. ♂. N. India. Presented by Lieutenant Hugo James.

b. c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

e. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

369. *LIMENITIS DANAVA*, Moore (Plate VIa, fig. 2, ♂ ♀).

Limenitis Danava, n. sp.—Male: UPPER-SIDE dark olive-brown from base to disc of fore-wing, and from base to one-third of hind-wing, the rest being pale olive-brown; a transverse dark band crossing the disc of fore-wing and middle of hind-wing; a dark submarginal wavy line and two indistinct inner wavy lines crossing both wings; some indistinct markings within the discoidal cell with paler centres,

and a small green patch at anal angle. UNDER-SIDE of a golden-glossed obscure ashy-brown ochreous colour: *fore-wing* with markings at the base; a transverse row of lunulated marks, and an indistinct submarginal row of very small spots, purple-ashy; also a whitish patch at the apex, and a narrow yellowish line across the disc: *hind-wing* with broad inner and submarginal band purple-ashy, the latter with small whitish spots along its middle; two marks within and one above the discoidal cell whitish. Female: UPPER-SIDE with portions at the base, as in male, dark brownish-green; the middle of both wings white, with broad greenish-brown band crossing the disc; exterior margins broadly paler greenish-brown, with green patch at the anal angle; markings at base of wings pale. UNDER-SIDE pale greenish-yellow, with greenish-golden gloss; bands and markings as in upper-side, but less distinct and whitish. Expanse from 3 to $3\frac{1}{2}$ inches.

- a. b. ♂.* Darjeeling. From Pearson's Collection.
- c. ♂.* Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- d. e. f. ♀.* N. India. Presented by Colonel Buckley.

370. LIMENITIS ZAYLA, E. Doubleday.

Limenitis Zayla, E. Doubleday, MS. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 276, n. 12, t. 35, f. 4 (1850).

- a. b. ♂.* Darjeeling. From Pearson's Collection.
- c. ♂.* N. India. Presented by Colonel Buckley.
- d. e. ♂.* Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus Nov. PANDITA, Moore.

Head rather small.

Eyes prominent, naked.

Labial palpi rather short, directed upwards to about the middle of the eyes, extended a little in front of the head, hairy; those at the base long; apical joint conic.

Antennæ rather longer than half the fore-wing, straight, with an elongated slender club.

Thorax moderately robust, covered with scales, the sides hairy.

Fore-wings elongate-trigonal, anterior margin arched, apical angle

rounded ; exterior margin two-thirds the length of anterior, more or less scalloped ; posterior margin about the same length as exterior, somewhat rounded near the base and the angle ; costal vein moderately strong, extending to the middle of anterior margin ; subcostal vein with its first branch arising at one-fourth of the wing from the base, and extending to the costal margin at one-third from the apex, second branch arising a little beyond the first, some distance before the extremity of the discoidal cell, and terminating one-fourth from the apex, third branch commencing at one-third from, and extending to, the apex, fourth branch at one-fifth of the wing, and terminating a little below the apex ; first disco-cellular vein very short, almost obsolete, commencing obliquely a little beyond the base of the third subcostal branch ; second disco-cellular vein five times the length of the first, descending in an oblique inward curve ; third disco-cellular vein rather slender, but distinct, long, descending straight, but uniting in a curve with the third median veinlet immediately above its origin, thus closing the discoidal cell.

Hind-wings quadrate-trigonal ; the extreme base of the anterior margin arched, the rest of it nearly straight ; exterior margin somewhat rounded, scalloped ; abdominal margin grooved, about the same length as anterior ; precostal vein curved outwardly ; costal vein arched to about the middle, and then straight to the apex ; subcostal vein branching off nearly straight, opposite the junction of the precostal ; first disco-cellular vein straight, oblique, arising a little beyond the subcostal ; lower disco-cellular vein obsolete ; discoidal cell open.

Fore-legs very small, with fine silky hairs ; femur rather longer than the tibia ; tarsus half the length of the tibia, simple : the other legs broken off.

Abdomen rather small, slender, hairy.

Remark.—This genus is somewhat allied to *Limenitis* and *Heterochroa*.

371. *PANDITA SINOPE*, Boisduval Sp. (Plate VIa, fig. 3).

Adolias Sinope, Boisduval, MS. Westwood, in Double-day and Hewitson's *Diurnal Lep.* p. 292, n. 25 (1850).

a. ♂. Java. From Dr. Horsfield's Collection.

Pandita Sinope, n. sp.—UPPER-SIDE bright fulvous : *fore-wing* with the costal margin and basal half of wing golden-brown ; exte-

rior margin, submarginal line, a pale inner parallel line, also a rather broad zigzag band, curving outward across the disc from the apical third of the costal margin to inner margin near angle, and transverse markings within and below discoidal cell, black: *hind-wing* with the base golden-brown; abdominal margin plain dull-brown; a broad exterior marginal band, a submarginal line, an inner broad band, also a rather zigzag ill-defined band across the disc, and markings about the discoidal cell, black, the black colour being darkest on the submarginal lines and discoidal markings. *UNDERSIDE* paler fulvous, the markings the same as in upper-side, but very indistinct; body and abdominal margin grey. Expanse of wings $2\frac{1}{10}$ inches. Sexes alike. Female in British Museum.

Stirps IV., with THYSANURIFORM Larva.

- THYSANURIFORM, *Horsfield*, *Cat. Lep. Mus. E.I.C.* p. 48 (1828).
- RASORIAL, or THYSANURIFORM, *Swainson*, *Cab. Cyclop.* pp. 58, 62 (1840).
- PAP. DANAÏ FESTIVI (*part*), *Linnæus*.
- PAP. NYMPHALES, GEMMATI et PHALERATI (*part*), *Linnæus*.
- LEP. TETRAPODA (*part*), *Latreille*, *Règ. Anim. V.* p. 378 (1829).
- NYMPHALIDÆ (*part*), *Swainson*, *Phil. Mag. I.* p. 187 (1827); id. *Cab. Cyclop.* pp. 86, 92 (1840). *Stephens*, *Brit. Ent. Haust. I.* p. 27 (1827); id. *List Brit. Lep. Brit. Mus. pt. I.* p. 6 (1850). *Westwood*, *Introd. to Ent. II.* p. 353 (1840); id. in *Doubleday and Hewitson's Diurnal Lep.* p. 144 (1852). *Dup. C.* p. 8 (1846). *Stainton*, *Manual Brit. Butt.* pp. 10, 21 (1856).
- NYMPHALIDES (*part*), *Boisduval*, *Spéc. Gén. Lép. I.* p. 165 (1836); id. *Ind. Méth. Eur. Lép.* p. 16. *Blanchard*, *Hist. Nat. des Ins. II.* p. 331.
- NYMPHALII et NYMPHALITES (*part*), *Blanchard*, *Hist. Nat. des Ins. II.* pp. 322, 329, 333 (1845).
- NYMPHALIDI (*part*), *Stephens*, *List Brit. Lep. Brit. Mus. pt. I.* p. 10 (1850). *Stainton*, *Manual Brit. Butt.* pp. 21, 33 (1856).
- APATURIDES, *Boisduval*, *Ind. Méth.* p. 24 (1840).
- MORPHIDES, *Boisduval*, *Spéc. Gén. Lép. I.* p. 166 (1836).
- MORPHITES, *Blanchard*, *Hist. Nat. des Ins. II.* p. 334 (1845).
- MORPHIDÆ, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 332 (1851).
- BRASSOLIDES, *Boisduval*, *Spéc. Gén. Lép. I.* p. 166 (1836).
- BRASSOLITES, *Blanchard*, *Hist. Nat. des Ins. II.* p. 334 (1845).
- BRASSOLIDÆ, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 350 (1851).
- SATYRIDES, *Boisduval*, *Spéc. Gén. Lép. I.* p. 166 (1836); id. *Ind. Méth. Eur. Lép.* p. 25.
- SATYRIDÆ, *Swainson*, *Cab. Cyclop.* pp. 86, 93 (1840). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 352 (1851).
- SATYRITES, *Blanchard*, *Hist. Nat. des Ins. II.* p. 335 (1845).
- SATYRIDI, *Stephens*, *List Brit. Lep. Brit. Mus. pt. I.* p. 6 (1850).
- BIBLIDES, *Boisduval*, *Spéc. Gén. Lép. I.* p. 167 (1836).

BIBLITES, *Blanchard*, *Hist. Nat. des Ins.* II. p. 332 (1845).

EURYTELIDÆ, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 403 (1851).

LIBYTHIDES, *Boisduval*, *Spéc. Gén. Lép.* I. p. 167 (1836); id. *Ind. Méth.* p. 23.

LIBYTHEITES, *Blanchard*, *Hist. Nat. des Ins.* II. p. 332 (1845).

LIBYTHEIDÆ, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 412 (1851).

Genus ADOLIAS, *Boisduval*.

ADOLIAS, *Boisduval*, *Spéc. Gén. Planches*, p. 2, t. 3, f. 2 (1836).
Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 289.

ACONTHEA, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 8, f. 6 (1829); *Zool. Journ.* V. p. 65 (1830).

SYMPHEDRA, EUTHALIA, CYMOCHOË, *pt. Hübner*.

NYMPHALIS, *pt. Godart*.

ITANUS, *E. Doubleday, MS.*

372. ADOLIAS ACONTHEA, *Cramer Sp.*

♂ ♀ Papilio Aconthea, *Cramer, Pap. Exot.* II. t. 134, f. D.E. ♂,
F.G. ♀ (1779).

Adolias Aconthea, *Boisduval*, *Spéc. Gén. Lép. List of Plates*, p. 2, t. 3, f. 2 (1836). *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 103. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 1.

Nymphalis Aconthea, *Godart, Enc. Méth.* IX. p. 383.

Cymothoë Aconthea, *Hübner, Verz. bek. Schmett.* p. 39.

♂ Nymphalis Disconthea, *Godart, Enc. Méth.* IX. p. 384 (1819).

Aconthea primaria, *Horsfield, Cat. Lep. Mus. E.I.C.* t. 8, f. 6 (1829).

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

The transformations of *Adolias Aconthea* are figured on Plate VI., fig. 1, 1a, from Java, and "feeds on the Mango."—(Horsfield, MS.)

373. ADOLIAS PARTA, *Moore*.

Adolias Parta, *n. sp.*—Male: UPPER-SIDE dark brown, with a vinaceous tinge: fore-wing with a paler transverse band, margined broadly on both sides with black, the anterior portion within, from

costal margin on both sides, with a series of small white patches; marks within discoidal-cell black, with dark-brown centres: *hind-wing* with an inner blackish band and outer row of small black spots. **UNDER-SIDE** paler, marked as above. Female: **UPPER-SIDE** pale brown: *fore-wing* with broad whitish curved transverse band, with patch on costa, and dark margins: *hind-wing* with rather broad inner band, and outer zigzag line, the point between each vein with a minute darker dot; space between inner band and base tinged with white; marks at base of wings blackish. **UNDER-SIDE** pale dull ochreous, with paler margins; markings as above, but very indistinct. Expanse of male $2\frac{3}{8}$, female, $2\frac{7}{8}$ inches.

a. ♀. Borneo. Presented by Edwin Shepherd, Esq.,
1857.

The male of *Adolias Parta* is in the Collection of W. W. Saunders, Esq.

374. ADOLIAS GARUDA, Moore.

Adolias Garuda, n. sp.—Male: **UPPER-SIDE** glossy greenish-brown: *fore-wing* with costal margin to its middle, a broad irregular band from thence to posterior margin, black, the latter bordered exteriorly, anteriorly from costal vein, with five white spots; two small white spots on costal margin one-fourth from the apex; exterior margin and submarginal band blackish; within discoidal-cell, first a short line, then two reniform marks, black: *hind-wing* with curved dentate blackish band from middle of anterior margin to near abdominal margin; a submarginal row of small deep-black spots; exterior margin near anal angle blackish; within discoidal-cell some black markings, and two small black spots without, one above, the other below the cell. **UNDER-SIDE** light chocolate-brown, greyish towards the base: *fore-wing* with black marks within discoidal-cell as above, and a small black spot below it; the row of five white spots and the two apical spots as above; from the latter, across the disc to posterior margin, runs a narrow interrupted black band; at the apex and along the exterior margin some bluish-grey spots: *hind-wing* with four lines within discoidal-cell, a small spot, and two oval marks above, black; an indistinct band across the disc; submarginal row of black spots as above; a patch of bluish-grey at anterior angle. Female pale brown, with a greenish gloss: **UPPER-SIDE** with markings as in male, but less defined; the row of white spots of the fore-wing are larger, and the submarginal row of black

spots on the hind-wing are also larger than in the male. UNDER-SIDE as in that of the male. Wings shaped as in *Adol. Aconthea*. Expanse of male $2\frac{3}{8}$, female, $2\frac{7}{8}$ inches.

- a. b. c. ♂ ♀. N. India. Presented by Colonel Sykes.
- d. e. ♂ ♀. N. India. Presented by Colonel Buckley.
- f. g. ♂ ♀. Madras. From Captain J. M. Jones's Collection.

The transformations of *Adolias Garuda* are figured on Plate VI., fig. 2, 2a, from General Hardwicke's original drawings in the British Museum.—“Feeds on *Trophis aspera* and on a species of *Bryonia*. March.”—(Hardwicke.)

The transformations of this species were also observed by A. Grote, Esq., of Calcutta, and are figured among his original drawings. It “feeds upon the Mango.”

375. ADOLIAS PHEMIUS, E. Doubleday Sp.

- ♂ Itanus Phemius, E. Doubleday, MS.
- ♂ Adolias Phemius, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 291, n. 13, t. 41, f. 4 (1850).
- a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Both sexes of *Adolias Phemius* are in the British Museum Collection.

376. ADOLIAS ANOSIA, Boisduval MS.

Adolias Anosia, n. sp.—Male: UPPER-SIDE dark ash-green, with the anterior margin of hind-wing broadly pinky-brown: fore-wing with broad transverse band of ashy-white irroration; black markings within discoidal-cell, and some below it, bordered with ashy-white irroration: hind-wing with ashy-white irroration on lower part of the disc, bordering the discoidal marks, and a spot above and below the cell, also bordering a submarginal row of black spots. UNDER-SIDE creamy ash-colour, palest on the anterior half, which is covered with rather indistinct darker short transverse striae; markings within and about discoidal-cell black; on the hind-wing an indistinct submarginal row of black spots. Female: UPPER-SIDE paler ash-green than the male; exterior margins brownish; markings disposed the same; fore-wing with a curved row of five white spots from

middle of costal margin ; the irrorated band paler and more clearly defined ; an indistinct inward oblique row of black spots from near apex to near middle of posterior margin : *hind-wing* with indistinct black curved band from middle of anterior to middle of abdominal margin ; also an indistinct submarginal row of black spots. *UNDER-SIDE* paler than in the male, marked as upper-side, with the curved row of five spots bordered inwardly with dark brown ; the indistinct oblique row of blackish spots from apex only to middle of the disc, the lower part being suffused with dark brown ; exterior margin dark brown : *hind-wing* with indistinct inner band and submarginal row of large dark-brown spots ; anterior wing in both sexes much falcated. Expanse of male $2\frac{1}{2}$, female, 3 inches.

a. b. c. d. ♂ ♀. N. India. Presented by Colonel Sykes.

377. ADOLIAS ALPHEDA, Godart Sp.

Nymphalis Alpheda, Godart, Enc. Méth. IX. p. 384 (1819).

Adolias Alpheda, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 291, n. 20.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

378. ADOLIAS ADONIA, Cramer Sp.

♀ Papilio Adonia, Cramer, Pap. Exot. III. t. 255, f. C.D. (1782).

Adolias Adonia, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 291, n. 11.

Euthalia Adonia, Hübner, Verz. bek. Schmett. p. 41.

Nymphalis Adonia, Godart, Enc. Méth. IX. p. 400.

♂ Aconthea Lubentina, apud Horsfield, Cat. Lep. Mus. E.I.C. t. 5, f. 5.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

379. ADOLIAS LUBENTINA, Cramer Sp.

♀ Papilio Lubentina, Cramer, Pap. Exot. II. t. 155, f. C.D. (1779). Fabricius, Ent. Syst. III. pt. I. p. 121 (1793). Donovan, Ins. of China, t. 36, f. 3, ♂.

Adolias Lubentina, Boisduval. E. Doubleday, List Lep.

Brit. Mus. pt. I. p. 103. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 291, n. 10.

Euthalia Lubentina, Hübner, Verz. bek. Schmett. p. 41.

Nymphalis Lubentina, Godart, Enc. Méth. IX. p. 400.

a. b. ♂ ♀. N. India. Presented by Colonel Buckley.

c. ♀. N. India. Presented by Colonel Sykes.

d. ♂. N. India. Presented by Edwin Shepherd, Esq.,
1857.

The male of *Adol. Lubentina* is correctly figured by Donovan, and the female by Cramer.

The transformations of *Adolias Lubentina* are figured among the original drawings of A. Grote, Esq., of Calcutta, and will be drawn on a subsequent Plate of this Catalogue.—“Feeds upon *Loranthus*.”

380. ADOLIAS SALIA, Moore.

Adolias Salia, n. sp.—Male: UPPER-SIDE deep dark brown: fore-wing from costal margin near apex to posterior margin near angle, a narrow zigzag white line, margined outwardly with black and then with dull blue; within the anterior portion of this line the ground-colour is paler; indistinct black markings at the base; narrow ciliae spotted with white: hind-wing, from anterior margin near angle, a rather broad white band, slightly curving to abdominal margin near anal angle, and which is margined outwardly with a zigzag black line, the points being inward; anterior angle pale brown; from anal angle outside the band, dull blue; narrow ciliae white. UNDER-SIDE dull ochreous; the transverse band of the fore-wing broad, and partly brown anteriorly, white posteriorly, and both wings margined within with dusky black, and outwardly with zigzag black line; markings at base of wing black; exterior margin dusky; body and abdominal margin greenish-ochreous. Female dull brown: fore-wing with broad tapering transverse white band, intersected by the veins, and suffused more or less with brown anteriorly on the inner portion, and margined outward with a narrow lanceolate black line, the points being inward; base of wing with indistinct black marks: hind-wing with broad white band, intersected by the veins, slightly curving from anterior margin to abdominal margin near anal angle, being margined outwardly with broad lanceolate black marks, the point being inward, these again being narrowly margined outwardly with white; base of wing with indistinct marks; narrow ciliae

spotted with white. UNDER-SIDE ochreous; markings as above, but the bands less distinct; body and abdominal margin greenish-grey. Shape of wings as in *Ad. Aconthea*; expanse of male $2\frac{1}{8}$, female, $2\frac{6}{8}$ inches.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

381. *ADOLIAS PALGUNA*, Moore.

Adolias Palguna, n. sp.—Male: UPPER-SIDE dull brown, dusky about the apex: *fore-wing* with a transverse band of irregular-shaped white spots, more or less suffused with brown, the outer margins of which have lanceolate black marks pointing inward; base of wing with indistinct marks: *hind-wing* with narrower curved white band from anterior margin to anal angle, margined outwardly with a broad lanceolate mark between each vein, and again by bluish-white; inner portion of disc pale, with indistinct black marks at base of wing. UNDER-SIDE pale greyish-white, tinged with ochreous at the base and apex; with markings as above, but very indistinct; abdominal margin greenish-grey. Female, above as in male, but paler, the under-side being tinged with very pale purple on the exterior half. Wings shaped as in *Ad. Trigerta*; expanse of male $2\frac{1}{4}$, female, $2\frac{3}{4}$ inches.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

382. *ADOLIAS PULASARA*, Moore.

Adolias Pulasara, n. sp.—Allied to *Adol. Palguna*, but differs in the male on the upper-side, in the *fore-wing*, in having the transverse band tinged with hyaline-blue; there is also a cyaneous tinge about the extremity of the cell: the band on the *hind-wing* is rather broader, and composed of less lanceolate portions. UNDER-SIDE very pale ochreous, band of *fore-wing* tinged with pale hyaline-blue: *hind-wing* with two rows of small deep-black lanceolate spots (representing the band of the upper-side); base of wing with a number of deep-black marks. Female: UPPER-SIDE as in male, and the under-side with paler markings. Expanse of male $2\frac{5}{8}$, female, 3 inches.

a. ♀. Pinang. Presented by Dr. Cantor.

Remark.—The male of *Ad. Pulasara* is in the Collection of W. W. Saunders, Esq.

383. *ADOLIAS PUSEDIA*, Moore.

Adolias Pusedia, n. sp.—Female: UPPER-SIDE brown, with a pale purple tinge: *fore-wing* with a broad band of irregular-shaped purplish hyaline-like spots, each spot being concave on the outward margin; also a submarginal row of ill-defined whitish spots; base of wing with indistinct black marks: *hind-wing* with two curved rows of indistinct whitish spots, with some indistinct blue spots between the two rows. UNDER-SIDE ochreous, whitish on the exterior margins; band of *fore-wing* bluish-white, bound outwardly narrowly with brown; distinct black marks within discoidal-cell: *hind-wing* with two narrow curved indistinct ochreous-brown bands. Expanse 3 inches.

a. ♀. Pinang. Presented by Dr. Cantor.

384. *ADOLIAS TRIGERTA*, Moore.

Adolias Trigerta, n. sp.—UPPER-SIDE dark brown: male, *fore-wing* with a transverse row of six whitish spots, more or less suffused with brown, from subcostal vein one-third from the apex to near posterior margin near the angle, the first two spots being long and oval, third the smallest, fourth and fifth larger than the third, and both alike, sixth smaller and heart-shaped, all of them terminating outward in a point, which is bounded by black; base of wing with indistinct black markings; posterior margin at the angle with a small blue patch: *hind-wing* with a band of seven clear white spots, curving from anterior margin near the angle to abdominal margin near the anal angle, these spots having within each on the outward portion a black lanceolate mark pointing outward, these being slightly margined laterally with blue, and those nearest the anal angle with a short black line crossing the vein above the lanceolate mark; base of wing with indistinct black markings; abdominal margin pale brown. UNDER-SIDE greyish: *fore-wing* with the costal margin and basal half ochreous-yellow; the band of spots less distinct than above, but bounded inwardly with black lunulated marks, and outwardly with a black spot at the point of each; the markings at the base of the wing broadly black: *hind-wing* with a purple-red tinge on the anterior half; the band of spots less defined, bounded inwardly with black lunulated marks, and outwardly with triangular black spots, those near the anal angle with a narrow line above; base of wing with broad black marks within, above, and below the discoidal-cell; body and legs ochreous. Female above as in the

male, but with the markings larger; under-side as in male, but the markings less distinct. Expanse of male $2\frac{1}{2}$, female, $2\frac{3}{4}$ inches. Hind-wings of the male rounded, as in female.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

385. *ADOLIAS IAPIS*, Godart Sp.

Nymphalis Iapis, Godart, Enc. Méth. IX. p. 382 (1819).

Lucas, Hist. Nat. Lep. Exot. t. 69, f. 1.

Adolias Iapis, Boisduval. E. Doubleday, List Lep. Brit.

Mus. pt. I. p. 104. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 291, n. 8.

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

386. *ADOLIAS AMBALIKA*, Moore.

Adolias Ambalika, n. sp.—Female: UPPER-SIDE dark brown: fore-wing with transverse band of white spots, each spot with a long black point outward, between the veins; also an outer row of less-defined lanceolated whitish marks; black markings at base of the wing: hind-wing with transverse band of smaller spots as in fore-wing, also with an outer row of lanceolate marks. UNDER-SIDE pale ochreous, marked as above, with the black marks to the white band broader. Variety: UPPER-SIDE much darker brown: fore-wing with the transverse band suffused with brown; space between the band to near exterior margin bluish: hind-wing with less-defined band of smaller lanceolate marks, between which to near exterior margin bluish, centred with whitish lanceolate marks. UNDER-SIDE dull ochreous, with brown margins; band of lanceolate marks as above, but less suffused with brown. Expanse 3 inches.

a. ♀ var. Borneo. Presented by Hugh Low, Esq.

387. *ADOLIAS JAHNU*, Moore.

Adolias Jahnu, n. sp.—Female: UPPER-SIDE obscure brown, glossed with green: fore-wing with two indistinct black zigzag lines across the disc, tinged with white anteriorly; markings at base of wing large and indistinct: hind-wing with two indistinct black zigzag lines across the disc, also indistinct black basal marks. UNDER-SIDE ochreous brown, with transverse zigzag black lines and

basal marks as above; apex of fore-wing with a white spot and a blackish patch; exterior margin dusky; basal half of hind-wing, and along the outer zigzag line, with a bluish-grey tinge. Anterior wings falcate; expanse $3\frac{1}{4}$ inches.

a. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

388. ADOLIAS SIKANDI, Moore.

Adolias Sikandi, n. sp.—Female: UPPER-SIDE obscure glossy pale greenish-brown: fore-wing with the disc from middle of anterior to middle of posterior margin white, sharply defined inwardly from the basal half by an irregular zigzag division; also parallel ill-defined brown and white spots; two large black marks within discoidal cell: hind-wing with narrow curved white band, tapering from middle of anterior to near middle of abdominal margin, and sharply defined inwardly from the basal half; a submarginal row of deep-brown lunular spots; blackish marks within discoidal-cell. UNDER-SIDE ashy-grey, tinged with blue, basal half dark, exterior half very pale; markings as above; on hind-wing a small red spot in centre of anterior basal mark. Shape of wings as in *Ad. Evelina*; expanse $3\frac{1}{2}$ inches.

a. ♀. Java. From Dr. Horsfield's Collection.

389. ADOLIAS EVELINA, Stoll Sp.

♂ *Papilio Evelina*, Stoll, *Suppl. Cramer's Pap. Exot.* t. 28, f. 2, 2B (1791).

Adolias Evelina, Boisduval. E' *Doubleday, List Lep. Brit. Mus.* pt. I. p. 104. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 7.

Nymphalis Evelina, Godart, *Enc. Méth.* IX. p. 401.

Adolias Derma, Kollar, *in Hügel's Kaschmir*, IV. pt. II. p. 436 (1844). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 292, n. 23.

a. b. c. d. ♂ ♀. Assam. From Griffith's Collection.

390. ADOLIAS COCYTINA, Horsfield Sp.

♂ *Aconthea Cocytina*, Horsfield, *Zool. Journ.* V. p. 67, t. 4, f. 3, 3a (1830).

Adolias Cocytina, Westwood, *in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 3.

- δ Papilio Cocytta, *Fabricius*,* *Ent. Syst.* III. pt. I. p. 127 (1793). *Jones, Icon.* IV. t. 64, f. 2.
Nymphalis Cocytta, *Godart, Enc. Méth.* IX. p. 382.
Adolias Cocytta, *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 104.
Adolias Godartii, *G. R. Gray, Cat. Lep. Ins. Nepal*, p. 14, t. 12, f. 2 (1833).
- a. δ .* Borneo. Presented by Hugh Low, Esq.
b. c. d. δ . Borneo. Presented by Edwin Shepherd, Esq., 1857.

391. ADOLIAS COCYTUS, *Fabricius Sp.*

- Papilio Cocytus, *Fabricius, Mant. Ins.* II. p. 29 (1787); *Ent. Syst.* III. pt. I. p. 55.
Adolias Cocytus, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 9.
Nymphalis Cocytus, *Godart, Enc. Méth.* IX. p. 368.
Adolias Sidera, *Boisduval, MS.*
- a. b. c. δ ♀.* N. India. Presented by Colonel Sykes.
d. e. δ . Canara, S. India. Presented by S. N. Ward, Esq.

392. ADOLIAS ADIMA, *Moore.*

Adolias Adima, n. sp.—Male: UPPER-SIDE yellowish olive-brown, with two indistinct zigzag lines crossing the disc of both wings, and markings about the base of the wings blackish. UNDER-SIDE pale ferruginous, palest at the base of fore-wings, with inner row of lunulated marks and outer row of spots crossing the disc of both wings, and markings at the base of the wings, black, the spots being deepest from anal angle, where they are margined with bluish-white; abdominal margin broadly, and about anal angle somewhat, green. Expanse $2\frac{1}{2}$ inches.

- a. δ .* Assam. From Griffith's Collection.

The male of *Adolias Adima* may at once be distinguished from *Ad. Apiades* by its plain brown upper-side.

* But not *Pap. Cocytus*, *Fabricius*.

393. *ADOLIAS APIADES*, Menetries.

♂ *Adolias Apiades*, *Menetries*, *Cat. Lep. Mus. Imp. Acad. of Sci. St. Petersburg*, pt. II. t. IX. f. 4 (ined.).

a. b. c. d. e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

394. *ADOLIAS SANCARA*, Moore.

Adolias Sancara, n. sp.—Male: UPPER-SIDE brown: fore-wing with an outward oblique white band from middle of costal margin to near posterior margin above the angle; also two small white spots near the apex; an indistinct blackish transverse band from near apex of fore-wing to abdominal margin; also an indistinct narrow submarginal blackish band on the hind-wing; black marks within the discoidal-cell. UNDER-SIDE greyish-brown, greyer at the base and along exterior margins; marked as above. Female darker, and marked as in male. Expanse of male $3\frac{1}{4}$, female, $3\frac{1}{2}$ inches.

a. ♂. N. India. Presented by Colonel Sykes.

b. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

395. *ADOLIAS IVA*, Moore.

Adolias Iva, n. sp.—Male: UPPER-SIDE very dark olive-green: fore-wing with oblique row of large, long greenish-white spots from middle of costal margin to near posterior angle, beneath which is a small narrow geminated spot near the middle of the posterior margin; also two small spots obliquely near the apex, and a small whitish patch at the posterior angle: hind-wing with a row of broadly-separated small round greenish-white spots curving from middle of costal margin to middle of the wing; also black marks within discoidal-cell. UNDER-SIDE deep greenish-grey, dusky about the anterior half, and blackish along the posterior margin; markings as above, but the spots on the hind-wing extending nearly to abdominal margin. Shape of wings as in *Adolias Epiona*; expanse 4 inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Adolias Iva may be known from *Adolias Epiona* by its very dark upper-side, and by the oblique band being composed of much longer spots, and by its deep greyish-green under-side.

396. *ADOLIAS EPIONA*, G. R. Gray Sp.

Aconthea Epiona, G. R. Gray, *List Lep. Nepal*, p. 13 (1833).

Adolias Doubledayii, *Boisduval*, MS. E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 104 (1844). G. R. Gray, *List Lep. Nepal*, p. 13, t. 13. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 15.

Adolias Patala, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 435 (1844).

a. b. c. d. e. ♂ ♀. N. India. Presented by Colonel Buckley.

397. *ADOLIAS DURGA*, Moore.

Adolias Durga, n. sp.—UPPER-SIDE dark olive-green, with a black marginal and submarginal band; a broad transverse band of irregular-shaped white spots crossing from middle of costal margin of fore-wing to beyond the middle of the hind-wing, near the anal angle, being margined exteriorly with blue from the fore-wing on its lower half to anal angle; the narrow space between marginal and submarginal bands also bluish; two small rounded white spots near the apex of fore-wing; marks within discoidal-cells black. UNDER-SIDE from inner margin of band to base greenish-grey, from its outer margin to extremity of wing greenish, with band and apical spots as above; a row of blackish marks from apical spots to posterior angle, which is whitish; also an indistinct dusky submarginal row on the hind-wing, and at the anal angle two patches of black; marks at the base of wings black; ciliae between the indentations on the upper and under-side white. Sexes alike. Expanse 4 inches.

a. b. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

398. *ADOLIAS TEUTA*, E. Doubleday.

♂ *Adolias Teuta*, E. Doubleday, MS. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 5, t. 44, f. 2 (1850).

a. b. c. d. ♂ ♀. Silhet. Presented by Colonel Buckley.

e. ♂. Java. From Dr. Horsfield's Collection.

399. *ADOLIAS FRANCIAE*, G. R. Gray.

Adolias Franciae, G. R. Gray, *List Lep. of Nepal*, p. 12, t. 14 (1833). *E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 104. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 14.

- a. b. ♂. Darjeeling. From Pearson's Collection.
- c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- d. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

400. *ADOLIAS CORESIA*, Hübner Sp.

♀ *Hypolimnas Coresia*, Hübner, *Sammel. Exot. Schmett. Band II.* t. (1806-27).

Adolias Coresia, E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 105.

♂ *Aconthea Apaturina*, Horsfield, *Zool. Journ. V.* p. 68, t. 4, f. 1, 1a (1830).

Adolias Apaturina, Westwood, *in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 4.

a. b. ♂ (*A. Apaturina*, Horsf.) Java. Presented by Dr. Horsfield.

c. ♂. Java. From Mr. Argent's Collection.

401. *ADOLIAS NICEA*, G. R. Gray.

Aconthea Nicea, G. R. Gray, *Lep. Ins. of Nepal*, p. 13, t. 12, f. 1 (1833).

Adolias Nicea, E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 105. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 291, n. 2.

a. b. ♂ ♀. N. India. Presented by Colonel Sykes.

c. ♀. N. India. From T. Tatum, Esq.

d. e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

402. *ADOLIAS NESIMACHUS*, Boisduval.

Adolias Nesimachus, Boisduval, *in Cuvier's Règ. Anim. edit. Crochard, Ins. t. 139, f. 1 (). E. Doubleday, List Lep. Brit. Mus. pt. I.* p. 105. *Westwood,*

in Doubleday and Hewitson's Diurnal Lep. p. 291,
n. 12.

Argynnис Hippomenes, Kaden, in Herr Schaeffer's Lep.
Exot. ser. II. fasc. 1 et 2, fig. 11, 12 (1853).

a. b. ♂. Assam. From Griffith's Collection.

c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

403. *ADOLIAS DIRTEA*, Fabricius Sp.

♀ Papilio Dirtea, Fabricius, Ent. Syst. III. pt. I. p. 59
(1793). *Jones, Icon. IV. t. 65, f. 1.*

Adolias Dirtea, E. Doubleday, List Lep. Brit. Mus.
pt. I. p. 104. G. R. Gray, List Lep. Nepal, p. 12,
t. 10, f. 1, ♀, 2, ♂. Westwood, in Doubleday and
Hewitson's Diurnal Lep. p. 291, n. 17, t. 44, f. 1, ♀.

Aconthea Boisduvalii, G. R. Gray, MS. Boisduval,
Spéc. Gén. Lép. I. t. 8, f. 2.

a. ♂. Assam. From Griffith's Collection.

b. ♂. N. India. From T. Tatum, Esq.

c. ♀. N. India. Presented by the Trustees of the
British Museum.

d. ♀. Pinang. Presented by Dr. Cantor.

e. f. ♂ ♀. Borneo. Presented by Hugh Low, Esq.

g. ♀. Borneo. Presented by Edwin Shepherd, Esq.

404. *ADOLIAS SIVA*, Westwood.

♀ Aconthea Doubledayi, Westwood, Cabinet Oriental Ent.
p. 76, t. 37, f. 4 (1847).

Adolias Siva, Westwood, in Doubleday and Hewitson's
Diurnal Lep. p. 291, n. 18 (1850).

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collec-
tion.

Genus EURIPUS, E. Doubleday.

EURIPUS, E. Doubleday, MS. Westwood, in Doubleday and Hewit-
son's Diurnal Lep. p. 293 (1850).

405. *EURIPUS HALITHERSES*, E. Doubleday.

Euripus Halitherses, E. Doubleday, MS. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 293, t. 51, f. 2 (1850).

- a. ♂. Darjeeling. From Pearson's Collection.
- b. c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- d. e. f. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

406. *EURIPUS HALLIROTHIUS*, Westwood.

Euripus Hallirothius, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 293 (1850).

- a. ♂. N. India. Presented by Colonel Buckley.
- b. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—The females of both these species are at present unknown.

Genus HERONA, E. Doubleday.

HERONA, E. Doubleday, MS. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 293 (1850).

407. *HERONA MARATHUS*, E. Doubleday.

Herona Marathus, E. Doubleday, MS. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 294, t. 51, f. 3 (1850).

- a. ♂. Assam. Presented by Colonel Hearsey.
- b. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus CASTALIA, Boisduval.

CASTALIA, Boisduval, MS.

408. *CASTALIA DICHROA*, Kollar Sp.

Limenitis dichroa, Kollar, in Hügel's *Kaschmir*, IV. pt. II. p. 429, t. 8, f. 1, 2 (1844).

Apatura dichroa, Westwood, in Doubleday and Hewitson's
Diurnal Lep. p. 305, n. 21.

Castalia Nyctis, Boisduval, MS.

a. ♂. Simla. Presented by the Trustees of the British Museum.

b. ♂. N. India. Presented by Colonel Buckley.

Castalia dichroa may be distinguished from *Castalia Chandra* by the exterior margin of the fore-wings being much less deflexed in the middle, and by having all the markings of the upper-side of the fore-wing orange-colour, and being without the black spots in the discoidal-cell of the hind-wing, and on the under-side by the pearly-white on the hind-wings. Expanse of wings $2\frac{3}{4}$ inches.

409. *CASTALIA CHANDRA*, Moore (Plate VIa, fig. 4).

Castalia Chandra, n. sp.—UPPER-SIDE: *fore-wing* jet-black, with a greenish gloss at the apex; a transverse curved row of four orange-coloured spots from costal margin one-fourth of the wing to middle of posterior margin; a straight row of five white spots from middle of costal margin to near posterior angle; also two small and one minute white spot near the apex; a marginal and submarginal row of spots, those at the apex whitish, the marginal ill-defined, the submarginal rather large and clearly defined, and those to the posterior angle small and bluish: *hind-wing* clear orange-colour, with rather broad black posterior margin, centred with a marginal row of small whitish spots; also a parallel inner zigzag black band from anal angle to anterior angle, and extending along the costal margin, the black extending along the veins upwards and downwards to marginal band; a small black spot nearly at the base of and within discoidal-cell, and another in the centre of the wing; on the thickest part of the inner band near anal angle is a small central bluish-white spot; abdominal margin broad dusky-white; body black; frontal tuft and neck spotted with white. UNDER-SIDE as above, but on the *fore-wing* the colours are brighter; the costal margin greenish-white, and there is also a whitish triangular spot at base of discoidal-cell: on the *hind-wing* along the marginal band is a row of bluish-white lunated spots; the spot on the thickest part of the inner band is large, as is also the central black spot within the discoidal-cell; at the base of the wing are a number of greenish-white spots; all the veins are broadly lined with black; body and abdomen on the sides spotted with white. Expanse of wings 3 inches.

a. b. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

c. d. e. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus nov. DILIPA, Moore.

This genus differs from the typical *Apatura* in having the discoidal cell closed in all the wings by a very delicate lower disco-cellular vein; the palpi and antennæ are, however, like those of *Apatura*.

410. *DILIPA MORGIANA*, Westwood Sp. (Plate VIa, fig. 5, ♂ ♀).

♂ *Apatura Morgiana*, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 305, n. 22 (1850).

a. ♂ (type). N. India. Presented by Colonel Buckley.

The female of *Dilipa Morgiana* is in the British Museum Collection, and differs from the male in being somewhat larger, and having all the spots of the fore-wing white, the broad band across the hind-wing being also white. Under-side as in male, but the spots white.

Genus APATURA, Fabricius.

APATURA, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 280, 1807).
Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 302.

CATARGYRIA, Doxocopa, Hübner, Verz. bek. Schmett. (1816).

CHLORIPPE, Boisduval, MS. E. *Doubleday, List Lep. Brit. Mus.* pt. I. p. 108 (1844).

NYMPHALIS, Blanchard.

411. *APATURA AMBICA*, Kollar.

♂ *Apatura Ambica*, Kollar, in Hügel's Kaschmir, IV. pt. II. p. 431, t. 8, f. 3, 4 (1844).

♂ *Apatura Namouna*, E. *Doubleday, Ann. Nat. Hist.* XVI. p. 178 (1845); *List Lep. Brit. Mus.* pt. I. App. p. 28. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 305, n. 18.

- a. b. ♂.* N. India. Presented by Colonel Buckley.
- c. ♂.* Darjeeling. From Pearson's Collection.
- d. ♂.* N. India. Presented by Lieutenant Hugo James.
- e. ♂.* Darjeeling. From Messrs. Schlagintweit's Collection.

A female of *Apatur. Ambica* is in the Collection at the British Museum, and is distinguished from the male in being paler, and having none of that beautiful gloss on the upper-side; the spots are yellowish-white, and those along the exterior margins are larger.

412. *APATURA PARVATA*, Moore (Plate VIa, fig. 6).

Apatura Parvata, n. sp.—Male: UPPER-SIDE dark ferruginous-brown, darkest at the base, and bright about discoidal cell and along the exterior margin: *fore-wing* with blackish marks in discoidal cell; marginal and submarginal lines, and indistinct narrow interrupted transverse series of pale ferruginous spots from middle of anterior to posterior margin; also a series of five small white spots from anterior margin near the apex, below which is an indistinct black spot: *hind-wing* with narrow pale ferruginous band from middle of anterior margin to near middle of abdominal margin; also, near anal angle, a black spot encircled with ferruginous; abdominal margin dull ferruginous-brown. UNDER-SIDE ferruginous, brightest on the fore-wings; markings disposed as above: *fore-wing* with the transverse series of spots distinct and white, being bounded inwardly with blackish patches; near the apex and posterior angle are some patches of black; the black spot on disc large; also a black patch on posterior base of the wing; marginal line ashy: *hind-wing* ferruginous-brown across the disc, with band as above white; on the lower part of the disc is a small black spot centred with blue and encircled with ferruginous; a marginal line and submarginal zigzag marks ashy. Expanse of wings 2 to $2\frac{1}{8}$ inches.

- a. b. ♂.* N. India. Purchased.

413. *APATURA PARISATIS*, Westwood.

Apatura Parisatis, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 305, n. 20 (1850).

- a. ♂.* Assam. From Griffith's Collection.
- b. ♀.* Canara, S. India. Presented by S. N. Ward, Esq.

c. d. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

g. ♂. Java. From Mr. Argent's Collection.

Apatura Parisatis may be distinguished by its small size, and by the uniform jet-black colour of the upper-side of the wings; its under-side being varied with rich red-brown and chestnut, and with a slight whitish macular fascia across the middle of the wings, and a minute eye-spot near the anal angle. The female is dull-ferruginous; both upper and under-side with markings nearly similar to those of the under-side of the male, and with a curved row of small white dots near the apex of the fore-wings. Expanse of wings $1\frac{3}{4}$ to nearly 2 inches.

414. APATURA NAKULA, Moore.

Apatura Nakula, n. sp.—Female: UPPER-SIDE dull pale reddish-brown, darkest at the base: *fore-wing* with a broad transverse whitish band from anterior to posterior margin; two blackish reniform marks and a dot within discoidal cell; a row of five small white dots from anterior margin near the apex, the three first being encircled with black, and a small sixth spot wholly black; a narrow marginal line blackish, and submarginal line dark brown: *hind-wing* with a broad nearly straight white band from anterior margin to near anal angle, where it is narrowest; an oval and two small black marks within discoidal cell; a narrow dark black marginal line; a submarginal row of dark-brown marks, bounded inwardly by a row of zigzag white marks; near the anal angle, between the middle and lower median veins, is a small black spot encircled with pale red; abdominal margin pale. UNDER-SIDE very pale ochreous-brown, with markings as above, but less distinct; the black spot on the fore-wing is rather large, but on the hind-wing this spot is very small. Expanse of wings $1\frac{8}{10}$ inches.

a. ♀. Java. From Dr. Horsfield's Collection.

Genus SYMPHÆDRA, Hübner.

SYMPHÆDRA, Hübner, Verz. bek. Schmett. (1816).

SYMPHEDRA, E. Doubleday.

LEXIAS, Boisduval, Voy. de l'Istrolube, Ins. pt. I. p. 125 (1832).

415. *SYMPHÆDRA THYELIA*, Fabricius Sp.

Papilio Thyelia, *Fabricius*, *Ent. Syst.* III. pt. I. p. 142 (1793). *Jones, Icon.* V. t. 83, f. 2.

Sympædra Thyelia, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 295, t. 42, f. 6.

Argynnus Thyelia, *Godart, Enc. Méth.* IX. p. 257. *Donovan, Ins. of India*, t. 31, f. 3.

Sympædra Thyelia, *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 105.

Sympædra Alcandra, *Hübner, Zuträge*, pt. I. p. 7, f. 1, 2, et *Verz. bek. Schmett.* n. 346.

a. b. c. ♂ ♀. N. India. From the Asiatic Society of Calcutta.

d. ♂. S. India. Presented by S. N. Ward, Esq.

Genus NYMPHALIS.

NYMPHALIS, *Latreille, Hist. Nat. des Ins.* XIV. p. (1805).

Godart, Enc. Méth. IX. p. 329. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 306 (1850).

CHARAXES, *Ochsenheimer, Schmett. von Eur.* IV. (1816). *Boisduval. E. Doubleday.*

IASIA, *Swainson, Zool. Ill. 2nd ser.* t. 90 (1831-2).

PAPHIA, sect. I., *Fabricius, Syst. Gloss. (Illiger's Mag.* VI. p. 282, 1807). *Horsfield (part).*

ERIBOEA et TIGRIDIA, *pt. Hübner, Verz. bek. Schmett.* (1816).

416. *NYMPHALIS FABIUS*, Fabricius Sp.

Papilio Fabius, *Fabricius, Spec. Ins.* II. p. 12 (1781); *Ent. Syst.* III. pt. I. p. 64.

Nymphalis Fabius, *Godart, Enc. Méth.* IX. p. 353. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 28.

Papilio Euphanes, *Esper, Ausl. Schmett.* I. p. 238, t. 59, f. 1 (1785).

Charaxes Fabius, *Boisduval. E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 110.

a. b. c. ♂ ♀. N. India. Presented by Colonel Buckley.

d. ♂. N. India. Presented by Colonel Hearsey.

Colonel Hearsey informs us that "he usually captured this species upon wounded apple-trees; the butterflies frequenting the wounds to suck up the fluids which were discharged therefrom."—(Westwood's Oriental Ent.)

Captain Mortimer J. Slater, in his MS. Notes, p. 41, states: "At Dinafore, I generally found *Charaxes Fabius* near, or on, pomegranate-trees. Flies rapidly. Uncommon."

417. NYMPHALIS ATHAMAS, Drury Sp.

Papilio Athamas, Drury, Ill. I. t. 2, f. 3, 4 (1770).

Cramer, Pap. Exot. I. t. 89, f. C.D.

Nymphalis Athamas, Godart, Enc. Méth. IX. p. 353.

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 309, n. 29.

Iasia Athama, Swainson, Zool. Ill. 2nd ser. t. 90.

Paphia Athamas, Horsfield, Cat. Lep. Mus. E.I.C. t. 8, f. 7, 7a.

Charaxes Athamas, Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 109.

Papilio Pyrrhus, Donovan, Ins. of India, t. 29, f. 3 (1800-3).

Eriboea Athamis, Hübner, Verz. bek. Schmett. n. 430 (1816).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. N. India. Presented by Colonel Sykes.

g. h. ♂ ♀. N. India. Presented by Colonel Buckley.

The larva and pupa of *Nymphalis Athamas* are figured on Plate VI., fig. 3, 3a, from Java, discovered by Dr. Horsfield.

418. NYMPHALIS SCHREIBERI, Godart.

Nymphalis Schreiberi, Godart, Enc. Méth. IX. App. p. 825 (1819). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 309, n. 30.

Paphia Schreibers, Horsfield, Cat. Lep. Mus. E.I.C. t. 6, f. 3, 3a.

Charaxes Schreiberi, Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 110.

a. ♀. Java. From Dr. Horsfield's Collection.

419. *NYMPHALIS DELPHIS*, E. Doubleday Sp.

Charaxes Delphis, E. Doubleday, *Annals Soc. Ent. France* (1843), p. 217, t. 7.

Nymphalis Delphis, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 33.

a. ♂. Silhet. Presented by the Trustees of the British Museum.

420. *NYMPHALIS EUDAMIPPOS*, E. Doubleday Sp.

Charaxes Eudamippus, E. Doubleday, *Annals Soc. Ent. France* (1843), p. 218, t. 8.

Nymphalis Eudamippus, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 32.

a. b. c. ♂ ♀. Darjeeling. From Pearson's Collection.

d. e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

421. *NYMPHALIS DOLON*, Westwood.

Charaxes Dolon, Westwood, *Cabinet Orient. Ent.* p. 55, t. 27, f. 2, 3 (1847).

Nymphalis Dolon, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 34.

a. b. c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

422. *NYMPHALIS MARMAX*, Westwood.

♂ ♀ Charaxes Marmax, Westwood, *Cabinet Orient. Ent.* p. 43, t. 21, f. 3, 4, 5 (1847).

Nymphalis Marmax, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 87.

a. ♂. Darjeeling. From Pearson's Collection.

b. ♂. Cherra Poonjee. From Colonel Buckley's Collection.

c. d. e. f. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

423. *NYMPHALIS BERNARDUS*, Fabricius Sp.

Papilio Bernardus, Fabricius, *Ent. Syst. III. pt. I.* p. 71 (1793). *Donovan, Ins. of China*, t. 34.

- Papilio Bernarda, *Jones, Icones*, IV. t. 65, f. 2 ().
Nymphalis Bernardus, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 309, n. 35.
Satyrus Bernardus, *Godart, Enc. Méth.* IX. p. 477.
Charaxes Bernardus, *E. Doubleday, List Lep. Brit. Mus.*
 pt. I. p. 112. *Kollar, in Hügel's Kaschmir*, IV.
 pt. II. p. 434, t. 11, f. 1, 2.
Papilio Polyxena, *Cramer, Pap. Exot.* I. t. 54, f. A.B.
 (1779).
Nymphalis Polyxo, *Godart, Enc. Méth.* IX. p. 399
 (1819).
Doxocopa Epilais, *Hübner, Verz. bek. Schmett.* p. 464
 (1816).
a. b. c. ♂♀. Silhet. From Colonel Buckley's Collection.
d. e. f. ♂. Darjeeling. From Indian Collection,
 Exposition Universelle at Paris, 1855.

Captain Mortimer J. Slater, in his MS. Notes, p. 144, informs us that " *Charaxes Bernardus* flies along about four feet from the ground, sometimes rising to ten feet or so; and in this manner traverses the open ground at a fast flight, seldom settling or pausing; and is consequently very difficult to capture, much less so in perfect condition."

424. NYMPHALIS BAYA, Moore.

Nymphalis Baya, n. sp.—Allied to *Nymph. Psaphon*, Westwood, Cabinet Orient. Ent. t. 21, f. 1, 2, from Ceylon, but differs in the male in having the black exterior border of the *fore-wing* much narrower, it not extending to the middle of the posterior margin, as in that species, but is confined to the angle: on the *hind-wing*, the apical patch is also smaller, and continued in a series of six white centred spots to anal angle; a small blackish spot at extremity of the discoidal-cell. UNDER-SIDE glossy greenish dusky-brown. Female paler above: *fore-wing* with a broad central yellowish band; a small round yellowish spot near the apex; a patch of black at extremity of discoidal-cell, and narrow zigzag line bordering the black band: *hind-wing* with the central upper portion of disc yellowish; submarginal row of spots clearly defined, and all centred with clear white. Under-side more variegated than the male, and much paler.

Tails two, short in the male; *both* long in the female. Expanse of male $2\frac{3}{4}$ inches, female rather more than 3 inches.

a. b. c. ♂ ♀ (type). Java. From Dr. Horsfield's Collection.

d. e. ♂. N. India. From Colonel Buckley's Collection.

The larva, pupa, and imago of *Nymph. Baya* are figured among the original drawings made by Mrs. Hamilton in Northern India, which are now in the possession of the Entomological Society of London. A copy of these will be drawn on a subsequent Plate of this Catalogue. Mrs. Hamilton's Notes state that the "larva was found on the 9th November, it changing to a pupa on the 20th, and the imago emerging on the 30th."

Genus KALLIMA, *E. Doubleday.*

KALLIMA, *E. Doubleday, MS. Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 324 (1850).

PAPHIA (*part*), *Horsfield. Hügel.*

AMATHUSIA (*part*), *Boisduval. Westwood.*

ZARETIS (*part*), *Hübner.*

425. KALLIMA PARALEKTA, *Horsfield Sp.*

♂ Paphia Paralekta, *Horsfield, Cat. Lep. Mus. E.I.C. t. 6, f. 4* (1829).

♀ Kallima Paralekta, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 325, n. 1, t. 52, f. 3.

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

Kallima Paralekta.—Female: fore-wings hooked at the tip; the broad oblique fascia white; base of fore-wing and whole of hind-wing ferruginous-brown.

426. KALLIMA INACHIS, *Boisduval Sp.*

Amathusia Inachis, *Boisduval, in Crochard's edit. of Cuv. Règ. Anim. Ins. t. 139, f. 3* (). Herrick-Schaeffer, *Lep. Exot. II. f. 7, 8.*

Kallima Inachis, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 325, n. 2.

Paphia Hügelii, *Kollar, in Hügel's Kaschmir, IV. pt. II. p. 432, t. 9* (1844).

Paphia Paralekta, *apud Royle*, *Ill. of Himal. Bot.* t. 10,
f. 3a, 3b.

Amathusia Paralekta, *apud E. Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 114.

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.

c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

e. f. g. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

427. *KALLIMA BISALTIDE*, Cramer Sp.

Papilio Bisaltide, *Cramer, Pap. Exot.* II. t. 102, f. C.D. *Suriwan* (1779). *Fabricius, Ent. Syst.* III. pt. I. p. 74.

Kallima Bisaltide, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 325, n. 8.

Apatura Bisaltide, *Hübner, Verz. bek. Schmett.* n. 294.

Nymphalis Bisaltide, *Godart, Enc. Méth.* IX. p. 370.

Amathusia Bisaltide, *E. Doubleday, List Lep. Brit. Mus.* pt. I. p. 114.

Papilio Polibetes, *Cramer, Pap. Exot.* III. t. 234, f. D.E. *Ambryna* (1782).

Zaretis Polibetes, *Hübner, Verz. bek. Schmett.* n. 453.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Pinang. Presented by Dr. Cantor.

f. g. ♂ ♀. N. India. Presented by Colonel Buckley.

h. i. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus AMATHUSIA, Fabricius.

AMATHUSIA, *Fabricius, Syst. Gloss.* (*Illiger's Mag.* VI. p. 279, 1807). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 326.

MERA (part), *Hübner, Verz. bek. Schmett.* (1816).

THORESSA, *Boisduval, MS.*

428. *AMATHUSIA PHIDIIPPUS*, Linnaeus Sp.

Papilio Phidippus, *Linnæus, Syst. Nat.* II. p. 752 (1767).

Fabricius, Ent. Syst. III. pt. I. p. 71. *Cramer, Pap. Exot.* I. t. 69, f. A.B.

Amathusia Phidippus, *Fabricius*, *Syst. Gloss.* (*Illiger's Mag.* VI. p. 279, 1807). *Horsfield*, *Cat. Lep. Mus. E. I. C. t. 7, f. 10.* *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 114.* *Westwood*, *in Doubleday and Hewitson's Diurnal Lep.* p. 327, n. 1, t. 54a, f. 2.

Morpho Phidippus, *Godart*, *Enc. Méth. IX.* p. 439.

Thoræssa Phidippus, *Boisduval*, *MS.*

Mœra Phidippe, *Hübner*, *Verz. bek. Schmett.* n. 477.

Var. *Amathusia Patalena*, *Westwood*, *Cab. Orient. Ent.* p. 40, t. 19, f. 1 (1847).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Pinang. Presented by Dr. Cantor.

The larva and pupa of *Amathusia Phidippus* are figured on Plate VI., fig. 4, 4a, from Java, where it "feeds on the young leaves of *Cocos nucifera*, from December to April. Common."—(Horsfield, MS.)

429. *AMATHUSIA AMYTHAON*, *E. Doubleday*.

Amathusia Amythaon, *E. Doubleday*, *Ann. Nat. Hist. XIX.* p. 175 (1847); *List Lep. Brit. Mus. App. p. 29.* *Westwood*, *Cab. of Orient. Ent.* p. 29, t. 19, f. 1, 2, 3; id. *in Doubleday and Hewitson's Diurnal Lep.* p. 327, n. 2.

Thoræssa Thessalus, *Boisduval*, *MS.*

a. ♀. N. India. Donor unknown.

Genus *ZEUXIDIA*, *Hübner*.

ZEUXIDIA, *Hübner*, *Samml. Exot. Schmett. II.* (1806–27). *Westwood*, *in Doubleday and Hewitson's Diurnal Lep.* p. 327.

AGLAURA, *Boisduval*, *MS.* *Herrich-Schäffer*.

430. *ZEUXIDIA LUXERI*, *Hübner*.

Zeuxidia Luxerii, *Hübner*, *Samml. Exot. Schmett. Band II. t.* (1806–27). *E. Doubleday*, *List Lep. Brit. Mus. pt. I. p. 114.* *Westwood*, *in Doubleday and Hewitson's Diurnal Lep.* p. 329.

a. ♂. Java. From Dr. Horsfield's Collection.

Genus DISCOPHORA, Boisduval.

DISCOPHORA, Boisduval, Spéc. Gén. Lép. t. 12, f. 3 (1836). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 329.

MCERA, ZERYNTHIA (*part*), Hübner.

MORPHO (*part*), Horsfield.

431. DISCOPHORA TULLIA, Cramer Sp.

♀ Papilio Tullia, Cramer, Pap. Exot. I. t. 81, f. A.B. (1779). Fabricius, Ent. Syst. III. pt. I. p. 98.

Discophora Tullia, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 114. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 331, n. 1.

Morpho Tullia, Godart, Enc. Méth. IX. p. 446.

Mcera Tullia, Hübner, Verz. bek. Schmett. n. 479.

a. b. c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

e. ♀. N. India. Presented by Colonel Sykes.

f. g. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

The transformations of *Discophora Tullia* are figured among the original drawings of A. Grote, Esq., of Calcutta, and are copied on Plate XII., fig. 15, 15a. It "feeds on *Bambusa*."

432. DISCOPHORA CELINDE, Stoll Sp.

Papilio Celinde, Stoll, Suppl. Cramer's Pap. Exot. V. t. 37, f. 1, 1a (1791).

Discophora Celinde, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 114. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 331, n. 2. Herr. Schaeffer, Lep. Exot. II. pt. 1 and 2, f. 5, 6 (1853).

Morpho Celinde, Godart, Enc. Méth. IX. p. 446. Horsfield, Cat. Lep. Mus. E.I.C. t. 6, f. 6, t. 7, f. 11.

Papilio Menetho, Fabricius, Ent. Syst. III. pt. I. p. 83 (1793). Jones, Icon. V. t. 61. Donovan, Ins. of India, t. 30, f. 1, ♀.

Morpho Menetho, Godart, Enc. Méth. IX. p. 446.

Var. ♀ ? Discophora Timora, E. Doubleday, Westwood and Hewitson's Diurnal Lep. t. 54, f. 2.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

The larva and pupa of *Discophora Celinde* are figured on Plate VI., fig. 5, 5a, from Java, where "it feeds on the young leaves of *Cocos nucifera*, from December to January. Common."—(Horsfield, MS.)

Genus ENISPE, *E. Doubleday*.

ENISPE, *E. Doubleday*, MS. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 292 (1850).

433. ENISPE EUTHYMIUS, *E. Doubleday*.

Adolias Euthymius, *E. Doubleday*, *Ann. Nat. Hist.* XVI. p. 179 (1845); *List Lep. Brit. Mus.* pt. I. App. p. 27.

Enispe Euthymius, *E. Doubleday*. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 292, t. 40, f. 2.

a. b. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

434. ENISPE CYCNUS, *Westwood*.

♂ Enispe Cyenus, *Westwood*, in Doubleday and Hewitson's *Diurnal Lep.* p. 330 (1851).

a. ♂. Darjeeling. From Pearson's Collection.

b. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Enispe Cyenus.—Male: UPPER-SIDE blackish-brown; base and exterior margins paler: fore-wing with an oblique band from middle of anterior margin terminating in a transverse series of lunular spots, and a parallel outer row of four quadrate spots, lilac-white: hind-wing with very indistinct marginal and submarginal pale spots. UNDER-SIDE with a transverse brown dentate band from middle of costal margin to anal angle; the base of the wings orange-yellow, with narrow irregular transverse markings; exterior half of wings light brown, with indistinct transverse lines: hind-wing with a small anterior black dot and a posterior white dot. Female: UPPER-SIDE, fore-wing with the base obliquely dull chocolate-brown, the rest brownish-black; the lilac-white oblique band and spots as in male,

but larger, with two inner rows and terminal spots ochreous-yellow : *hind-wing* with the basal half to abdominal angle dull chocolate-brown, with a marginal, submarginal, an inner third and a short fourth row of ochreous-yellow spots, the spaces between blackish. *UNDER-SIDE* yellow throughout, with a slight greenish tinge ; transverse dentate band and markings as in male. *Expanse* of male $3\frac{1}{8}$, female $3\frac{5}{8}$ inches.

435. *ENISPE LEPIDA*, Moore.

Enispe Lepida, n. sp.—Female: *UPPER-SIDE* ochreous-brown, more dusky about the anterior half: *fore-wing* with broad oblique transverse band from middle of anterior margin, and inner row of spots, lilac-white; an outer row of submarginal and indistinct marginal spots ochreous: *hind-wing* with exterior margin and three inner rows of spots ochreous. *UNDER-SIDE* pale dull dirty ochreous, with ill-defined transverse dentate line and markings about the base brown; indistinct rows of paler spots; *hind-wing* with an anterior brown spot centred with a white dot, and half margined with black; a posterior small white spot encircled with brown. *Expanse* 4 inches.

a. ♀. Canara, S. India. Presented by S. N. Ward, Esq.

Genus CLEROME, Boisduval.

CLEROME, *Boisduval, MS.* *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 332 (1851).

FAUNIS (*part*), Hübner.

436. *CLEROME ARCESILAUS*, Fabricius Sp.

Papilio Arcesilaus, *Fabricius, Ent. Syst.* III. pt. I. p. 153 (1793). *Donovan, Ins. of India*, t. 30, f. 2.

Clerome Arcesilaus, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 334, t. 54, f. 5.

Satyrus Arcesilaus, *Godart, Enc. Méth.* IX. p. 497.

Faunis canens, *Hübner, Samml. Exot. Schmett. Band II. tab.* (1806-27).

Morpho Leonteus, *Zinken-Sommer, Nova Acta Acad. Nat. Cur.* XV. p. 170, t. 16, f. 14, 15.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

437. CLEROME PHAON, Erichson Sp.

Drusilla Phaon, Erichson, in *Nova Acta Acad. Nat. Cur. XVI. Suppl.* p. 277, t. 50, f. 1, 2a (1834).

Clerome Phaon, Westwood, *Trans. Ent. Soc. IV. n. s. p.*

Thaumantis Phaon, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 337, n. 4.

a. b. ♂ ♀. Borneo. Presented by Hugh Low, Esq.

Genus DRUSILLA, Swainson.

DRUSILLA, Swainson, *Zool. Ill. 1st ser. t. 11* (1820). Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 334.

HYADES, Boisduval.

TENARIS (*part*), Hübner, *Verz. bek. Schmett.*

438. DRUSILLA HORSFIELDI, Swainson.

Drusilla Horsfieldii, Swainson, *Zool. Ill. 1st ser. t. 11* (1820). E. Doubleday, *List Lep. Brit. Mus. pt. I. App. p. 29.* Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 335.

Hyades Horsfieldii, Boisduval, *Spéc. Gén. Lép. t. 13, f. 1* (1836).

Morpho Urania, Zinken-Sommer, *Nova Acta Acad. Nat. Cur. XV.* p. 167.

a. ♂. Java. From Dr. Horsfield's Collection.

Genus THAUMANTIS, Hübner.

THAUMANTIS, Hübner, *Sammel. Exot. Schmett.* (1806-27). Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 335.

MORPHO (*part*), Godart. Horsfield. Zinken-Sommer.

439. THAUMANTIS ODANA, Godart Sp.

Morpho Odana, Godart, *Enc. Méth. IX.* p. 445 (1819). Horsfield, *Cat. Lep. Mus. E. I. C. t. 6, f. 5, 5a.*

Thaumantis Odana, Hübner, *Sammel. Exot. Schmett. Band II. tab.* (1806-27). Boisduval, *Spéc. Gén. Lép. I. t. 12 (8B), f. 1.* E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 115. Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 336, n. 1.

δ Morpho Klugius, *Zinken-Sommer*, in *Nova Acta Acad. Nat. Cur.* XVI. t. 15, f. 12, 13 (1834).

a. b. c. d. δ ♀. Java. From Dr. Horsfield's Collection.

440. *THAUMANTIS DIORES*, E. Doubleday.

Thaumantis Diores, E. Doubleday, *Ann. Nat. Hist.* XVI. p. 234 (1845). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 337, n. 2.

a. b. c. δ ♀. Cherra Poonjee. Presented by Colonel Buckley.

441. *THAUMANTIS RAMDEO*, Moore.

Thaumantis Ramdeo, n. sp.—Differs from *Thaum. Diores* in being larger, and in having the brilliant blue patch very large, it being an inch in diameter either way, and which extends over the whole of the middle of the wings. The under-side is paler than in *Th. Diores*, and wavy line running near the exterior margins of *Diores* is nearly obsolete in *Th. Ramdeo*; the two ocelli on the hind-wing are very small. Expanse of male 4, female nearly $4\frac{3}{4}$ inches.

a. b. c. δ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

442. *THAUMANTIS NOURMAHAL*, Westwood.

♀ *Thaumantis Nourmahal*, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 337, n. 8 (1851); id. *Trans. Ent. Soc.* IV. n. s. p. , t. , f. 1.

a. b. c. d. δ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

443. *THAUMANTIS LUCIPOR*, Westwood.

Thaumantis Lucipor, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 337, n. 5 (1851); id. *Trans. Ent. Soc.* IV. n. s. p. , t. , fig. 1, δ ; 2, ♀.

a. δ . Borneo. Presented by Hugh Low, Esq.

444. *THAUMANTIS NOUREDDIN*, Westwood.

Thaumantis Noureddin, Westwood, in *Doubleday and*

Hewitson's Diurnal Lep. X. p. 337, n. 6 (1851);
 id. *Trans. Ent. Soc.* IV. n. s. p. , t. , fig. 1, ♂;
 2, ♀.

a. ♀. Borneo. Presented by Hugh Low, Esq.

445. *THAUMANTIS CAMADEVA*, *Westwood*.

Thaumantis Camadeva, *Westwood*, *Cabinet of Orient.*
Ent. p. 9, t. 4, f. 1, 2 (1847); id. in *Doubleday and*
Hewitson's Diurnal Lep. p. 337, n. 7, t. 55, f. 2; id.
Trans. Ent. Soc. IV. n. s. p. .

a. b. ♂ ♀. Darjeeling. From Pearson's Collection.
 c. d. e. f. ♂ ♀. Darjeeling. From Messrs. Schlagint-
 weit's Collection.

Genus *DEBIS*, *Boisduval*.

DEBIS, *Boisduval*, *MS.* *Westwood*, in *Doubleday and Hewitson's*
Diurnal Lep. p. 358 (1851).

LETHE, *Hübner*, *Verz. bek. Schmett.* (1816).

OREAS MARMOREA (*part*), *Hübner*, *Samml. Exot. Schmett.*

CYLO (*part*), *E. Doubleday*.

446. *DEBIS EUROPA*, *Fabricius Sp.*

Papilio Europa, *Fabricius*, *Syst. Ent.* p. 500 (1775);
Ent. Syst. III. pt. I. p. 76.

Debis Europa, *Westwood*, in *Doubleday and Hewitson's*
Diurnal Lep. p. 359, n. 1.

Lethe Europa, *Hübner*, *Verz. bek. Schmett.* n. 534.

Satyrus Europa, *Godart*, *Enc. Méth.* IX. p. 478.

Oreas-Marmorea-Europa, *Hübner*, *Samml. Exot. Schmett.*
Band I. tab.

Cyllo Europa, *E. Doubleday*, *List Lep. Brit. Mus.* pt. I.
 p. 120.

♀ *Papilio Beroe*, *Cramer*, *Pap. Exot.* I. t. 79, f. C. D.
 (1779).

♂ *Papilio Arete*, *Cramer*, *Pap. Exot.* IV. t. 313, f. E.F.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Col-
 lection.

e. f. g. h. ♂ ♀. Purchased.

447. *DEBIS NEILGHERRIENSIS*, Guérin-Méneville Sp.

Satyrus (Cyllo) *Neelgheriensis*, *Guérin-Méneville*, in *Delessert's Voy. dans l'Inde*, pt. II. p. 74, t. 21, f. 1 (1843).

- a. ♂. Bootan. From Pemberton's Collection.
- b. c. ♂ ♀. N. India. Presented by Colonel Buckley.
- d. e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

448. *DEBIS ROHRIA*, Fabricius Sp.

Papilio Rohria, *Fabricius*, *Mant. Ins.* II. p. 45 (1787); *Ent. Syst.* III. I. p. 75.

Debis Rohria, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 360, n. 4.

Satyrus Rohria, *Godart*, *Ene. Méth.* IX. p. 479.

Satyrus Isana, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 448, tab. 16, f. 3, 4 (1844).

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
- e. f. ♂ ♀. N. India. Presented by Lieut. Hugo James.
- g. h. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

449. *DEBIS VERMA*, Kollar Sp.

Satyrus Verma, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 447, tab. 16, f. 1, 2 (1844).

Debis Verma, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 360, n. 5.

- a. b. ♂ ♀. Bootan. From Pemberton's Collection.
- c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

450. *DEBIS BHAIKAVA*, Moore.

Debis Bhairava, n. sp.—Female: UPPER-SIDE olive-brown: fore-wing with a transverse sharply-defined dusky line bordered with a series of small white spots, disposed obliquely from beyond the

middle of anterior to near posterior margin, one-fourth from the angle; also a subapical row of four small yellow spots, parallel with exterior margin: *hind-wing* with four submarginal round black spots encircled with pale brown, and medially disposed from near anterior to near anal angle. *UNDER-SIDE* luteous-brown: *fore-wing* with two transverse ferruginous-brown lines, the inner one bifid across discoidal-cell, the outer straight, and bordered by the white spots; a submarginal row of four pale ocelli, also an anterior terminal yellow spot: *hind-wing* with two transverse ferruginous-brown lines and a submarginal row of six ocelli; a narrow dark marginal line to both wings. Expanse $2\frac{5}{8}$ inches.

a. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

451. *DEBIS SCANDA*, Moore.

Debis Scanda, *n. sp.*—Male: *UPPER-SIDE* blackish, glossed with blue; along the margins pale purplish-blue: *hind-wing* with indistinct submarginal black spots. *UNDER-SIDE* deep ferruginous-brown: *fore-wing* with a transverse paler band across discoidal-cell, and another across the disc; a submarginal row of pale ocelli: *hind-wing* with two transverse darker lines, and a submarginal series of six, and a geminated seventh ocellus; margins pale, with a narrow darker line. Expanse $2\frac{1}{2}$ inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

452. *DEBIS NADA*, Moore.

Debis Nada, *n. sp.*—Female: *UPPER-SIDE* dark brown: *fore-wing* with two indistinct yellowish subapical marks: *hind-wing* with row of four small black submarginal spots. *UNDER-SIDE* deep bright ferruginous: *fore-wing* with transverse band across discoidal-cell, and another across the disc, yellowish; a submarginal row of pale ocelli: *hind-wing* with two transverse darker lines and a row of submarginal ocelli; margins pale, with a darker narrow line; ciliae white. Expanse $2\frac{5}{8}$ inches.

a. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Remark.—This may possibly be only the female of the preceding species.

453. *DEBIS ARCADIA*, Cramer Sp.

♀ Papilio Arcadia, Cramer, Pap. Exot. II. t. 116, f. E.F. (1779).

Debis Arcadia, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 359, n. 2.

Temenis Arcadia, Hübner, Verz. bek. Schmett. n. 279.

Satyrus Caumas, Godart, Enc. Méth. IX. p. 479, n. 7.

a. b. ♂. Java. From Dr. Horsfield's Collection.

454. *DEBIS MEKARA*, Moore.

Debis Mekara, n. sp.—Male: UPPER-SIDE glossy olive-brown: hind-wing with four round spots encircled with red. UNDER-SIDE with two transverse lines: fore-wing with five submarginal pale ocelli: hind-wing with six brown ocelli, encircled by a yellow and then a black line, the centre being minutely spangled with white. Expanse $2\frac{5}{8}$ inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

b. c. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

455. *DEBIS CHANDICA*, Moore.

Debis Chandica, n. sp.—Male: UPPER-SIDE dark brown, about the disc blackish: hind-wing with indistinct submarginal spots. UNDER-SIDE: fore-wing pale drab-brown, and having a purple tinge; along costal margin, and broadly on exterior margin, yellowish; transverse red line crossing from middle of discoidal-cell, and another irregular brown line across the disc; five pale ocelli; straight marginal and wavy submarginal line: hind-wing drab-brown; two irregular zigzag transverse lines, anteriorly within yellowish, posteriorly ferruginous-brown; six ocelli encircled with a yellow and brown line, the three anterior, and last posterior, black, the other two drab, all minutely spangled with white; exterior margin yellowish; marginal line, and submarginal row of ill-defined lunular marks, ferruginous-brown. Female: UPPER-SIDE: fore-wing with the basal half obliquely brownish-red, anterior half brown, with a band obliquely from middle of costal margin, terminated by two spots, white: also a small white subapical spot: hind-wing brownish-red, anterior angle and along exterior margin yellowish; a submarginal row of indistinct spots and marginal line black. UNDER-SIDE as in male, excepting that

the oblique band of fore-wing is more distinct. Expanse of male $2\frac{1}{2}$, female $2\frac{5}{8}$ inches.

- a. b. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- c. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.
- d. ♂. Java. From Mr. Argent's Collection.

456. *DEBIS KANSA*, Moore.

Debis Kansa, n. sp.—Male: UPPER-SIDE glossy olive-brown; hind-wing with caudal appendage somewhat long; marginal row of blackish spots indistinct. UNDER-SIDE drab, with two transverse dark lines; fore-wing with four pale ocelli: hind-wing with six small black ocelli, encircled by a yellow, brown, and then a lavender-coloured line, the sixth minute and geminated. Female: UPPER-SIDE dark brown: fore-wing with darker line across the disc: hind-wing with five large black spots, narrowly encircled with red, from anterior angle. UNDER-SIDE brown, darkest within the outer transverse line; the inner line within, about the apex of fore-wing, glossed with lavender-colour; ocelli pale: hind-wing with the anterior first and fifth ocellus large, that at the anal angle geminated, all encircled with a glossy lavender band. Expanse $2\frac{6}{8}$ inches.

- a. b. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus ZOPHOESSA, Doubleday.

ZOPHOESSA, E. Doubleday. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 362 (1851).

457. *ZOPHOESSA SURA*, E. Doubleday.

Zophoessa Sura, E. Doubleday, MS. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 362, t. 61, fig. 1 (1851).

- a. ♂. Bootan. From Pemberton's Collection.
- b. c. ♂. Darjeeling. From Pearson's Collection.
- d. ♂. N. India. Presented by Lieut. Hugo James.
- e. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

458. *ZOPHOESSA YAMA*, Moore.

Zophoessa Yama, n. sp.—Differs from *Zoph. Sura* in having no caudal appendage to the hind-wing; the under-side having on the fore-wing three pale bands crossing the discoidal-cell, and four submarginal ocelli: the hind-wing having the basal portion irrorated with minute pale dots, the transverse streaks more oblique, and the ocelli, seven in number, are black, encircled with a pale and then a dark line, centred with a minute white dot, that at the anal angle being, as it were, geminated; ciliae seen from upper-side yellowish-white. Sexes alike. Expanse 3 inches.

a. b. ♂. Bootan. From Pemberton's Collection.

c. ♀. N. India. Presented by Colonel Buckley.

Genus CYLLO, Boisduval.

CYLLO, *Boisduval*, *Voy. de l'Astrolabe*, *Entom. pt. I.* p. 140 (1832).

HIPIO, *pt. Hübner*, *Verz. bek. Schmett.* (1816).

OREAS, *pt. Hübner*.

CYLLO, *pt. E. Doubleday*. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 360.

459. *CYLLO CONSTANTIA*, Cramer Sp.

Papilio Constantia, *Cramer*, *Pap. Exot. II. t. 133, f. A.B.* (1779).

Cyllo Constantia, *Boisduval*, *Voy. de l'Astrolabe*, *Ent. pt. I.* p. 141. *E. Doubleday*, *List Lep. Brit. Mus.* *pt. I.* p. 120. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 361, n. 5.

Hipio Constantia, *Hübner*, *Verz. bek. Schmett.* n. 537.

Satyrus Constantia, *Godart*, *Enc. Méth. IX.* p. 477.

a. ♂. N. India. Purchased.

b. c. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

460. *CYLLO CRISHNA*, Westwood.

Cyllo Crishna, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 361, n. 9 (1851).

a. Java. From Mr. Argent's Collection.

Genus MELANITIS, *Fabricius*.

MELANITIS, *Fabricius*, *Syst. Gloss.* (*Illiger's Mag.* VI. p. 282, 1807).
HIPIO, *pt.* *Hübner, Verz. bek.* Schmett.

CYLO, *pt.* *Boisduval.* *E. Doubleday.* *Westwood* (*Diurnal Lep.*).

461. MELANITIS LEDA, *Linnæus* Sp.

Papilio Leda, *Linnæus*, *Syst. Nat.* II. p. 773 (1767).

Fabricius, *Syst. Ent.* III. *pt.* I. p. 108. Cramer,
III. *Pap. Exot.* t. 196, f. C.D.; 292, f. A. Drury,
Ill. Ent. I. t. 15, f. 5, 6.

Melanitis Leda, *Fabricius*, *Syst. Gloss.* (*Illiger's Mag.* VI. p. 282).

♂ Oreas-Marmorea-Leda, *Hübner, Samml. Exot.* Schmett.
Band I. tab.

Satyrus Leda, *Godart, Enc. Méth.* IX. p. 478.

Hipparchia Leda, *Horsfield, Cat. Lep. Mus.* E.I.C.
t. 8, f. 9.

Cyllo Leda, *Boisduval, Voy. de l'Astrolabe, Ent. pt.* I.
p. 142. *E. Doubleday, List Lep. Brit. Mus.* pt. I.
p. 121. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 361, n. 1.

Papilio Solandra, *Fabricius, Syst. Ent.* p. 500. *Donovan, Ins. of New Holland*, t. 23, f. 1.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

g. ♂. Borneo. Presented by Edwin Shepherd, Esq., 1857.

The larva and pupa of *Melanitis Leda* are figured on Plate VI., fig. 8, 8a, from Java.

462. MELANITIS BANKSIA, *Fabricius* Sp.

Papilio Banksia, *Fabricius, Syst. Ent.* p. 499 (1775).
Donovan, Ins. of New Holland, t. 25, f. 1.

Hipio Leda, *Hübner, Verz. bek.* Schmett. n. 538.

Satyrus Banksia, *Godart, Enc. Méth.* IX. p. 477.

Cyllo Banksia, *E. Doubleday, List Lep. Brit. Mus.* pt. I.

p. 121. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 361, n. 3.

Papilio Ismene, *Cramer*, *Pap. Exot.* I. t. 26, f. A.B. (1779).

Papilio Mycena, *Cramer*, *Pap. Exot.* IV. t. 291, f. F. (1782).

Papilio Phedima, *Cramer*, *Pap. Exot.* IV. t. 292, f. B.

Papilio Arcensia, *Cramer*, *Pap. Exot.* IV. t. 292, f. C.

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

f. g. ♂ ♀. N. India. From the Asiatic Society of Bengal.

h. . Darjeeling. From Pearson's Collection.

i. N. India. Presented by Colonel Sykes.

463. *MELANITIS VAMANA*, Moore.

Melanitis Vamana, n. sp.—Differs from *Melanitis Banksia* in being larger, the subapical ferruginous patch on fore-wing occupying a larger space, the deep-black patch extends to the costal vein, the upper white spots smaller, the lower being very indistinct; two white dots on hind-wing. UNDER-SIDE marked as in some specimens of *Melanitis Banksia*. Expanse $3\frac{1}{2}$ inches.

a. b. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

c. ♂. N. India. Presented by Colonel Sykes.

464. *MELANITIS AMBASARA*, Moore.

Melanitis Ambasara, n. sp.—UPPER-SIDE brown, having an olive gloss: fore-wing with an ill-defined oblique ochreous streak, bounded anteriorly on both sides with dusky-black; the streak with two small indistinct spots: hind-wing with very white spots. UNDER-SIDE marked as in some specimens of *Melanitis Banksia*. Expanse $3\frac{1}{8}$ inches.

a. ♂. Java. From Dr. Horsfield's Collection.

b. ♀. Java. From Mr. Argent's Collection.

465. *MELANITIS BELA*, Moore.

Melanitis Bela, n. sp.—UPPER-SIDE deep dusky-brown: fore-wing with small indistinct ferruginous streak, below which the space is

blackish; the two usual white spots: *hind-wing* with one minute white dot. UNDER-SIDE deep mottled ferruginous-brown: triangular patch near apex of *fore-wing* paler: *hind-wing* with six spots. Expanse $2\frac{7}{8}$ inches.

a. ♂. N. India. Presented by Lieutenant Hugo James.

b. ♂. Assam. From McClelland's Collection.

c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

466. *MELANITIS SUYUDANA*, Moore.

Melanitis Suyudana, n. sp.—UPPER-SIDE deep dark-brown, paler on the margins: a single whitish spot near apex of *fore-wing*, and a minute dot on *hind-wing*. UNDER-SIDE deep mottled ferruginous-brown, with indistinct darker usually transverse streaks; a triangular space of mottled greyish-white from costal margin near the apex: *hind-wing* with six small pale spots centred with a white dot. Expanse $2\frac{5}{8}$ inches.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

467. *MELANITIS VARAHA*, Moore.

Melanitis Varaha, n. sp.—UPPER-SIDE dusky-brown, ashy-brown along exterior margins; subapical patch indistinctly black, with two white spots. UNDER-SIDE ashy, tinged with olive from the base to the middle of wing: near apex of *fore-wing* two small brown dots: *hind-wing* with submarginal row of indistinct paler spots. Expanse $2\frac{3}{8}$ inches.

a. ♂. Canara. Presented by S. N. Ward, Esq.

468. *MELANITIS GOKALA*, Moore.

Melanitis Gokala, n. sp.—UPPER-SIDE brown, suffused with ashy about the apex and exterior margins; an indistinct blackish subapical patch (without spots). UNDER-SIDE glossy olive-brown, washed on the basal two-thirds with pulverulent ashy-white; a small white dot on lower disco-cellular vein, and another near anal angle of *hind-wing*. Expanse 3 inches.

a. ♂. Canara, S. India. Presented by S. N. Ward, Esq.

469. *MELANITIS SURADEVA*, Moore.

Melanitis Suradeva, n. sp.—Male: UPPER-SIDE deep vinaceous-brown, darkest at the apex, and having in some lights a purple gloss: fore-wing with a large black patch in middle; a short narrow obliquely-curved yellow subapical streak. UNDER-SIDE dull-ochreous, suffused with brown, and covered with short, narrow, undulating striae, as in *Melanitis Leda*; a transverse darker streak across middle of both wings; four small whitish submarginal spots on fore-wing, and two near anal angle. Expanse 3 inches.

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus CŒLITES, Boisduval.

CŒLITES, Boisduval, MS. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 367.

470. *CŒLITES EPIMINTHIA*, Boisduval.

Cœlites Epiminthia, Boisduval, MS. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 368 (1851).

a. ♂. Borneo. Presented by Hugh Low, Esq.

Genus ORINOMA, E. Doubleday.

ORINOMA, E. Doubleday, in Gray's Lep. of Nepal, p. 14 (1846). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 368.

471. *ORINOMA DAMARIS*, G. R. Gray Sp.

Pieris Damaris, G. R. Gray, Lep. of Nepal (1833).

Orinoma Damaris, E. Doubleday, in Gray's Lep. of Nepal, p. 14, t. 7, f. 2, 2a. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 369, t. 63, f. 3.

Satyrus Damaris, E. Doubleday, List Lep. Brit. Mus. pt. I. p. 122.

a. b. c. ♂. Silhet. From Colonel Buckley's Collection.

d. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus NEORINA, Westwood.

NEORINA, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 369 (1851).

472. *NEORINA HILDA*, Westwood.

Neorina Hilda, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 370, t. 65, f. 2 (1851).

a. b. c. d. ♂ ♀. Darjeeling. From Pearson's Collection.

e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus *RAGADIA*, *Westwood*.

NEONYMPHA (*RAGADIA*), *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 376 (1851).

473. *RAGADIA MAKUTA*, Horsfield Sp.

Hipparchia Makuta, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 5, f. 9, 9a (1829).

Euptychia Crisia, *Hübner*, *Zuträge*, f. 675, 676 (1832-7).

Neonympha (*Ragadia*) *Crisia*, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 376, n. 26.

a. b. ♂. Java. From Dr. Horsfield's Collection.

c. ♂. Java. From Mr. Argent's Collection.

Genus *EREBIA*, *Dalman*.

EREBIA, *p. Dalman, P. S.* p. 81 (1816).

EPIGEA, *MELAMPIAS*, *Hübner*, *Verz. bek. Schmett.* (1816).

OREINA, *Westwood*, *Brit. Butt.* p. 76 (1840).

EREBIA, *Boisduval*. *Herr.-Schäffer*. *E. Doubleday*. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 376.

474. *EREBIA SCANDA*, Kollar.

Erebia Scanda, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 452, t. 17, f. 3, 4 (1844).

a. ♂. N. India. Presented by Colonel Buckley.

b. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

475. *EREBIA ANNADA*, Moore.

Erebia Annada, *n. sp.*—Differs from *Erebia Scanda* in being larger, and in having the anal angle more falcate. On the under-side, the

fore-wing is of a bright ferruginous: the *hind-wing* is mottled with white nearly throughout, and from near anal angle, below the ocelli, extends a curved white line. *Expanse* $2\frac{1}{2}$ inches.

a. b. c. ♂ ♀. Bootan. From Pemberton's Collection.

Genus LASIOMMATA, *Westwood.*

LASIOMMATA, *Westwood*, *Brit. Bult.* p. 65 (1840); id. *Diurnal Lep.* p. 385.

DIRA, PARARGE, EPINEPHILE, *pt. Hübner.*

476. LASIOMMATA SCHAKRA, *Kollar Sp.*

Satyrus Schakra, *Kollar*, in *Hügel's Kaschmir*, IV. *pt. II.* p. 446, *t. 15, f. 3, 4* (1844).

Lasiommata Schakra, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 387, *n. 12.*

a. b. ♂. N. India. Presented by Colonel Hearsey.

c. d. ♂ ♀. N. India. Presented by Colonel Buckley.

e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

477. LASIOMMATA? PULAHĀ, *Moore.*

Lasiommata? Pulaha, n. sp.—UPPER-SIDE chocolate-brown, dusky about the apex and margins: *fore-wing*, with the veins at the base, a stripe at extremity of discoidal-cell, and two transverse series of spots, ochreous-yellow: *hind-wing* with two rows of ochreous-yellow spots; ciliae spotted with yellowish-white. UNDER-SIDE dark brown: *fore-wing* with three bars across the discoidal-cell and subapical spots tinged with white; lower series of spots and space below the cell yellow: *hind-wing* with submarginal row of eight ocelli and a single ocellus at abdominal angle; an irregular dark-brown band from middle of anterior to middle of abdominal margin; markings about anterior margin tinged with white. *Expanse* $2\frac{5}{8}$ inches.

a. ♂. Bootan. From Pemberton's Collection.

Remark.—This species is somewhat allied in form to *Las. Merope*.

478. LASIOMMATA? BHADRA, *Moore.*

Lasiommata? Bhadra, n. sp.—UPPER-SIDE blackish-brown, suffused with chocolate-brown at the base of fore-wing and along abdominal margin: *fore-wing* with two transverse bars within discoidal-cell and

two rows of spots across the wing yellowish, the lower inner spot being long: *hind-wing* with two rows of irregular-shaped spots, and small space about extremity of discoidal-cell deep ochreous-yellow, the last spot on each row near anal angle encircling a spot of black. **UNDER-SIDE** dark brown, tinged in parts with pink: *fore-wing* with markings as above; also with a subapical eye-spot: *hind-wing* with a series of seven ocelli and a smaller spot at abdominal angle; base of the wing varied with pale ochreous-white and dark-brown markings. Expanse $3\frac{1}{2}$ inches.

a. b. c. d. ♂ ♀. Darjeeling. From Indian Collection,
Exposition Universelle at Paris, 1855.

Remark.—The two preceding species may be generically separated from LASIOMMATA, under the name of ENOPE, Moore.

Genus XENICA, Westwood.

LASIOMMATA (XENICA), Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 387.

479. XENICA ACHANTA, Donovan Sp.

Papilio Achanta, Donovan, *Ins. of New Holland*, t. 22, f. 2 (1805).

Lasiommata (Xenica) Achanta, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 387, n. 22.

Satyrus Achanta, Godart, *Enc. Méth. IX.* p. 500. Boisduval, *Voy. de l'Astrolabe, Ent. pt. I.* p. 147.

Tisiphone Acanthe, Hübner, *Zuträge*, f. 267, 268 (1818).

a. ♂. Sumatra? From Raffles's Collection.

Genus SATYRUS, Latreille.

SATYRUS, pt. Latreille.

HIPPARCHIA, pt. Fabricius.

HIPPARCHIA, MINOIS, EUMENIS, OENEIS, PYRONIA, EPINEPHILE, TISIPHONE, et ENODIA, pt. Hübner.

480. SATYRUS PADMA, Kollar.

Satyrus Padma, Kollar, in Hügel's *Kaschmir*, IV. pt. II. p. 445, t. 15, f. 1, 2.

a. ♂. N. India. Presented by Edwin Shepherd, Esq.

481. *SATYRUS AVATARA*, Moore.

Satyrus Avatara, n. sp.—Differs from *Sat. Padma* in having the transverse maculated band much narrower, and there being no spots running obliquely to costal margin. It is less in size than *Pat. Padma*, but larger than *Satyrus Swaha* and *Saraswati*. Expanse $3\frac{2}{10}$ inches.

a. b. ♂. N. India. Presented by Colonel Buckley.

482. *SATYRUS SWAHA*, Kollar.

Satyrus Swaha, Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 444, t. 14, f. 1, 2 (1844). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 390, n. 18.

a. ♂. N. India. From the Asiatic Society of Bengal.

b. c. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

483. *SATYRUS SARASWATI*, Kollar.

Satyrus Saraswati, Kollar, in *Hügel's Kaschmir*, IV. pt. II. p. 445 (March, 1844). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 390, n. 19.

♂ ♀ *Satyrus Brahminus*, Blanchard, in *Jacquemont's Voy. dans l'Inde*, Ins. p. 22, t. 2, f. 4, 5, 6 (1844). *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 390, n. 16.

Satyrus Jacquemontii, Boisduval, MS.

a. b. ♂ ♀. N. W. India. Presented by Col. Hearsey.

Genus ERITES, Boisduval.

ERITES, Boisduval, MS.

SATYRUS (ERITES), Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 392 (1851).

484. *ERITES MADURA*, Horsfield Sp.

Hipparchia Madura, Horsfield, *Cat. Lep. Mus. E.I.C.* t. 5, f. 8, 8a (1829).

Satyrus (Erites) Madura, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 392.

Erites Medura, Boisduval, MS.

a. ♀. Java. From Dr. Horsfield's Collection.

Genus MYCALESIS, Hübner.

MYCALESIS, *Hübner, Verz. bek. Schmett.* (1816). *E. Doubleday, List Lep. Brit. Mus.* p. 138. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 392.

485. *MYCALESIS POLYDECTA*, Fabricius Sp.

Papilio Polydecta, *Fabricius, Spec. Ins.* II. p. 85 (1781); *Ent. Syst.* III. pt. I. p. 107. *Cramer, Pap. Exot.* II. t. 144, f. E.F.

Mycalesis Polydecta, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 391, n. 1.

Satyrus Polydecta, *Godart, Enc. Méth.* IX. p. 511.

a. ♂. China. Presented by Dr. Cantor.

486. *MYCALESIS OTREA*, Cramer Sp.

Papilio Otrea, *Cramer, Pap. Exot.* IV. t. 314, f. A.B. (1782).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂. N. India. From the Asiatic Society of Bengal.

g. h. ♂ ♀. Bootan. From Pemberton's Collection.

487. *MYCALESIS FRANCISCA*, Cramer Sp.

Papilio Francisca, *Cramer, Pap. Exot.* IV. t. 326, f. E.F. (1782).

a. ♂. Assam. From McClelland's Collection.

b. c. ♂. Canara. Presented by S. N. Ward, Esq.

488. *MYCALESIS VISALA*, Moore.

Mycalesis Visala, n. sp.—This species is larger than the preceding, and has the apex of the fore-wing sharply angulated. The under-side has two transverse lines across both wings, the outer line across the middle, the inner towards the base, of the wing; the row of dots minute, white, the third from anal angle being black. Expanse $2\frac{1}{4}$ inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

b. ♀. Bootan. From Pemberton's Collection.

c. ♀. N. India. Presented by Colonel Buckley.

489. *MYCALESIS SANATANA*, Moore.

Mycalesis Sanatana, n. sp.—UPPER-SIDE brown: fore-wing with a large lower, and a smaller upper eye-spot, the latter with a minute one above and below it. UNDER-SIDE ochreous-brown; the series of spots white. Expanse 2 inches.

- a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

490. *MYCALESIS MAMERTA*, Cramer Sp.

Papilio Mamerta, Cramer, Pap. Exot. IV. t. 326, f. D. (1782).

- a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

491. *MYCALESIS DRUSIA*, Cramer Sp.

Papilio Drusia, Cramer, Pap. Exot. I. t. 84, f. C.D. (1779).

Var. *Papilio Justinia*, Cramer, Pap. Exot. IV. t. 326, f. C.

? *Papilio Mineus*, Linnæus, Syst. Nat. II. p. 768.

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. g. ♂ ♀. N. India. Presented by Colonel Sykes.

h. ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

i. ♂. (*Justinia*, Cramer.) N. India.

492. *MYCALESIS MALSARA*, Moore.

Mycalesis Malsara, n. sp.—UPPER-SIDE dark brown: fore-wing with a transverse brownish-white band, and upper and lower ocelli: hind-wing with very indistinct transverse brownish-white streak, and two small ocelli. UNDER-SIDE with transverse yellowish-white band well defined across both wings: fore-wing with five ocelli, the first and third minute: hind-wing with seven larger ocelli, the third and seventh smallest. Expanse $2\frac{1}{4}$ inches.

- a. b. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

493. *MYCALESIS GOTAMA*, Moore.

Mycalesce Gotama, n. sp.—UPPER-SIDE pale brown : fore-wing with upper and lower ocelli : hind-wing with a minute ocellus. UNDER-SIDE pale greyish-brown : fore-wing with transverse yellowish-white band, and the two ocelli as above : hind-wing with transverse yellowish-white band ; three ocelli from anterior margin near angle, the two lower very small ; and three near anal angle, the two lower also very small. Sexes alike. Expanse, ♂ $1\frac{5}{8}$, ♀ 2 inches.

a. b. ♂ ♀. China. Presented by Dr. Cantor.

494. *MYCALESIS PATNIA*, Moore.

Mycalesce Patnia, n. sp.—UPPER-SIDE ochreous-brown : fore-wing with a lower black spot, centred with white, and broadly bordered on the upper half (but not on the lower) with pale ochreous ; a streak from thence to base pale ochreous ; near apex a small longitudinal white spot ; two narrow marginal lines to both wings. UNDER-SIDE ochreous-yellow : fore-wing with the upper longitudinal white spot ; also the large lower ocellus, the lower border of which is suffused with black ; five transverse lines commencing from base, and circle round the upper spot and lower ocellus, ochreous-red ; a silvery line across the middle of wing and round the ochreous-red circle : hind-wing with single upper and two lower white spots, distantly encircled with a black line, around this and two transverse basal lines ochreous-red ; a transverse line across middle of wing, and round the outer circle of the ocelli, silvery. Sexes alike. Expanse $1\frac{5}{8}$ to $1\frac{3}{4}$ inches.

a. b. ♂ ♀. Ceylon. From M. Jonville's Collection.

495. *MYCALESIS ANAPITA*, Moore.

Mycalesce Anapita, n. sp.—UPPER-SIDE ochreous-red : fore-wing with the costal, exterior, and to middle of posterior margin, dark brown ; a small lower black spot centred with white and encircled with red : hind-wing with two narrow marginal and broader submarginal lines dark brown, with three lower ocelli. UNDER-SIDE yellow, with two ochreous-red transverse lines crossing middle of both wings and bordering the ocelli on both wings ; marginal lines narrow, black. Expanse $1\frac{3}{4}$ inches.

a. ♂. ? Sumatra. ? From Raffles's Collection.

Remark.—This species is closely allied to *Myc. Remulia*, Cramer, t. 237, f. F. G.

496. *MYCALESIS HERI*, Moore.

Mycalesis Heri, n. sp.—UPPER-SIDE brown: *fore-wing* with large upper and larger lower ocellus: *hind-wing* with two larger and third smaller ocelli. UNDER-SIDE paler, the base covered with short brown undulated striae; transverse line, inner and outer border to the ocelli pale ochreous: *fore-wing* with the two ocelli as above: *hind-wing* with six ocelli; marginal lines brown and pale ochreous. Expanse $2\frac{3}{8}$ inches.

a. b. c. ♂. Bootan. From Pemberton's Collection.

497. *MYCALESIS SAFITZA*, Hewitson.

Mycalesis Safitza, Hewitson, MS. Westwood, in Double-day and Hewitson's Diurnal Lep. p. 394, n. 10, t. 66, f. 3 (1851).

a. ♂. Africa. Presented by Edwin Shepherd, Esq., 1857.

498. *MYCALESIS SAMBA*, Moore.

Mycalesis Samba, n. sp.—UPPER-SIDE dark brown, exterior margins paler; narrow marginal line black. UNDER-SIDE with transverse white line: *fore-wing* with two ocelli: *hind-wing* with seven ocelli, the anterior second and third very small; exterior margins pale, with narrow marginal lines. Expanse $1\frac{3}{4}$ inch:

a. ♂. N. India. Donor unknown.

Remark.—This species is somewhat allied to *Myc. Hesione*, from which it may be distinguished by having seven ocelli instead of three on the under-side of the hind-wing.

499. *MYCALESIS MANDATA*, Moore.

Mycalesis Mandata, n. sp.—UPPER-SIDE brown, with transverse band of under-side indistinctly seen through the wing. UNDER-SIDE with broad transverse very white band crossing both wings; *fore-wing* with two small ocelli: *hind-wing* with three ocelli, the anterior being small, and all encircled with a silvery line. Expanse $1\frac{1}{2}$ inch.

a. ♂. Canara. Presented by S. N. Ward, Esq.

Remark.—Differs from *Myc. Hesione* on the under-side in having the transverse band three times the width that it is in that species.

500. *MYCALESIS HESIONE*, Cramer Sp.

Papilio Hesione, Cramer, *Pap. Exot.* I. t. 11, f. C.D. (1779). *Fabricius*, *Ent. Syst.* III. pt. I. p. 100.

Mycalesis Hesione, E. *Doubleday*, *List Lep. Brit. Mus.* pt. I. p. 139. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 393, n. 4.

Satyrus Hesione, Godart, *Enc. Méth.* IX. p. 510. *Boisduval*, *Voy. de l'Astrolabe*, *Ent. pt. I.* p. 154.

Papilio Doris, Cramer, *Pap. Exot.* IV. t. 362, f. C.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. ♂. N. India. Presented by Colonel Sykes.

501. *MYCALESIS RUNEKA*, Moore.

Mycalesis Runeka, n. sp.—UPPER-SIDE uniform dark brown. UNDER-SIDE dark brown, paler along the exterior margins; a transverse yellowish-white band crossing both wings: fore-wing with two, and hind-wing with three, pale-brown ocelli.

a. b. c. d. ♂ ♀. Assam. From McClelland's Collection.

502. *MYCALESIS JANARDANA*, Moore.

Mycalesis Janardana, n. sp.—Male: UPPER-SIDE dusky-brown, palest on the hind-wing, which has two rather indistinct black spots. UNDER-SIDE brown, the basal parts being covered with small narrow striae; two transverse lines, the outer margined exteriorly with white: fore-wing with six and hind-wing with seven small ocelli. Female paler on the UPPER and UNDER-SIDE, the latter nearly covered with small dark striae; the transverse lines paler; the ocelli somewhat larger than in male. Expanse of ♂ $1\frac{1}{2}$; ♀ $1\frac{7}{10}$ inch.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

503. *MYCALESIS? HIMACHALA*, Moore.

Mycalesis? Himachala, n. sp.—UPPER-SIDE deep brown, paler broadly along exterior margins: fore-wing with six and hind-wing with five ocelli; marginal line deep brown. UNDER-SIDE as above, but the hind-wing with six ocelli, the anterior ocellus large, and the posterior with two white dots. Expanse 2 $\frac{3}{4}$ to 3 inches.

a. b. c. d. e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

This species may be separated under the generic name of THEOPE, Moore.

Genus YPHTHIMA, Hübner.

YPHTHIMA, Hübner, *Zuträge*, f. 83 (1818). E. Doubleday. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 394.

504. YPHTHIMA LYSANDRA, Cramer Sp.

Papilio Lysandra, Cramer, *Pap. Exot. IV.* t. 293, f. G.II. (1782).

Papilio Baldus, Fabricius, *Mant. Ins. II.* p. 34 (1787); *Ent. Syst. III. pt. I.* p. 323. Donovan, *Ins. of India*, t. 36, f. 2.

Yphthima Baldus, E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 139. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 395.

Satyrus Baldus, Godart, *Enc. Méth. IX.* p. 551.

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. h. i. j. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

505. YPHTHIMA PHIOMELA, Hübner.

Yphthima Philomela, Hübner, *Zuträge*, f. 83, 84 (1818). E. Doubleday, *List Lep. Brit. Mus. pt. I.* p. 138. Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 395, n. 3.

Satyrus Chenu, Guérin-Méneville, in Delessert's *Souv. Voy. Ind.* p. 77, t. 21, f. 2 (1843).

Yphthima Chenu, Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 396, n. 9.

a. ♂. N. India. From the Asiatic Society of Bengal.

506. YPHTHIMA PANDOCUS, Boisduval MS.

Yphthima Pandocus, n. sp.—UPPER-SIDE as in *Yph. Lysandra*, but with only two ocelli on hind-wing, the lower ocellus being at anal angle and very small. UNDER-SIDE with the hind-wing having only two ocelli. Expanse of ♂ $1\frac{1}{2}$; ♀ $1\frac{3}{4}$ inch.

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. h. ♂. N. India? From the Asiatic Society of Bengal.

507. *YPHTHIMA HYAGRIVA*, Moore.

Yphthima Hyagriva, n. sp.—UPPER-SIDE with only two ocelli, one being at apex of *fore-wing*, the other on *hind-wing* near anal angle. UNDER-SIDE yellowish: *hind-wing* covered with short narrow undulated striae, with two anterior and two posterior ocelli; a marginal line and submarginal streak blackish. Expanse 1½ inch.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

508. *YPHTHIMA SAKRA*, Moore.

Yphthima Sakra, n. sp.—UPPER-SIDE with five ocelli, one being on *fore-wing*, three near anal angle, and one (but indistinct) near anterior angle of *hind-wing*. UNDER-SIDE yellow, covered with short narrow dark-brown striae: *hind-wing* with geminated anterior, and three single posterior ocelli. Expanse 2 inches.

a. b. c. d. ♂ ♀. Darjeeling. From Pearson's Collection.

e. f. ♂. Bootan. From Pemberton's Collection.

g. h. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

509. *YPHTHIMA NARASINGHA*, Moore.

Yphthima Narasingha, n. sp.—UPPER-SIDE with only one ocellus on *fore-wing*. UNDER-SIDE greenish-grey, covered with short undulated darker striae; the one ocellus on *fore-wing* only, beneath which are two descending dark lines to posterior margin. Expanse 2 inches.

a. b. c. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus ELYMNIAS, Hübner.

ELYMNIAS, Hübner, Verz. bek. Schmett. (1816).

MELANITIS, pt. Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 282).

MELANITIS, Horsfield. Boisduval. E. Doubleday. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 403.

BIBLIS, pt. Godart.

510. *ELYMNIAS LAIS*, Fabricius Sp.

Papilio Lais, *Fabricius*, *Spec. Ins.* II. p. 102 (1781);
Ent. Syst. III. pt. I. p. 58. *Cramer, Pap. Exot.* II.
t. 110, f. A.B. (1779).

Elymnias Lais, *Hübner, Verz. bek. Schmett.* n. 326.

Biblis Lais, *Godart, Enc. Méth.* IX. p. 326.

Melanitis Lais, *Boisduval. E. Doubleday, List Lep.*
Brit. Mus. pt. I. p. 143. *Westwood, in Doubleday*
and Hewitson's Diurnal Lep. p. 404, n. 1.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

511. *ELYMNIAS UNDULARIS*, Fabricius Sp.

♂ Papilio undularis, *Fabricius, Spec. Ins. App.* p. 504
(1781); *Ent. Syst.* III. pt. I. p. 127. *Drury,*
Ill. Exot. Ins. II. t. 10, f. 1, 2. *Cramer, Pap.*
Exot. III. t. 256, f. A.B. (*Tanysp.*)

Biblis undularis, *Godart, Enc. Méth.* IX. p. 326.

Melanitis undularis, *Horsfield, Cat. Lep. Mus. E.I.C.*
t. 3, f. 24, t. 8, f. 8. E. Doubleday, List Lep. Brit.
Mus. pt. I. p. 144. *Westwood, in Doubleday and*
Hewitson's Diurnal Lep. p. 404, n. 2.

♀ Papilio Protogenia, *Cramer, Pap. Exot.* II. t. 189, f. F.G.
Fabricius, Ent. Syst. III. pt. I. p. 117.

Elymnias Protogenia, *Hübner, Verz. bek. Schmett.* n. 323.

Biblis Protogenia, *Godart, Enc. Méth.* IX. p. 327.

♂ Elymnias Jynx, *Hübner, Zuträge*, f. 37, 38.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. N. India. Presented by Colonel Sykes.

g. h. i. j. k. ♂ ♀ var. N. India. Presented by
Colonel Buckley.

l. ♂ var. Borneo. Presented by Edwin Shepherd, Esq.

The larva and pupa of *Elymnias undularis* are figured on Plate VI.,
fig. 7, 7a, from Java. "Feeds on the Gobbang (*Corypha umbraculifera*). May."—(Horsfield, MS.)

512. *ELYMNIAS DUSARA*, Horsfield Sp.

Melanitis Dusara, Horsfield, Cat. Lep. Mus. E.I.C. t. 5, f. 7 (1829). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 405, n. 5.

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

g. ♂ var. Borneo. Presented by Hugh Low, Esq.

513. *ELYMNIAS VASUDEVA*, Moore.

Elymnias Vasudeva, n. sp.—Male: UPPER-SIDE, *fore-wing* black, glossed with slaty-blue; costal margin with small bluish-white dots; longitudinal spots in middle of wing, and spots from near apex to posterior angle bluish-white: *hind-wing*, from abdominal margin to middle of wing, white, anterior and posterior border black, the latter with some marginal white spots. UNDER-SIDE white, veins black: *fore-wing* covered throughout with deep-black striæ: *hind-wing* with exterior margin covered with deep-black striæ; base of wing black, with numerous small white spots, and a large red spot below the cell; abdominal margin broadly suffused with yellow. Expanse $2\frac{3}{4}$ inches.

a. ♂. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

514. *ELYMNIAS PATNA*, Westwood Sp.

Melanitis Patna, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 405, n. 6, t. 68, f. 2 (1851).

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

515. *ELYMNIAS LEUCOCYMA*, Godart Sp.

Biblis Leucocyma, Godart, Enc. Méth. IX. p. 326 (1819). *Melanitis Leucocyma*, Boisduval. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 144. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 404, n. 4.

a. b. c. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

d. e. f. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

516. *ELYMNIAS KAMARA*, Moore.

Elymnias Kamara, n. sp.—Male: UPPER-SIDE dark ferruginous-brown, glossed with purple: *hind-wing* with three rows of ochreous-white spots along exterior margin. UNDER-SIDE paler brown, suffused with dusky-black, and covered with darker striae; a dark line, with pale inner border, along exterior margin. Female: UPPER-SIDE ferruginous-brown: *fore-wing* with a series of ochreous-white submarginal spots from posterior angle: *hind-wing* with a series of ochreous-white markings between the veins. UNDER-SIDE ochreous, suffused with pale brown, and covered with dark-brown striae. EXPANSE of ♂ $2\frac{3}{4}$; ♀ $3\frac{1}{8}$ inches.

a. b. ♂ ♀. Java. From Mr. Argent's Collection.

Remark.—This species is somewhat allied to *Elymnias Casiphone*, Hübner.

Genus HYPANIS, Boisduval.

HYPANIS, Boisduval, Spéc. Gén. Lép. t. IX. f. 6 (1836); id. Lép. Madagas. t. 7, f. 5. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 410.

BIBLIA, pt. Hübner.

517. *HYPANIS ILITHYIA*, Drury Sp.

Papilio Ilithyia, Drury, Ill. II. t. 17, f. 1, 2 (1773).

Biblis Ilithyia, Godart, Enc. Méth. IX. p. 327.

Papilio Polinice, Cramer, Pap. Exot. IV. t. 375, f. G. II. (1782).

Hypanis Polinice, Boisduval, Spéc. Gén. Lép. I. t. 9, f. 6. E. Doubleday, List Lep. Brit. Mus. pt. I. p. 145. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 411, n. 2.

a. ♂. N. India. Presented by Colonel Sykes.

b. ♂. N. India. From Captain Boys's Collection.

c. ♀. N. India.

Genus LIBYTHEA, Fabricius.

LIBYTHEA, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 1, 1807).

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 412.

HECAERGE, Hübner, Samml. Exot. Schmett. (1806-27).

518. *LIBYTHERA MYRRHA*, Godart.

Libythea Myrrha, Godart, *Enc. Méth.* IX. p. 171 (1819).
Hübner, *Zuträge*, f. 789, 790. *G. R. Gray*, *Lep.*
Ins. Nepal, p. 15, t. 12, f. 4. *Boisduval*, *Spéc. Gén.*
Lép. I. t. 10, f. 8. *E. Doubleday*, *List Lep. Brit.*
Mus. pt. I. p. 146. *Westwood*, in *Doubleday and*
Hewitson's Diurnal Lep. p. 413, n. 1.

- a. ♂. S. India. Presented by S. N. Ward, Esq.
- b. ♀. N. India. Presented by Colonel Buckley.
- c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

519. *LIBYTHERA LEPITA*, Moore.

Libythea Lepita, n. sp.—Differs from *Libyth. Myrrha* in the ferruginous streak from base of fore-wing being divided into two portions, the first being within the discoidal-cell, the second a round terminal spot beyond its extremity.

- a. b. ♂. N. India. Presented by Colonel Buckley.
- c. d. e. ♂ ♀. Bootan. From Pemberton's Collection.

Stirps V., with ANOPLURIFORM Larva.

ANOPLURIFORM, *Horsfield, Cat. Lep. Mus. E.I.C.* p. 50 (1828).

NATATORIAL, or ANOPLURIFORM, *Swainson, Cabinet Cyclop.* pp. 58, 60 (1840).

LEP. PLEBEII-RURALES, *pt. Linnæus.*

LEP. TETRAPODA, *pt. Latreille, Règ. Anim.* V. p. 378 (1829).

NAPAÆE-RURALES, *Hübner.*

SUCCINCTI et SUCCINCTÆ, *pt. Boisduval.*

ERYCINIDÆ, *pt. Swainson, Zool. Ill. 2nd ser.* III. t. 111 (1832-3); id. *Cabinet Cyclop.* pp. 86, 94.

ERYCINIDES, *Boisduval, Spéc. Gén. des Lép.* I. p. 164 (1836); id. *Index Méth. Eur. Lép.* p. 14.

ERYCINIDÆ, *Westwood, Introd.* II. p. 357 (1840); id. in *Doubleday and Hewitson's Diurnal Lep.* p. 415. *Stephens, Cat. Brit. Lep. Brit. Mus. pt. I.* p. 16 (1850).

ERYCINITES, *Blanchard, Hist. Nat. des Ins.* II. p. 343 (1845).

ERYCINÆ, *Swainson, Zool. Ill. 2nd ser.* II. t. 33 (1829); id. III. t. 111 (1832-3).

PAP. PLEBEII-URBICOLÆ, *Linnæus.*

PAPILIONES GENTILES ASTYCI, *Hübner.*

INVOLUTI, *Boisduval, Spéc. Gén. Lép.* I. p. 162 (1836). *Westwood, Introd. to Ent.* II. p. 347; id. *Synopsis*, p. 88.

INVOLUTÆ, *Boisduval, Ind. Méth. Eur. Lép.* p. 34.

ERYNNIS, *Schrantz, Fauna Boeia,* II. p. 157 (1801).

HESPERIDES, *Latreille, Gen. Crust. et Ins.* IV. p. 187 (1809); *Enc. Méth. IX.* p. 706. *Boisduval, Spéc. Gén. Lép.* I. p. 167.

HESPERIDÆ, *Leach. Swainson, Cabinet Cyclop.* pp. 86, 97 (1840). *Boisduval, Ind. Méth.* p. 34. *Stephens, Cat. Brit. Lep. Brit. Mus. pt. I.* p. 21. *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 505.

HESPERIIDÆ, *Stephens, Cat. Brit. Ent. Haust.* I. p. 26 (1829). *Westwood, Introd. to Ent.* II. p. 360; id. *Synopsis*, p. 88 (1840).

HESPERII, *Blanchard, Hist. Nat. des Ins.* II. pp. 322, 346 (1845).

URANIIDÆ, *pt. Westwood, Intr.* II. p. 369 (1840).

CYDIMONII, *Blanchard, Hist. Nat. des Ins.* II. pp. 322, 348 (1845). *Walker, List Lep. Brit. Mus. Het.* pt. I. p. 4 (1854).

Genus ZEMEROS, Boisduval.

ZEMEROS, *Boisduval*, Spéc. Gén. Lép. I. t. 21, f. 5 (1836). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 418.

HAMANUMIDA, pt. Hübner.

520. ZEMEROS FLEGYAS, Cramer Sp.

Papilio Flegyas, Cramer, Pap. Exot. III. t. 280, f. E.F. (1782).

Zemeros Fleygas, E. Doubleday, List Lep. Brit. Mus. pt. II. p. 1. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 419, t. 69, f. 5.

Papilio Allica, Fabricius, Ent. Syst. III. pt. I. p. 244 (1793). Donovan, Ins. of India, pl. 37, f. 2.

Hamanumida Allica, Hübner, Verz. bek. Schmett. p. 18, n. 105.

Erycina Allica, Godart, Enc. Méth. IX. p. 566.

Zemeros Allica, Boisduval, Spéc. Gén. Lép. t. 21, f. 5.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Assam. From McClelland's Collection.

Genus TAXILA, E. Doubleday.

TAXILA, E. Doubleday, List Lep. Brit. Mus. pt. II. p. 2 (1847). Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 421.

EMESIS, Horsfield. Boisduval (nec. Fabr.)

521. TAXILA DURGA, Kollar Sp.

Melitaea Durga, Kollar, in Hügel's Kaschmir, IV. pt. II. p. 441, t. 13, f. 3, 4 (1844).

Taxila Durga, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 422.

Nemeobius? Demeter, E. Doubleday, List Lep. Brit. Mus. pt. II. p. 2 (1847).

a. b. c. ♂. Simla. Presented by Colonel Hearsey.

522. TAXILA EGEON, Boisduval Sp.

Emesis Egeon, Boisduval, MS. (1847).

Taxila Egeon, E. Doubleday, List Lep. Brit. Mus. pt. II.

p. 2 (1847). Westwood, in Doubleday and Hewitson's *Diurnal Lep.* p. 422, n. 2, t. 69, f. 2.

a. b. c. ♂. Bootan. From Pemberton's Collection.

d. ♀. Silhet. Presented by Colonel Buckley.

523. *TAXILA FATNA*, Boisduval Sp.

Emesis Fatna, *Boisduval, MS.* (1847).

Taxila Fatna, E. Doubleday, *List Lep. Brit. Mus. pt. II. p. 2 (1847).*

a. b. ♂. Bootan. From Pemberton's Collection.

c. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

524. *TAXILA ERATO*, Boisduval Sp.

Emesis Erato, *Boisduval, MS.* (1847).

Taxila Erato, E. Doubleday, *List Lep. Brit. Mus. pt. II. p. 2 (1847).*

a. b. c. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

525. *TAXILA ECHERIUS*, Stoll Sp.

Papilio Echerius, Stoll, *Suppl. Cramer's Pap. Exot. V. t. 31, f. 1, a.b.* (1791).

Taxila Echerius, E. Doubleday, *List Lep. Brit. Mus. pt. II. p. 3. Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 422, n. 3.

Erycina Echerius, Godart, *Enc. Méth. IX.* p. 566.

Hesperia Coriolanus, Fabricius, *Ent. Syst. III. pt. I. p. 284 (1793).* Jones, *Icon. VI. t. 48, f. 1.*

Erycina Coriolanus, Godart, *Enc. Méth. IX.* p. 828.

Lycæna Xenodice, Hübner, *Verz. bek. Schmett.* p. 23.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. ♀. Java. From Mr. Argent's Collection.

526. *TAXILA NEOPHRON*, Boisduval Sp.

Emesis Neophron, *Boisduval, MS.* (1847).

Taxila Neophron, E. Doubleday, *List Lep. Brit. Mus. pt. II. p. 3 (1847).*

a. ♀. Java? From Mr. Argent's Collection.

527. *TAXILA DRUPADI*, Horsfield Sp.

Emesis Drupadi, *Horsfield*, *Cat. Lep. Mus. E.I.C.* t. 2, f. 3, 3a (1828). *Boisduval*, *Spéc. Gén. Lép.* I. t. 7, f. 2.

Taxila Drupadi, *E. Doubleday*, *List Lep. Brit. Mus. pt. II.* p. 2. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 422, n. 6.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

528. *TAXILA FYLLA*, Boisduval Sp.

Emesis Fylla, *Boisduval*, *MS.* (1847).

Taxila Fylla, *E. Doubleday*, *List Lep. Brit. Mus. pt. II.* p. 2 (1847). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 422, n. 8, t. 69, f. 3.

a. b. c. ♂ ♀. Bootan. From Pemberton's Collection.

d. e. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus *GONILOBA*, *Westwood*.

GONILOBA, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 511 (1852).

EPARGYREUS, *CRETEUS*, *TALIDES*, *ASTRAPTES*, *THRACIDES*, *TELEGONUS*, et *PROTEIDES*, *pt. Hübner*.

EUDAMUS, *pt. Boisduval*.

529. *GONILOBA CHROMUS*, Cramer Sp.

Papilio Chromus, *Cramer*, *Pap. Exot. III.* t. 284, f. E. (1782).

Goniloba Chromus, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 513, n. 32.

Hesperia Chromus, *Latrelle*, *Enc. Méth. IX.* p. 744.

a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

d. N. India. Presented by Colonel Hearsey.

e. f. Borneo. Presented by Edwin Shepherd, Esq.

530. *GONILOBA SENA*, Moore.

Goniloba Sena, Moore, Proc. Zool. Soc. (ined.).

- a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.
- d. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

531. *GONILOBA DERMA*, Moore.

Goniloba Derma, Moore, Proc. Zool. Soc. (ined.).

- a. ♂. Java. From Dr. Horsfield's Collection.

532. *GONILOBA BADRA*, Moore.

Goniloba Badra, Moore, Proc. Zool. Soc. (ined.).

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The transformations of *Goniloba Badra* are figured on Plate VII., fig. 3, 3a, from Java. "Feeds upon a leguminous plant bearing the native name of *Tungkul*. March."—(Horsfield, MS.)

The transformations of this species were also observed by Mrs. Hamilton, and are figured among her original drawings, now in the possession of the Entomological Society of London.

533. *GONILOBA JAPETUS*, Cramer Sp.

Papilio Japetus, Cramer, Pap. Exot. IV. t. 365, f. E.F. (1782). Drury, Ins. III. t. 17, f. 1, 2.

Hesperia Nepos, Fabricius, Ent. Syst. III. pt. I. p. 340 (1793). Jones, Icon. VI. t. 76, f. 2. Latreille, Enc. Méth. IX. p. 787. Donovan's Drawings in Bibl. Hope, Oxford.

Goniloba Nepos, Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 513, n. 56.

- a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Goniloba Japetus* are figured on Plate VII., fig. 2, 2a, from Java. "Feeds upon the Uwi-leggi, *Dioscorea oppositifolia*. April, May. Rather scarce."—(Horsfield, MS.)

534. *GONILOBA GANA*, Moore.

Goniloba Gana, Moore, Proc. Zool. Soc. (ined.).

- a. b. c. ♂ ♀. Java. From Dr. Horsfield's Collection.

535. *GONILOBA RAVI*, Moore.

Goniloba Ravi, *Moore, Proc. Zool. Soc. (ined.)*.

a. Pinang. Presented by Dr. Cantor.

536. *GONILOBA TAPANA*, Moore.

Goniloba Tapana, *Moore, Proc. Zool. Soc. (ined.)*.

a. Borneo. Presented by Edwin Shepherd, Esq.

537. *GONILOBA MENAKA*, Moore.

Goniloba Menaka, *Moore, Proc. Zool. Soc. (ined.)*.

? Hesperia Eson, *Boisduval, MS.*

a. b. c. d. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

538. *GONILOBA GOPALA*, Moore.

Goniloba Gopala, *Moore, Proc. Zool. Soc. (ined.)*.

a. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

539. *GONILOBA MADHAVA*, Moore.

Goniloba Madhava, *Moore, Proc. Zool. Soc. (ined.)*.

a. Darjeeling. From Messrs. Schlagintweit's Collection.

540. *GONILOBA PRALAYA*, Moore.

Goniloba Pralaya, *Moore, Proc. Zool. Soc. (ined.)*.

a. b. Java. From Dr. Horsfield's Collection.

541. *GONILOBA BHAGAVA*, Moore.

Goniloba Bhagava, *Moore, Proc. Zool. Soc. (ined.)*.

a. N. India. Presented by Colonel Buckley.

542. *GONILOBA SAMBARA*, Moore.

Goniloba Sambara, *Moore, Proc. Zool. Soc. (ined.)*.

a. b. c. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus ISMENE, Swainson.

Ismene, Swainson, Zool. Ill. 1st ser. t. 16 (1820-1).

Thymele, Boisduval, Fauna Madagascar.

Castnius, pt. Hübner.

543. *ISMENE OEDIPODEA*, Swainson.

♂ ♀ Ismene Oedipodea, Swainson, Zool. Ill. 1st ser. t. 16 (1820-21). Thon, Ent. Archiv. p. 91, pl. 4, fig. 5. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 515, n. 3.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Bootan. From Pemberton's Collection.

f. ♂. Silhet. Presented by Colonel Buckley.

g. ♀. Canara, S. India. Presented by S. N. Ward, Esq.

The larva and pupa of *Ismene Oedipodea* are figured on Plate VII., fig. 4, 4a, from Java. "Feeds upon *Gärtnera Javensis*, which bears the native name of *Kakkas*. August."—(Horsfield, MS.)

544. *ISMENE JAINA*, Moore.

Ismene Jaina, Moore, Proc. Zool. Soc. (ined.).

a. ♂. Darjeeling. From Messrs. Schlagintweit's Collection.

545. *ISMENE HARISA*, Moore.

Ismene Harisa, Moore, Proc. Zool. Soc. (ined.).

a. b. c. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

546. *ISMENE AMARA*, Moore.

Ismene Amara, Moore, Proc. Zool. Soc. (ined.).

a. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

547. *ISMENE VASUTANA*, Moore.

Ismene Vasutana, Moore, Proc. Zool. Soc. (ined.).

a. Darjeeling. From Pearson's Collection.

b. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

548. *ISMENE BENJAMINI*, Guérin-Méneville Sp.

Hesperia (Thymele) Benjaminii, *Guérin-Méneville*, in
Delessert's Voy. dans l'Inde, pt. 2, p. 79, t. 22,
f. 2, 2a (1843).

Ismene Benjaminii, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 515, n. 4.

Hesperia Xanthopogon, *Kollar*, in *Hügel's Kaschmir*,
 IV. pt. II. p. 453, t. 18, f. 1, 2 (1844).

a. N. India. Presented by Colonel Buckley.

b. Darjeeling. From Indian Collection, Exposition
 Universelle at Paris, 1855.

549. *ISMENE JAYADEVA*, Moore.

Ismene Jayadeva, *Moore*, *Proc. Zool. Soc.* (ined.).

a. Darjeeling. From Indian Collection, Exposition
 Universelle at Paris, 1855.

550. *ISMENE SINTA*, Moore.

Ismene Sinta, *Moore*, *Proc. Zool. Soc.* (ined.).

a. Java. From Dr. Horsfield's Collection.

551. *ISMENE DASA*, Moore.

Ismene Dasa, *Moore*, *Proc. Zool. Soc.* (ined.).

a. Java. From Dr. Horsfield's Collection.

552. *ISMENE LADON*, Cramer Sp.

Papilio Ladon, *Cramer*, *Pap. Exot.* III. t. 284, f. G.
 (1782).

Ismene Ladon, *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 515, n. 11.

Hesperia Ladon, *Latreille*, *Enc. Méth.* IX. p. 749.

a. b. Java. From Dr. Horsfield's Collection.

c. d. N. India. From the Asiatic Society of Bengal.

e. N. India. Presented by Colonel Hearsey.

553. *ISMENE AMBASA*, Moore.

Ismene Ambasa, *Moore*, *Proc. Zool. Soc.* (ined.).

a. b. ♂. Java. From Dr. Horsfield's Collection.

c. ♀. N. India. Presented by Lieutenant Hugo James.

The larva and pupa of *Ismene Ambasa* are figured on Plate VII., fig. 6, 6a, from Java. "Feeds upon a species of Pandanus, bearing the native name of *Pandan-duri*. May. Rather scarce."—(Horsfield, MS.)

554. ISMENE CHAON, Boisduval.

Ismene Chaon, Boisduval, MS.

a. b. c. d. Java. From Dr. Horsfield's Collection.

e. Pinang. Presented by Dr. Cantor.

555. ISMENE BATARA, Moore.

Ismene Batara, Moore, Proc. Zool. Soc. (ined.).

a. b. c. Java. From Dr. Horsfield's Collection.

Genus PYRGUS, Hübner.

Pyrgus, Hübner. Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 516.

Syrichtus, Boisduval.

556. PYRGUS AGAMA, Moore.

Pyrgus Agama, Moore, Proc. Zool. Soc. (ined.).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Pyrgus Agama* are figured on Plate VII., fig. 1, 1a, from Java. "Feeds upon a species of Microcos, bearing the native name of *Talok*. December to February. Rather common."—(Horsfield, MS.)

557. PYRGUS SUPERNA, Moore.

Pyrgus Superna, Moore, Proc. Zool. Soc. (ined.).

a. b. N. India. From the Asiatic Society of Bengal.

c. d. N. India. Presented by Colonel Hearsey.

558. PYRGUS DANNA, Moore.

Pyrgus Danna, Moore, Proc. Zool. Soc. (ined.).

a. b. c. Bootan. From Pemberton's Collection.

559. *PYRGUS? PURENDRA*, Moore.

Pyrgus Purendra, Moore, Proc. Zool. Soc. (ined.).

a. Canara, S. India. Presented by S. N. Ward, Esq.

Genus *NISONIADES*, Hübner.

NISONIADES, Hübner.

THANAO, Boisduval.

560. *NISONIADES SALSALA*, Moore.

Nisoniades Salsala, Moore, Proc. Zool. Soc. (ined.).

a. b. c. Canara (S. India). Presented by S. N. Ward, Esq.

561. *NISONIADES DAHA*, Moore.

Nisoniades Daha, Moore, Proc. Zool. Soc. (ined.).

a. b. c. Java. From Dr. Horsfield's Collection.

562. *NISONIADES DIOCLES*, Boisduval.

Thanaos Diocles, Boisduval, MS.

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

563. *NISONIADES? AMERTA*, Moore.

Nisoniades? Amerta, Moore, Proc. Zool. Soc. (ined.).

a. Java. From Dr. Horsfield's Collection.

b. N. India. Presented by Colonel Buckley.

Genus *PAMPHILA*, Fabricius.

PAMPHILA, Fabricius, Syst. Gloss. (Illiger's Mag. VI. p. 287, 1807).

Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 521.

CÆLENORRHINUS, CYCLOPIDES, STEROPES, AUGIADES, et THYMELICUS, pt. Hübner.

HESPERIA, Boisduval.

564. *PAMPHILA VEDANGA*, Moore.

Pamphila Vedanga, Moore, Proc. Zool. Soc. (ined.).

a. b. c. ♂. Java. From Dr. Horsfield's Collection.

d. ♂. Bootan. From Pemberton's Collection.

- e. ♂. N. India. Presented by Colonel Hearsey.
 f. ♂. Canara, S. India. Presented by S. N. Ward, Esq.
 g. h. ♂ ♀. N. India. Presented by Colonel Buckley.

565. *PAMPHILA NARANATA*, Moore.

Pamphila Naranata, Moore, Proc. Zool. Soc. (ined.).

- a. Java. From Dr. Horsfield's Collection.

566. *PAMPHILA AUGIAS*, Linnæus Sp.

Papilio Augias, Linnæus, Syst. Nat. I. II. p. 794. *Fabricius*, Mant. Ins. II. p. 84. *Donovan*, Ins. of India, t. 48, f. 1.

Pamphila Augias, Westwood, in *Doubleday and Hewitson's Diurnal Lep.* p. 522, n. 14.

Hesperia Augias, Latreille, Enc. Méth. IX. p. 767.

♂ *Pamphila Pheres*, Boisduval, MS.

- a. b. c. d. e. f. g. h. ♂ ♀. Java. From Dr. Horsfield's Collection.

567. *PAMPHILA SUMBADRA*, Moore.

Pamphila Sumbadra, Moore, Proc. Zool. Soc. (ined.).

- a. b. Java. From Dr. Horsfield's Collection.

Genus *ACHLYODES*, Hübner.

Achlyodes, Hübner, Zuträge.

Eantis, Boisduval.

568. *ACHLYODES?* *CHANDRASA*, Moore.

Achlyodes? *Chandrasa*, Moore, Proc. Zool. Soc. (ined.).

- a. Java. From Dr. Horsfield's Collection.

569. *ACHLYODES?* *SURA*, Moore.

Achlyodes? *Sura*, Moore, Proc. Zool. Soc. (ined.).

- a. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

570. *ACHLYODES?* *VASAVA*, Moore.

Achlyodes? *Vasava*, *Moore*, *Proc. Zool. Soc. (ined.)*.

- a. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

Genus *HESPERIA*, *Fabricius*.

HESPERIA, *Fabricius*. *Swainson*, *Zool. Ill. 1st ser.* *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 525.

571. *HESPERIA FOLUS*, Cramer Sp.

Papilio Folus, *Cramer*, *Pap. Exot. I. t. 74, f. E.* (1779).

Hesperia Cicero, *Fabricius*, *Ent. Syst. III. pt. I. p. 358* (1793). *Latreille*, *Enc. Méth. IX.* p. 787. *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 526, n. 16.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. N. India. From the Asiatic Society of Bengal.

f. S. India. Presented by S. N. Ward, Esq.

The larva and pupa of *Hesperia Folus* are figured on Plate VII., fig. 7, 7a, copied from the original drawings in the possession of A. Grote, Esq., of Calcutta.

572. *HESPERIA FATIH*, Kollar.

Hesperia Fatih, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 454, t. 18, f. 5, 6 (1844). *Westwood*, in *Doubleday and Hewitson's Diurnal Lep.* p. 526, n. 19.

a. N. India. Presented by Colonel Buckley.

b. c. d. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

573. *HESPERIA PULOMAYA*, Moore.

Hesperia Pulomaya, *Moore*, *Proc. Zool. Soc. (ined.)*.

a. b. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

574. *HESPERIA LEUCOCERCA*, Kollar.

Hesperia leucocerca, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 454, t. 18, f. 3, 4 (1844). *Westwood*, in

Doubleday and Hewitson's Diurnal Lep. p. 526,
n. 20.

a. Bootan. From Pemberton's Collection.

575. *HESPERIA PUTRA*, Moore.

Hesperia Putra, Moore, *Proc. Zool. Soc.* (*ined.*).

a. b. c. Java. From Dr. Horsfield's Collection.

576. *HESPERIA ASMARA*, Moore.

Hesperia Asmara, Moore, *Proc. Zool. Soc.* (*ined.*).

a. Java. From Dr. Horsfield's Collection.

577. *HESPERIA ALYSOS*, Boisduval.

Hesperia Alykos, Boisduval, MS.

a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

578. *HESPERIA PRABA*, Moore.

Hesperia Praba, Moore, *Proc. Zool. Soc.* (*ined.*).

a. Java. From Dr. Horsfield's Collection.

579. *HESPERIA INDRANI*, Moore.

Hesperia Indrani, Moore, *Proc. Zool. Soc.* (*ined.*).

a. ♂. N. India. Presented by Colonel Hearsey.

b. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

580. *HESPERIA DAN*, Fabricius.

Hesperia Dan, *Fabricis, Mant. Ins.* II. p. 88 (1787);
Ent. Syst. III. I. p. 341.

Goniloba Dan, *Westwood, in Doubleday and Hewitson's Diurnal Lep.* p. 513, n. 52.

Hesperia Eacus, *Latreille, Enc. Méth.* IX. p. 738 (1819).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

581. *HESPERIA SEMA*, Moore.

Hesperia Sema, Moore, *Proc. Zool. Soc.* (*ined.*).

a. Java. From Dr. Horsfield's Collection.

582. *HESPERIA THRAX*, *Linnæus Sp.*

- Papilio Thrax, *Linnæus, Syst. Nat.* I. II. p. 794 (1767).
Fabricius, Mant. Ins. II. p. 87? *Donovan, Ins. of India*, t. 49, 2.
Hesperia Thrax, *Latreille, Enc. Méth.* IX. p. 748.
Westwood, in Doubleday and Hewitson's Diurnal Lep. p. 527, n. 38.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. f. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

The larva and pupa of *Hesperia Thrax* are figured on Plate VII., fig. 5, 5a, from Java. "Feeds on the Pisang (*Mura paradisiaca*). November to February. Common."—(Horsfield, MS.)

583. *HESPERIA IRAVA*, *Moore*.

- Hesperia Irava*, *Moore, Proc. Zool. Soc.* (ined.).
Cælænorrhinus Thrax, *Hübner, Zuträge*, f. 875-6.
a. b. c. d. Java. From Dr. Horsfield's Collection.

584. *HESPERIA PANDIA*, *Moore*.

- Hesperia Pandia*, *Moore, Proc. Zool. Soc.* (ined.).
a. Java. From Dr. Horsfield's Collection.
b. Canara (S. India). Presented by S. N. Ward, Esq.

The larva and pupa of *Hesperia Pandia* are figured on Plate VII., fig. 10, 10a, copied from the original drawings made by A. Grote, Esq.

585. *HESPERIA ESA*, *Moore*.

- Hesperia Esa*, *Moore, Proc. Zool. Soc.* (ined.).
a. Java. From Dr. Horsfield's Collection.

586. *HESPERIA DRUNA*, *Moore*.

- Hesperia Druna*, *Moore, Proc. Zool. Soc.* (ined.).
a. Java. From Dr. Horsfield's Collection.

587. *HESPERIA ARIA*, *Moore*.

- Hesperia Aria*, *Moore, Proc. Zool. Soc.* (ined.).
a. b. Java. From Dr. Horsfield's Collection.

588. *HESPERIA CHAYA*, Moore.

Hesperia Chaya, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. b. Java. From Dr. Horsfield's Collection.
- c. Pinang. Presented by Dr. Cantor.
- d. e. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- f. N. India. Presented by Colonel Hearsey.

589. *HESPERIA AGNA*, Moore.

Hesperia Agna, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. b. c. Java. From Dr. Horsfield's Collection.
- d. e. f. g. Canara, S. India. Presented by S. N. Ward, Esq.

590. *HESPERIA MANGALA*, Moore.

Hesperia Mangala, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. b. Pinang. Presented by Dr. Cantor.
- c. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

591. *HESPERIA CINNARA*, Moore.

Hesperia Cinnara, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. b. Java. From Dr. Horsfield's Collection.

592. *HESPERIA KUMARA*, Moore.

Hesperia Kumara, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. Canara, S. India. Presented by S. N. Ward, Esq.

593. *HESPERIA SUTAPA*, Moore.

Hesperia Sutapa, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. Java. From Dr. Horsfield's Collection.

594. *HESPERIA DIVODASA*, Moore.

Hesperia Divodasa, Moore, *Proc. Zool. Soc.* (*ined.*).

- a. Canara, S. India. Presented by S. N. Ward, Esq.

Remark.—The above enumerated new species of *Hesperia* form the subjects of a memoir which is now in course of preparation to be submitted to the Zoological Society of London.

Genus NYCTALEMON, *Dalman.*

NYCTALEMON, *Dalman*, *Acta Holm.* (1824), p. 407. *Westwood*.

Walker, *List Lep. Brit. Mus. Het. pt. I.* p. 7.

ORONTES, *Swainson*, *Zool. Ill. 2nd ser. pl. 125* ().

URANIA, *pt. Fabricius*. *Latreille*.

595. *NYCTALEMON PATROCLUS*, *Linnaeus Sp.*

Papilio Patroclus, *Linnaeus*, *Mus. Lud. Ulr.* p. 204
 (); *Syst. Nat.* II. p. 749. *Clerck*, *Icon. pl. 37*,
f. 1. *Fabricius*, *Spec. Ins.* II. p. 13; *Mant. Ins.* II.
 p. 7; *Ent. Syst.* III. p. 2. *Seba*, *Mus.* IV. *pl. 47*,
f. 13, 16. *D'Aubent*, *Miscell.* *pl. 17*, *f. 1, 2*.
Olivier, *Enc. Méth. Ins. Pap.* *pl. 8, f. 1*. *Cramer*,
Pap. Exot. III. p. 9, *pl. 198, f. A.* *pl. 109, f. A.B.*
Herbst, *Pap.* *pl. 54, f. 2, 3, pl. 55, f. 1*.

Nyctalemon Patroclus, *Westwood*. *Drury*, *Ins.* I. p. 13,
pl. 7, f. 1, pl. 8, f. 1. *Walker*, *List Lep. Brit. Mus. Het. pt. I.* p. 8.

Noctua Patroclus, *Fabricius*, *Ent. Syst.* III. 2, 8, 2.

Urania Patroclus, *Latreille*. *Latreille et Godart*, *Enc. Méth.* IX. p. 710. *Lamarek*, *Hist. Nat. Anim. Invert.* IV. p. 194. *Boisduval*, *Voy. de l'Astrolabe*,
pt. I. Lép. p. 254.

Lyssa Achillaria et L. Patroclaria, *Hübner*, *Verz. bek. Schmett.* p. 289.

a. Silhet. From Mr. Argent's Collection.

"This insect is common in Collections from China, Assam, Silhet, and Arakan."—(E. Blyth, J. A. S. Beng. (1854), p. 217.)

Tribe II. SPHINGES.

SPHINX, *pt. Linnæus* (*Sphinges legitimæ*, sec. *, **, *** *pt.*).

LEP. CREPUSCULARIA, *Latreille*, *Gén. Crust. et Ins.* IV. pp. 187, 209 (1809). *Stephens*, *Cat. Brit. Lep. Brit. Mus.* *pt. I.* p. 24 (1850).

FUSICORNES seu CLOSTEROERES, *pt. Dumeril*, *Consid. Gén. des Ins.* pp. 139, 163, 322 (1823).

HETEROCERA, *pt. Boisduval et Leconte*, *Icon. Lép.* I. p. 39 (1836). *Boisduval*, *Ind. Méth.* p. 39 (1840). *Westwood*, *Introd.* II. p. 363 (1840). *Stephens*, *Cat. Lep. Brit. Mus.* *pt. I.* p. 24 (1850). *Walker*, *Cat. Lep. Brit. Mus. Het.* *pt. I.* p. 1 (1854). *Stainton*, *Insecta Brit. Lep. Tineina*, p. 2 (1855); id. *Manual Brit. Butt.* p. 72 (1856).

CHALINOPTERA, *pt. Blanchard*, *Hist. Nat. des Ins.* II. pp. 322, 349. *Chenu*, *Enc. d'Hist. Nat. Pap.* 232.

SPH. LEGITIMÆ, INTEGRALES, et DENTALES, *Haworth*, *Lep. Brit.* p. 55 (1803).

SPHINGII, *pt. Blanchard*, *Hist. Nat. des Ins.* II. pp. 323, 356. *Chenu*, *Enc. d'Hist. Nat. Pap.* p. 252. *Walker*, *Cat. Lep. Brit. Mus. Het.* *pt. I.* p. 2 (1854); id. *pt. VIII.* p. 76.

SPHINGIDÆ, *Leach*, *Edin. Encycl.* IX. p. (1815). *Horsfield*, *Cat. Lep. Mus. E.I.C.* pp. 20, 22 (1828). *Stephens*, *Ill. I.* p. 110 (1828); id. *Cat. Brit. Lep. Brit. Mus.* *pt. I.* p. 25 (1850). *Westwood*, *Introd.* II. p. 364 (1840). *Walker*, *Cat. Lep. Brit. Mus. Het.* *pt. I.* p. 76 (1856). *Stainton*, *Manual Brit. Butt.* pp. 75, 81 (1856).

SPHINGIDES, *Latreille*, *Gén. Crust. et Ins.* IV. pp. 188, 209.

SPHINGES, *pt. Hübner*, *Zuträge*, p. 4 (1818).

SPHINGIDES, *Boisduval*, *Ind. Méth. Lép. Eur.* p. 45 (1840).

SPHINGIDÆ, *pt. Swainson*, *Cab. Cycl.* p. 99 (1840).

SPHINGINA, *Stainton*, *Insecta Brit. Lep. Tineina*, p. 2 (1855); id. *Manual Brit. Butt.* pp. 72, 75 (1856).

CASTNII et CASTNITES, *pt. Blanchard*. *Walker*.

The SPHINGES constitute in this Catalogue a distinct Tribe, equal in rank and value to the PAPILIONES, or to the other Tribes which will be subsequently defined, although more limited in the number of individuals.

The metamorphosis varies in the different Stirpes. The chrysalis is generally naked, of a cylindrico-conical form, deposited on the surface, under the earth, or rarely enveloped in a slight web.

The antennæ are prismatic, somewhat fusiform, or thickened in the middle and attenuated at the ends. The individuals are provided with a proboscis, varying in size and length in the different Stirpes.

This Tribe is subdivided into five distinct Stirpes, four of which derive their name from the form of the Caterpillar.

The first Stirps forms a natural junction between the Hesperidæ, the last subdivision of the Anopluriform Stirps of the PAPILIONES, and the second Stirps of the SPHINGES. The typical genera are *Castnia* and *Euschemon*. Their reciprocal affinity manifests itself in the following characters:—*First*, in the form of the antennæ: these organs in *Castnia* resemble those of *Hesperia* in the length of the club and its attenuation towards the base, while they approach to *Sphinx* in the prismatic form of the club and its terminal hook. *Secondly*, the fore-legs are perfect in both sexes of *Hesperia* and *Castnia*.

The habit and aspect are very similar in both groups. In the Systems of Lamarck and Latreille, *Castnia* and *Sphinx* stand contiguous.

Mr. Swainson, who observed *Castnia* in its native country, states that “they fly by day, and much in the same manner as the typical *Hesperidæ*. The antennæ are intermediate in form between a *Sphinx* and a *Hesperia*.”—(Swainson and Shuckard’s Treatise on Insects, p. 104.)

The metamorphosis is as yet a desideratum in Entomology.

The second Stirps, named *Elongatæ* or *Langleib-raupen*, Wien. Verz. and Schrank, has a cylindrical larva, more slender and elongated than in the other Stirpes. It is generally without lateral ocelli; the head is globular, and the abdominal horn short and rigid. The perfect insect has short wings, and flies by day, and the abdomen is terminated by a broad or depressed brush-like appendage. The proboscis is long. In their metamorphosis they exhibit a slight affinity to the Diurna. The typical genera are *Sesia* and *Macroglossa*. The metamorphosis is illustrated on Plate VIII., in figures 1, 2, 3, and 4.

The *third* Stirps, named *Acrocephalæ* or *Spitzohopf-raupen*, Wien. Verz. and Schrank, has a larva with a head almost triangular or acuminate above, the body being obliquely striated, generally with yellow,—naked, and somewhat rugose; the abdominal horn of moderate size, smooth, and marked with a peculiar colour. The perfect insect is strikingly distinguished by its angulated or excavated wings. The proboscis is very short or obsolete. The typical genus is *Smerinthus*. Of this Stirps no transformations are contained in the Company's Museum.

The *fourth* Stirps, named *Amblocephalæ* or *Stumpfkopf-raupen*—obtuse-headed larva,—has a larva with an ovate truncated head; it is nearly naked and even on the surface; the abdominal horn is lengthened, tuberculated, and curved. The perfect insect has entire lanceolate wings; the abdomen is marked with oblique lateral striae. The proboscis is large and strong, and this character is strikingly manifested in the pupa (see Plate IX., fig. 3a). The typical genera are *Acherontia* and *Sphinx*. The metamorphosis is illustrated on Plate VIII., fig. 6, and Plate IX., figs. 1, 2, 3, and 4.

The *fifth* Stirps, characteristically named in the Wiener Verzeichnis *Augen-raupen*, larvae *Ophthalmicæ*, has a larva with a small globose retractile head, and is moreover strikingly distinguished by the ocelli with which it is marked, and which are also observed on the chrysalis. These ocelli, or eyes, are placed in some genera behind the head only, while in others they exist in a regular row along the entire larva, and in some cases are very prominent and brilliant, as appears in the figures. The metamorphosis takes place on the surface of the earth, the pupa being in some cases covered with a slight web. The pupa is enlarged towards the head, and is generally provided with an annular case, covering the proboscis. The proboscis is not so large as in the preceding Stirps. In the perfect insect the abdomen is long and acute, and the individuals in many cases have the habit of a *Vespertilio*.

The typical genera are *Deilephila* and *Chærocampa*. The metamorphoses are illustrated on Plate IX., fig. 5, and Plates X. and XI.

Stirps II. Larvæ ELONGATÆ.

SESIÆ, sect. TECTÆ et (DENUDATÆ, div. I.), Haworth, Lep. Brit. pp. 55, 65 (1803).

MACROGLOSSIADÆ, Harris.

SESIARLÆ, pt. Boisduval, Ind. Méth. Lép. Eur. p. 41 (1840).

SESIIDÆ, Stephens, Ill. I. p. 132 (1828); id. Cat. Brit. Lep. Brit. Mus. pt. I. p. 29 (1850). Stainton, Manual Brit. Butt. pp. 75, 97 (1856).

SESIIDES, pt. Blanchard, Hist. Nat. des Ins. II. p. 352 (1845).

Genus SESIA, Fabricius.

SESIÆ, Fabricius, Ent. Syst. III. I. p. 379 (1775). Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 79 (1856).

SPHINX, pt. Linnæus.

HEMARIS, Dalman, Sp. S. p. 215 ().

MACROGLOSSA, pt. Ochsenheimer. Boisduval.

CEPHONODES, pt. Hübner, Verz. bek. Schmett. p. 131.

596. SESIA HYLAS, Linnæus Sp.

Sphinx Hylas, Linnæus, Mant. I. p. 539 (). *Fabricius, Spec. Ins. II. p. 154; id. Mant. Ins. II. p. 99. Gmelin, Syst. Nat. I. 5, 2387, 95.*

Sesia Hylas, Fabricius, Ent. Syst. III. pt. I. p. 379. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 84.

Cephonodes Hylas, Hübner, Verz. bek. Schmett. p. 131.

Sphinx Picus, Cramer, Pap. Exot. II. p. 83, pl. 148 f. B. (1779).

Macroglossa Picus, Kollar, in Hügel's Kaschmir, IV. pt. II. p. 458.

- a. N. India. Presented by Colonel Sykes.
- b. N. India. Presented by Colonel Hearsey.
- c. Canara, S. India. Presented by S. N. Ward, Esq.
- d. e. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.
- f. Pinang. Presented by Dr. Cantor.

The larva and pupa of *Sesia Hylas* are figured on Plate VIII., fig. 1, 1 α , copied from General Hardwicke's original drawings, fig. 179.

Genus SATASPES, *Moore.*

MACROGLOSSA (Group 3), *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 95.*

Differs from *Macroglossa* in having the antennæ stout and strongly serrate-setose; abdomen broad to the tip; the fore-wings very narrow and pointed, and the exterior border extremely oblique.

597. *SATASPES INFERNALIS*, *Westwood Sp.*

Sesia infernalis, *Westwood, Cabinet Orient. Ent. p. 61, pl. 30, f. 3 (1847).*

Macroglossa infernalis, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 95.*

a. Silhet. Presented by the Trustees of the British Museum.

Genus MACROGLOSSA, *Ochsenheimer.*

MACROGLOSSUM, *Scopoli, Ent. Carn. p. 414 (1763). Swainson, Zool. Ill. 1st ser. t. 64.*

MACROGLOSSA, *Ochsenheimer, Schmett. von Eur. IV. p. 41 (1816). Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 85.*

SPHINX, *pt. Linnaeus.*

SESIAS, *pt. Fabricius, Ent. Syst.*

PSITHYROS et CEPHONODES, *pt. Hübner.*

598. *MACROGLOSSA STELLATARUM*, *Linnaeus Sp.*

Sphinx Stellatarum, Linnaeus, Syst. Nat. I. p. 493 (1758).

Macroglossa Stellatarum, Ochsenheimer, Schmett. von Eur. IV. p. 42. Stephens, Ill. Brit. Ent. Haust. I. p. 133; Cat. Brit. Lep. Brit. Mus. pt. I. p. 29. Boisduval, Ind. Méth. p. 45. Curtis, Brit. Ent. XVI. pl. 747. Stainton, Manual Brit. Lep. I. p. 98. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 86.

Sesia Stellatarum, Fabricius, Ent. Syst. III. pt. I. p. 380.

Psithyros Stellatarum, Hübner, Verz. bek. Schmett. p. 132.

a. China. Presented by Dr. Cantor.

599. *MACROGLOSSA CORYTHUS*, Boisduval.

Macroglossa Corythus, *Boisduval*, *MS.* *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* *p. 92* (1856).

Macroglossum arcuatum, *Horsfield*, *MS.* (1826).

a. b. c. d. Java. From Dr. Horsfield's Collection.

e. Canara. Presented by S. N. Ward, Esq.

f. g. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

h. Ladakh. Presented by Captain R. Strachey.

The larva and pupa of *Macrog. Corythus* are figured on Plate VIII., fig. 2, 2*a*, from Java.

600. *MACROGLOSSA GILIA*, Boisduval.

Macroglossa Gilia, *Boisduval*, *MS.* *Herrich-Schaeffer*, *Lep. Exot. series I. pt. 6, f. 107* (1854). *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* *p. 93* (1856).

Macroglossum maculatum, *Horsfield*, *MS.* (1826).

a. b. c. d. e. f. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Macrog. Gilia* are figured on Plate VIII., fig. 3, 3*a*, from Java. "Feeds on the Patji or Bengkudu (*Morinda citrifolia*). January and February. Tolerably common."—(Horsfield, MS. Notes.)

601. *MACROGLOSSA GYRANS*, Boisduval.

Macroglossa gyrans, *Boisduval*, *MS.* *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* *p. 91* (1856).

a. b. N. India. Presented by Colonel Sykes.

c. d. N. India. Presented by Colonel Hearsey.

602. *MACROGLOSSA SITIENE*, Boisduval.

Macroglossa Sitiene, *Boisduval*, *MS.* *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* *p. 92* (1856).

a. N. India. Presented by Colonel Buckley.

603. *MACROGLOSSA PASSALUS*, Drury Sp.

Sphinx Passalus, *Drury*, *Ins. II.* *p. 52, pl. 29, f. 2* (1773).

Macroglossa Passalus, *Westwood*, in 2nd ed. *Drury's Ins. Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 92.
Sphinx Belis, *Cramer*, *Pap. Exot. II.* p. 147, pl. 94, f. C. (1782).

Macroglossum Rhebus, *De Cerisy*, *MS.* (1826).

- a. Java. (*Mac. Rhebus*, *De Cerisy*). From Dr. Horsfield's Collection.
- b. c. N. India. Presented by Colonel Sykes.
- d. e. N. India. Presented by Colonel Hearsey.
- f. Canara, S. India. Presented by S. N. Ward, Esq.

604. *MACROGLOSSA DIVERGENS*, *Walker*.

Macroglossa divergens, *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 94 (1856).

- a. b. Canara, S. India. Presented by S. N. Ward, Esq.

605. *MACROGLOSSA NYCTERIS*, *Kollar*.

Macroglossa Nycteris, *Kollar*, in *Hügel's Kaschmir*, IV. pt. II. p. 458, pl. XIX. f. 5 (1844).

Macroglossa volucris, *Boisduval*, *MS. Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 94 (1856).

- a. N. India. From Captain Boys's Collection.

606. *MACROGLOSSA FARO*, *Cramer Sp.*

Sphinx Faro, *Cramer*, *Pap. Exot. III.* p. 165, pl. 285 f. C. (1782).

- a. Java. From Dr. Horsfield's Collection.

Remark.—This is a very distinct and well-defined species, and is correctly figured by Cramer.

Genus *LOPHURA*, *Boisduval*.

LOPHURA, *Boisduval*, *MS. Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 105 (1856).

ENYO, *pt. Hübner*, *Verz. bek. Schmett.*

607. *LOPHURA HYAS*, *Boisduval*.

Lophura Hyas, *Boisduval*, *MS. Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 107 (1856).

Macroglossum geometricum, *Horsfield*, *MS.* (1826).

a. b. c. d. e. f. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Lophura Hyas* are figured on Plate VIII., fig. 4, 4a, from Java. "Feeds on a species of Pæderia, bearing the native name of *Simbuæn*."—(Horsfield, MS.)

Stirps III. Larvæ ACORCEPHALÆ.

Genus SMERINTHUS, Latreille.

SMERINTHUS, Latreille, *Hist. Nat. Ins.* III. p. 401 (1802). Walker, *List Lep. Brit. Mus. Het. pt.* VIII. p. 239.

Sphinx, pt. Linnæus.

MERINTHUS, Meigen.

PAONIAS, MIMAS, et POLYPTYCHUS, Hübner, Verz. bek. Schmett.

608. SMERINTHUS DRYAS, Boisduval.

Smerinthus Dryas, Boisduval, MS. Walker, *List Lep. Brit. Mus. Het. pt.* VIII. p. 250 (1856).

Smerinthus parallelis, Horsfield, MS. (1826).

Smerinthus Horsfieldii, De Cerisy, MS. (1826).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

609. SMERINTHUS DENTATUS, Cramer Sp.

Sphinx dentatus, Cramer, *Pap. Exot.* II. p. 42, pl. 125,

f. G. (1779). Fabricius, *Spec. Ins.* II. p. 43, n. 16;

Mant. Ins. II. p. 94, n. 18; *Ent. Syst.* III. I. p. 361, n. 19.

Smerinthus dentatus, Walker, *List Lep. Brit. Mus. Het. pt.* VIII. p. 252.

Polyptychus dentatus, Hübner, Verz. bek. Schmett. p. 141.

Sphinx Timesius, Stoll, *Suppl. Cramer, Pap. Exot.* V. p. 172, t. 40, f. 1.

a. ♂. N. India. Presented by Colonel Hearsey.

b. ♀. N. India. From Captain Boys's Collection.

Stirps IV. Larvæ AMBLOCEPHALÆ.

Genus LEUCOPHLEBIA, *Westwood.*

LEUCOPHLEBIA, *Westwood*, *Cabinet Orient. Ent.* p. 46 (1847).
Walker, *List Lep. Brit. Mus. Het. pt. IX.* p. 136.

610. LEUCOPHLEBIA LINEATA, *Westwood.*

Leucophlebia lineata, *Westwood*, *Cabinet Orient. Ent.*
 p. 46, pl. 22 (1847). *Walker*, *List Lep. Brit. Mus.*
Het. pt. IX. p. 136.

a. Java. From Dr. Horsfield's Collection.

b. N. India. From the Asiatic Society of Bengal.

The larva of this interesting genus is figured on Plate VIII., fig. 5, copied from the original drawing made by — Frith, Esq. "Feeds on the Ootoo, or common thatch-grass (*Saccharum cylindricum*). Rather common."—(Frith's MS. Notes.)

Remark.—This genus is arranged by Mr. F. Walker, in his tribe NOCTUITES, at the end of the family Leucanidæ, who remarks, that it "is very remote from the other Leucanidæ, and has some affinities with the BOMBYCITES, but hardly belongs to any described family of Lepidoptera heterocera."

Genus BASIANA, *Walker.*

BASIANA, *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 236 (1856).

611. BASIANA CERVINA, *Walker.*

Basiana cervina, *Walker*, *List Lep. Brit. Mus. Het.*
pt. VIII. p. 237 (1856).

a. b. ♂ ♀. Madras. From Captain J. M. Jones's Collection.

c. ♀. N. India. Presented by the Trustees of the British Museum.

Genus AMBULYX, *Boisduval.*

AMBULYX, *Boisduval*, MS. *Westwood*, *Orient. Ent.* (1847). *Walker*, *List Lep. Brit. Mus. Het. pt. VIII.* p. 120 (1856).

PHOLUS et AMBLYPTERUS, *pt. Hübner.*

612. *AMBULYX SUBSTRIGILIS*, Westwood.

Sphinx (*Ambulyx*) *substrigilis*, *Westwood*, *Orient. Ent.* p. 61, *pl. 30, f. 2* (1847).

Ambulyx substrigilis, *Walker*, *List Lep. Brit. Mus. Het. pt. VIII*. *p. 122*.

Ambulyx Moorei, *Boisduval*, *MS. Mus. E.I.C.*

a. Java. (*A. Moorei*, Boisd.) From Dr. Horsfield's Collection.

b. Canara, S. India. Presented by S. N. Ward, Esq.

Genus *CALYMNIA*, *Boisduval*.

CALYMNIA, *Boisduval*, *MS. Walker*, *List Lep. Brit. Mus. Het. pt. VIII*. *p. 123* (1856).

AMBLYPTERUS, *pt. Hübner*.

613. *CALYMNIA PANOPUS*, Cramer Sp.

Sphinx Panopus, *Cramer*, *Pap. Exot. III*. *p. 50, pl. 224, f. A.B.* (1782).

Calymnia Panopus, *Boisduval*. *Walker*, *List Lep. Brit. Mus. Het. pt. VIII*. *p. 124*.

Amblypterus Panopus, *Hübner*, *Verz. bek. Schmett.* *p. 133*.

Smerinthus? *Panopus*, *Westwood*, *Cabinet Orient. Ent.* *p. 13, pl. 6, f. 2*.

a. b. c. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Calymnia Panopus* are figured on Plate VII., fig. 6, 6a, from Java. "Feeds on the Manga (*Mangifera indica*). May."—(Horsfield, MS.)

Genus *ACHERONTIA*, *Ochsenheimer*.

ACHERONTIA, *Ochsenheimer*, *Schmett. von Eur. IV*. *p. 44* (1816). *Walker*, *List Lep. Brit. Mus. Het. pt. VIII*. *p. 233*.

SPHINX, *pt. Linnæus*.

ACHERONTIA, *pt. Hübner*, *Verz. bek. Schmett.* (1816).

BRACHYGLOSSA, *Boisduval*.

614. *ACHERONTIA STYX*, Westwood.

Acherontia Styx, *Westwood*, *Cabinet Orient. Ent.* *p. 88, pl. 42, f. 3* (1847).

Acherontia Atropos, var., *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 234.*

Acherontia Medusa, *De Cerisy, MS. Mus. E.I.C. (1826).*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♂. Pinang. Presented by Dr. Cantor, 1854.

f. ♀. Dukhun. Presented by Colonel Sykes.

g. h. ♂ ♀. Madras. From Captain J. M. Jones's Collection.

The larva and pupa of *Ach. Styx* are figured on Plate IX. fig. 1, 1a, from India, copied from the original drawings made by Lady Isabella Rose Gilbert.

615. *ACHERONTIA SATANAS*, Boisduval.

Acherontia Satanas, *Boisduval, Spéc. Gén. des Lép. Planches, p. 5, pl. 16, f. 1 (1836).*

Acherontia Lethe, *Westwood, Cabinet Orient. Ent. p. 87, pl. 42, f. 2 (1847).*

Sphinx Atropos, var., *Cramer, Pap. Exot. III. p. 74, pl. 237, f. A.*

? *Acherontia Morta*, *Hübner, Verz. bek. Schmett. p. 140.*

Acherontia Circe, *De Cerisy, MS. Mus. E.I.C. (1826).*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. f. ♂ ♀. Silhet. Presented by Colonel Buckley.

g. ♀. China. Presented by Dr. Cantor.

The larva and pupa of *Ach. Satanas* are figured on Plate IX., fig. 2, 2a, from Java. "Feeds on the Tambaku, Kechubung (*Nicotiana Tabacum*). January and February. Common."—(Horsfield, MS.)

Genus SPHINX, Linnæus.

SPHINX, *Linnæus, Syst. Nat. I. II. p. 796 (1767). Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 211.*

AGRIUS, LETHIA, et PHLEGETHONTIUS, *pt. Hübner.*

616. *SPHINX CONVOLVULI*, Linnæus.

Sphinx Convolvuli, *Linnæus, Syst. Nat. I. 2, p. 798 (1767). Drury, Ins. I. pl. 25, f. 1. Cramer, Pap. Exot. III. p. 55, pl. 225, f. D. Fabricius, Spec. Ins. II. p. 150; Mant. Ins. II. p. 97; Ent. Syst.*

- III. I. p. 374. *Stephens, Ill. Brit. Ent. Haust.*
 I. p. 119; *Cat. Lep. Brit. Mus. pt. I. p. 27. Westwood, Brit. Moths, I. p. 12, pl. 3, f. 1-3. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 212.*
Agrius Convolvuli, Hübner, Verz. bek. Schmett. p. 140.
a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. ♀. Pinang. Presented by Dr. Cantor.
f. g. ♂ ♀. Dukhun. Presented by Colonel Sykes.
h. i. ♂ ♀. Madras. From Captain J. M. Jones's Collection.

The larva and pupa of *Sph. Convolvuli* are figured on Plate IX., fig. 3, 3a, from Java. "Feeds on *Phaseolus Max.* January."—(Horsfield, MS.)

Genus MACROSILA, *Boisduval.*

MACROSILA, *Boisduval, MS. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 198 (1856).*

COCYTIUS, PHLEGETHONTIUS, ACHERONTIA, et HYLOICUS, *pt. Hüb.*

617. *MACROSILA NYCTIPHANES, Boisduval.*

Macrosila Nyctiphanes, *Boisduval, MS. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 209 (1856).*

a. b. c. ♂ ♀. Silhet. Presented by Colonel Buckley.

618. *MACROSILA DISCISTRIGA, Walker.*

Macrosila discistriga, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 209 (1856).*

Sphinx emarginata, *Horsfield, MS. (1826).*

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. ♀. Canara, S. India. Presented by S. N. Ward, Esq.

f. g. ♂ ♀. N. India. Presented by Colonel Buckley.

Genus ZONILIA, *Boisduval.*

ZONILIA, *Boisduval, MS. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 192 (1856).*

NEPHELE, *pt. Hübner.*

619. *ZONILIA MORPHEUS*, Cramer Sp.

Sphinx Morpheus, *Cramer, Pap. Exot. II.* p. 84, pl. 149, f. D. (1779).

Zonilia Morpheus, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 194.

Nephele Morpheus, *Hübner, Verz. bek. Schmett.* p. 133.

Sphinx Hespera, *Fabricius, Spec. Ins. II.* p. 152; *Mant. Ins. II.* p. 91; *Ent. Syst. III. pt. I.* p. 372.

? Sphinx Chiron, *Cramer, Pap. Exot. II.* pl. 137, f. E.

a. b. Dukhun. Presented by Colonel Sykes.

c. d. N. India. Presented by Colonel Hearsey.

e. Canara. Presented by S. N. Ward, Esq.

f. g. Ceylon. From M. Jonville's Collection.

Stirps V. Larvæ OPHTHALMICÆ.

Genus PANACRA, Walker.

PANACRA, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 154 (1856).

620. *PANACRA AUTOMEDON*, Boisduval Sp.

Deilephila Automedon, *Boisduval, MS.*

Panaera Automedon, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 154 (1856).

a. Silhet. Presented by the Trustees of the British Museum.

621. *PANACRA SCAPULARIS*, Horsfield Sp.

Deilephila scapularis, *Horsfield, MS.* (1826).

Panacra scapularis, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 157 (1856).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Pan. scapularis* are figured on Plate IX., fig. 5, 5a, from Java. "Feeds on a species of Arum, bearing the native name of *Lumba*. February and March. Solitary in April and May."—(Horsfield, MS.)

622. *PANACRA VIGIL*, Guérin-Méneville Sp.

Sphinx (Deilephila) Vigil, *Guérin-Méneville, in Delesser's Souv. Voy. dans l'Inde, pt. II. p. 80, pl. 23, f. 1* (1843).

Panacra lignaria, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 156* (1856).

a. b. Ceylon. From M. Jonville's Collection.

The larva, pupa, and imago of this species are figured among Mr. E. Layard's Ceylonese drawings.

623. *PANACRA BUSIRIS*, Boisduval Sp.

Deilephila Busiris, *Boisduval, MS.*

Panaera Busiris, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 158* (1856).

a. Silhet. Presented by the Trustees of the British Museum.

b. Canara, S. India. Presented by S. N. Ward, Esq.

Genus *PHILAMPELUS*, Harris.

PHILAMPELUS, *Harris, Amer. Journ. Sci. XXXVI. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 173.*

PHOLUS, ARGEUS, et DUPO, *pt. Hübner, Verz. bek. Schmett.*

ORNEUS et DAPHNIS, *pt. Hübner, Verz. bek. Schmett.*

ENYO, *pt. Hübner. Walker.*

624. *PHILAMPELUS ANCEUS*, Cramer Sp.

Sphinx Anceus, *Cramer, Pap. Exot. IV. p. 124, pl. 355, f. A.* (1782).

Enyo Anceus, *Hübner, Verz. bek. Schmett. p. 132. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 119.*

a. b. ♂ ♀. Java. From Dr. Horsfield's Collection.

c. ♀. Pinang. Presented by Dr. Cantor.

The larva and pupa of *Phil. Anceus* are figured on Plate IX., fig. 4, 4a, from Java. "Feeds on a species of *Cissus*, bearing the native name of *Jambul*. March."—(Horsfield, MS.)

625. *PHILAMPELUS SERICEUS*, Walker.

Philampelus sericeus, Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 181.

? *Sphinx anceus*, Cramer.

- a. Silhet. Presented by the Trustees of the British Museum.
- b. c. d. e. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

626. *PHILAMPELUS NAGA*, Moore.

Philampelus Naga, n. sp.—Differs from *Phil. sericeus* in being of a paler colour and more silky in appearance: the fore-wing has an oblique blackish streak from middle of costal margin to near middle of outer margin, another from the middle of this to near base of posterior margin, and one tapering along exterior margin from near apex; also various small narrow undulating lines. Expanse $4\frac{1}{2}$ inches.

- a. b. c. d. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus DARAPSA, Walker.

DARAPSA, Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 182 (1856).

DAPHNIS, pt. Hübner.

CHÆROCAMPA, pt. Harris.

627. *DARAPSA HYPOTHOÜS*, Cramer Sp.

Sphinx Hypothoüs, Cramer, Pap. Exot. III. p. 165, pl. 285, f. F. (1782).

Darapsa Hypothoüs, Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 186.

Daphnis Hypothoüs, Hübner, Verz. bek. Schmett. p. 134.

- a. b. c. d. e. f. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Darapsa Hypothoüs* are figured on Plate X., fig. 2, 2a, from Java. "Feeds on the Malati-kosta (*Cadamba jasminiflora*),—April; and on a species of *Contorta*, bearing the native name of *Alluk-antung*. January to April."—(Horsfield, MS.)

Genus DAPHNIS, Hübner.

DAPHNIS, Hübner, Verz. bek. Schmett. p. 134. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 187.

DEILEPHILA, pt. Boisduval, Ind. Méth.

CHÆROCAMPA, Curtis.

628. DAPHNIS NERII, Linnæus Sp.

Sphinx Nerii, Linnæus, Syst. Nat. I. 2, p. 798 (1767).

Fabricius, Spec. Ins. II. p. 142; Mant. Ins. II. p. 93; Ent. Syst. III. I. p. 360. Cramer, Pap. Exot. III. p. 51, pl. 224, f. D.

Daphnis Nerii, Hübner, Verz. bek. Schmett. p. 134. Stephens, Cat. Brit. Lep. p. 29. Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 188.

Deilephila Nerii, Boisduval, Ind. Méth. p. 47.

Chærocampa Nerii, Curtis, Brit. Ent. XIII. pl. 626.

a. b. c. ♂ ♀. Dukhun. Presented by Colonel Sykes.

c. N. India. Presented by Colonel Hearsey.

d. Madras. From Captain J. M. Jones's Collection.

The larva and pupa of *Daphnis Nerii* are figured on Plate X., fig. 3, 3a, copied from the original drawings of General Hardwicke in the British Museum. "Larva found full-grown at Dum-Dum in the beginning of November, which changed to a pupa on the 20th, the imago coming forth on December 21st. Another full-grown larva found on May 7th, changing to pupa on the 11th, the perfect insect emerging on the 24th. Feeds on *Nerium Oleander*."—(Hardwicke, MS. Note.)

Genus PERGESA, Walker.

PERGESA, Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 149 (1856).

DEILEPHILA, pt. Ochsenheimer. Boisduval. Stephens.

THERETRA et OREUS, pt. Hübner.

CHÆROCAMPA, pt. Westwood. Stephens.

629. PERGESA ACTEUS, Cramer Sp.

Papilio Acteus, Cramer, Pap. Exot. III. p. 93, pl. 248, f. A. (1782).

Pergesa Acteus, Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 153.

Oreus Acteus, Hübner, Verz. bek. Schmett. p. 136.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. N. India. From the Asiatic Society of Bengal.

The larva and pupa of *Pergesa Acteus* are figured on Plate X., fig. 1, 1 α , from Java. "Feeds on a species of Arum, bearing the native names of *Senti* and *Suwuy*. January and February. Scarce." —(Horsfield, MS.)

The transformations of this species were also observed by Captain J. Mortimer Slater at Dinapore, in July, 1848, and are figured on page 182 of his original drawings. Also by General Hardwicke (see his Drawings, fig. 172 α).

630. PERGESA CASTOR, Boisduval Sp.

Deilephila Castor, *Boisduval*, MS.

Pergesa Castor, Walker, *List Lep. Brit. Mus. Het.* pt. VIII. p. 153 (1856).

a. b. c. d. ♂ ♀. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

e. f. g. h. ♂ ♀. Darjeeling. From Messrs. Schlagintweit's Collection.

Genus ELIBIA, Walker.

ELIBIA, Walker, *List Lep. Brit. Mus. Het.* pt. VIII. p. 148 (1856).

631. ELIBIA DOLICHUS, Westwood Sp.

Chærocampa Dolichus, Westwood, *Cabinet Orient. Ent.* p. 61, pl. 30, f. 1 (1847).

Elibia Dolichus, Walker, *List Lep. Brit. Mus. Het.* pt. VIII. p. 149.

a. Silhet. Presented by the Trustees of the British Museum.

b. N. India. From Mr. Argent's Collection.

Genus DEILEPHILA, Ochsenheimer.

DEILEPHILA, Ochsenheimer, *Schmett. von Eur.* IV. p. 42 (1816).

Curtis. Stephens. Boisduval. Walker.

PHRYXUS, HYLES, et THAUMAS, pt. *Hübner*.

632. *DEILEPHILA LATHYRUS*, Boisduval.

Deilephila Lathyrus, *Boisduval*, MS. *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 172 (1856).

a. N. India. From Captain J. M. Jones's Collection.

633. *DEILEPHILA LIVORNICA*, Esper Sp.

Sphinx Livornica, *Esper, Ausl. Schmett.* II. pp. 87, 196, pl. 8, fig. 4 (1785); id. II. *Suppl.* 41, pl. 46, f. 3—7. *Hübner, Sphing.* p. 96, pl. 12, f. 65, pl. 23, f. 112.

Deilephila Livornica, *Stephens, Cat. Brit. Lep. Brit. Mus. pt. I.* p. 28. *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 164.

Phryxus Livornica, *Hübner, Verz. bek. Schmett.* p. 137.

Sphinx lineata, *Fabricius, Spec. Ins.* II. p. 147; *Mant. Ins.* II. p. 96; *Ent. Syst. III.* I. p. 368.

Sphinx lineata, *Stephens, Ill. Brit. Lep. Haust.* I. p. 126, pl. 12, f. 1.

a. b. c. d. ♂ ♀. Landoor, N. W. India. Presented by Colonel Hearsey.

Genus CHÆROCAMPA, *Duponchel.*

CHÆROCAMPA, *Duponchel, Hist. Nat. Lép. France, Suppl.* II. p. 159 (1835). *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 126.

DEILEPHILA, *pt. Ochsenheimer. Boisduval.*

OREUS, HIPPOTION, ISOPLES, AMPHION, THERETRA, et THAUMAS, *pt. Hübner.*

METOPSILUS, *pt. Duncan, Brit. Moths*, p. 154.

634. *CHÆROCAMPA CELERIO*, Linnæus Sp.

Sphinx Celerio, *Linnæus, Syst. Nat.* I. 2, p. 800 (1767). *Fabricius, Spec. Ins.* II. p. 151. *Cramer, Pap. Exot.* II. p. 42, pl. 25, f. E.

Chærocampa Celerio, *Stephens, Cat. Brit. Lep.* p. 28. *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 128. *Stainton, Brit. Butt.* I. p. .

Deilephila Celerio, *Stephens, Ill. Brit. Ent. Haust.* I. p. 128. *Boisduval, Ind. Méth. Eur. Lép.* p. 47.

Hippotion Celerio, *Hübner, Verz. bek. Schmett.* p. 135.

Deilephila inquilina, *Harris, Ex.* p. 93, pl. 28, f. 1.

Hippotion Ocys, *Hübner, Verz. bek. Schmett.* p. 135.

a. b. c. d. Java. From Dr. Horsfield's Collection.

e. N. India. Presented by Colonel Sykes.

f. g. h. N. India. Presented by Colonel Hearsey.

The larva and pupa of *Chær. Celerio* are figured on Plate XI., fig. 1, 1 α , copied from fig. 165b of the original drawings of General Hardwicke in the British Museum.

635. CHÆROCAMPA ALECTO, Linnæus Sp.

Sphinx Alecto, *Linnæus, Mus. Lugd. Ulr.* p. 357 ().

Drury, Ins. Exot. II. p. 48, *pl.* 27, *f.* 4. *Cramer,*
Pap. Exot. II. p. 62, *pl.* 137, *f. D.* *Fabricius, Spec.*
Ins. II. p. 152; *Mant. Ins.* II. p. 97; *Ent. Syst.*
III. I. p. 376. *Menetries, Cat. Obj. Zool. Caucase,*
p. 258.

Chærocampa Alecto, *Walker, List Lep. Brit. Mus. Het.*
pt. VIII. p. 130.

Isopes Alecto, *Hübner, Verz. bek. Schmett.* p. 135.

Deilephila Alecto, *Boisduval, Ind. Méth.* p. 46.

Deilephila Cretica, *Boisduval, Icon. Hist. Lep. Sphingidæ,*
p. 20, ♀. *Duponchel, Hist. Nat. Lép. France,* III.
Suppl. p. 15, *pl.* 1, *f.* 3.

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.

e. N. India. Presented by Colonel Buckley.

f. Darjeeling. From Indian Collection, Exposition
 Universelle at Paris, 1855.

"The larva and pupa of *Chær. Alecto* are figured on Plate X.,
 fig. 4, 4 α , from Java."—(Horsfield.)

The transformations of this species were also observed by Captain J. Mortimer Slater at Dinapore, July, 1849, who states that "the larva is common on the vine with *Chær. Clotho*."

The transformations are also figured in General Hardwicke's drawings (fig. 173) in the British Museum.

636. CHÆROCAMPA SUFFUSA, Walker.

Chærocampa suffusa, *Walker, List Lep. Brit. Mus. Het.*
pt. VIII. p. 146 (1856).

a. China. Presented by Dr. Cantor.

b. Borneo. Presented by Hugh Low, Esq.

637. *CHÆROCAMPA PALLICOSTA*, Boisduval Sp.

Deilephila pallicosta, *Boisduval*, *MS.*

Chærocampa pallicosta, *Walker*, *List Lep. Brit. Mus. Het. pt. VIII. p. 145* (1856).

a. b. c. Silhet. Presented by Colonel Buckley.

638. *CHÆROCAMPA THYELIA*, Linnæus Sp.

Sphinx Thyelia, *Linnæus*, *Mus. Lugd. Ulr. p. 360* ();

Syst. Nat. I. 2, p. 803. Clerck, Icon. pl. 46, f. 7, 8.

Cramer, Pap. Exot. I. p. 58, pl. 226, f. E.F. Fabricius, Spec. Ins. II. p. 153; Mant. Ins. II. p. 98; Ent. Syst. III. I. p. 378.

Isopes Thyelia, *Hübner, Verz. bek. Schmett. p. 135.*

Var. *Sphinx Boërhaviae*, *Fabricius, Spec. Ins. II. p. 148; Mant. Ins. II. p. 96; Ent. Syst. III. I. p. 371. Sulzer, Hist. Ins. pl. 20, f. 3.*

Var. *Sphinx Pluto*, *Fabricius, Spec. Ins. II. p. 148.*

Var. *Sphinx Eson*, *Cramer, Pap. Exot. III. p. 57, pl. 226, f. C.*

Isopes Eson, *Hübner, Verz. bek. Schmett. p. 135.*

Chærocampa Eson, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 137.*

Deilephila Rafflesii, *Horsfield, MS. (1826).*

a. b. c. d. Java. From Dr. Horsfield's Collection.

e. China. Presented by Dr. Cantor.

f. N. India. Presented by Colonel Hearsey.

g. h. i. j. Canara. Presented by S. N. Ward, Esq.

639. *CHÆROCAMPA LINEOSA*, Walker.

Chærocampa lineosa, *Walker, List Lep. Brit. Mus. Het. pt. VIII. p. 144* (1856).

a. b. c. Darjeeling. From Indian Collection, Exposition Universelle at Paris, 1855.

d. e. Darjeeling. From Messrs. Schlagintweit's Collection.

640. *CHÆROCAMPA NESSUS*, Drury Sp.

Sphinx Nessus, *Drury, Ill. Exot. Ins. II. p. 46, pl. 27, f. 1* (1773). *Cramer, Pap. Exot. III. pl. 226, f. D.*

Chærocampa Nessus, *Walker*, *List Lep. Brit. Mus. Het.*
pt. VIII. p. 140.

a. b. c. d. e. f. Java. Presented by Dr. Horsfield.

The larva and pupa of *Chær. Nessus* are figured on Plate XI., fig. 2, 2a, from Java. "Feeds on the Uwi (*Dioscorea oppositifolia*). December to March."—(Horsfield, MS.)

641. *CHÆROCAMPA CLOTHO*, *Drury* Sp.

Sphinx Clotho, *Drury*, *Ill. Exot. Ins. II.* p. 48, *pl. 28, f. 1* (1773). *Fabricius*, *Syst. Ent.* p. 540; *Mant. Ins. II.* p. 97; *Ent. Syst. III. I.* p. 376.

Chærocampa Clotho, *Walker*, *List Lep. Brit. Mus. Het.*
pt. VIII. p. 141.

Deilephila Cyrene, *Westwood*, *Cabinet Orient. Ent.* p. 13,
pl. 6, f. 1 (1847).

a. b. c. d. ♂ ♀. Java. From Dr. Horsfield's Collection.
e. China. Presented by Dr. Cantor.

642. *CHÆROCAMPA LUCASI*, *Boisduval* Sp.

Deilephila Lucasii, *Boisduval*, MS.

Chærocampa Leucasii, *Walker*, *List Lep. Brit. Mus. Het.*
pt. VIII. p. 141 (1856).

Deilephila spilota, *Horsfield*, MS. (1826).

a. b. c. d. Java. From Dr. Horsfield's Collection.
e. Canara, S. India. Presented by S. N. Ward, Esq.

The larva and pupa of *Chær. Lucasi* are figured on Plate XI., fig. 3, 3a, from Java. "Feeds on the Galing (? *Cissus*). December to April. Most abundant."—(Horsfield, MS.)

643. *CHÆROCAMPA LYCETUS*, *Cramer* Sp.

Sphinx Lyceetus, *Cramer*, *Pap. Exot. I.* *pl. 61, f. D.* (1779). *Fabricius*, *Ent. Syst. III. I.* p. 371.

Chærocampa Lyceetus, *Walker*, *List Lep. Brit. Mus. Het.*
pt. VIII. p. 143.

Xylophanes Lyceetus, *Hübner*, *Verz. bek. Schmett.* p. 136.
? *Sphinx Draneus*, *Cramer*, *Pap. Exot. II.* p. 56, *pl. 132,*
f. F.

a. N. India. From the Asiatic Society of Bengal.
b. c. N. India. Presented by Colonel Hearsey.

644. *CHÆROCAMPA OLDENLANDIÆ*, Fabricius Sp.

Sphinx Oldenlandiæ, *Fabricius*, *Spec. Ins.* II. p. 148 (1781); *Mant. Ins.* II. p. 96; *Ent. Syst.* III. I. p. 370. *Lewin, Prodr. Ent.* p. 4, pl. 3. *Thon, Ent. Archiv.* pt. I. p. 6, pl. 1, f. 3.

Chærocampa Oldenlandiæ, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 142.

Xylophanes Gortys, *Hübner, Samml. Exot. Schmett.* pp. 58, 513–14.

a. b. c. d. e. ♂ ♀. Java. From Dr. Horsfield's Collection.

The larva and pupa of *Chær. Oldenlandiæ* are figured on Plate XI., fig. 4, 4a, from Java. "Feeds on a species of *Cissus*, bearing the native name of *Galing*. December."—(Horsfield, MS.)

645. *CHÆROCAMPA BISECTA*, Horsfield Sp.

Deilephila Silhetensis, *Boisduval, MS.**

Chærocampa Silhetensis, *Walker, List Lep. Brit. Mus. Het. pt. VIII.* p. 143.

Deilephila bisecta, *Horsfield, MS.* (1826).

a. b. c. d. Java. From Dr. Horsfield's Collection.

e. N. India. Presented by Colonel Buckley.

The larva and pupa of *Chær. bisecta* are figured on Plate XI., fig. 5, 5a, from Java. "Feeds on a species of *Boërhavia*, bearing the native name of *Chakas-ayam*. February and March."—(Horsfield, MS.)

* This name is cancelled, as being inapplicable.

APPENDIX.

Page.

17. Add the following to Tribe I. :—

PAPILIONES, *Swainson, Zool. Ill. 2nd ser. pl. 114* (1832-3); id.
Cabinet Cyclop. p. 58 (1840).

ACHALINOPTERA, *pt. Blanchard, Hist. Nat. des Ins. II. pp. 322-3*
(1845).

SUSPENSI, *Boisduval, Spéc. Gén. Lép. I. pp. 162, 164* (1836).

PENDULÆ, *Boisduval, Ind. Méth. Eur. Lép. p. 15* (1840).

NUDI, *Westwood, Introd. to Ent. II. p. 347*; id. *Syn. p. 87*
(1840).

18. Add the following to Stirps I. :—

ONISCIFORM, *Westwood.*

SUCTORIAL, or VERMIFORM, *Swainson, Cab. Cyclop. pp. 58, 61*
(1840).

LEP. HEXAPODA, *pt. Latreille, Règ. Anim. V.* (1829).

LYCÄNINÆ, *Swainson, Cabinet Cyclop. pp. 87, 89* (1840).

LYCENITES, *Blanchard, Hist. Nat. des Ins. II. p. 342* (1845).

ERYCINII, *pt. Blanchard, id. pp. 322, 341.*

21. No. 7. The larva and pupa of *Polyommatus Laius* are figured on Plate XII., fig. 1, 1 a , copied from the original drawings in the possession of A. Grote, Esq., of Calcutta.

44. No. 67. The larva and pupa of *Amblypodia Timoleon* are figured on Plate XII., fig. 3, 3 a , copied from the original drawings in the possession of A. Grote, Esq., of Calcutta.

49. No. 81. The larva and pupa of *Myrina Triopas* are figured on Plate XII., fig. 5, 5 a , from A. Grote, Esq.

No. 82. The larva and pupa of *Myrina Etolus* are figured on Plate XII., fig. 6, 6 a , from A. Grote, Esq.

52. No. 93. The larva and pupa of *Anops Thetys* are figured on Plate XII., fig. 7, 7 a , from A. Grote, Esq.

55. Add the following to Stirps II. A :—

PAPILIONII, *pt. Blanchard.*

PAP. PARNASII, *pt. Linnæus.*

Page.

56. No. 99. The larva and pupa of *Callidryas Philippina* are figured on Plate XII., fig. 8, 8a, from A. Grote, Esq.
72. No. 138. The larva and pupa of *Pieris Mesentina* are figured on Plate XII., fig. 9, 9a, from A. Grote, Esq.; also figured among General Hardwicke's drawings (fig. 89) in the British Museum.
73. No. 142. Add page 103 and plate 44.
74. No. 143. Add page 103 and plate 44.
76. No. 148. Add page 102 and plate 44.
78. No. 153. Add page 102 and plate 44.
79. No. 155. Add page 103 and plate 44.
80. No. 160. The transformations of *Pieris Eucharis* were observed by Captain Mortimer Slater, and are figured on page 396 of his "Notes." "The larva was taken October 1st, changing to a pupa the next day, and the imago emerging on the 13th; the larva feeds upon the *Guava*."

The transformations of this species were also observed by Mrs. Hamilton in Moulmein, in 1853, and are figured among that lady's original drawings, now in the possession of the Entomological Society of London. The "larva was taken on the Lime-tree, December 12."

Also observed by A. Grote, Esq., and figured among his original drawings.

85. Genus *Leptocircus*. Add as syn. LAMPROPTERA, G. R. Gray, Griffith's edit. Cuvier, Ins. II. t. 102, f. 4 (1832).
88. No. 178. "Common at Dacea, 1845, and at Darjeeling, being partial to the feathery scarlet-flowered plant about which they hover, and may be easily caught by the hand, as its flight is heavy."—(Captain Mortimer Slater's Notes, p. 390.)
91. No. 187. The transformations of the female of this species were also observed by Mrs. Hamilton at Moulmein, and figured among that lady's drawings.
97. No. 198. Add page 104 and plate 45.
99. No. 202. Captain Mortimer Slater, in his MS. "Notes," p. 420, remarks: "This butterfly is very active, and difficult to catch, flying in and out of deep forest shades with great velocity, and generally high."
105. No. 211. "Very common throughout the Bengal presidency; the larva feeding on the Lime, Orange, and "Bel" tree, and has the two processes (above the head) very pliable and completely retractile. The scent of these processes is that

Page.

of very rotten oranges, and is overpowering in the extreme, and does not leave the hand, when once infected, for many hours, even after several washings. The larva, when touched, shrugs up its head, and only displays the nauseous filaments when much or repeatedly irritated, and then it endeavours to smear the irritating body by dragging them over and about it."—(Captain Mortimer Slater, MS. "Notes," p. 438.)

The transformations were also observed by A. Grote, Esq., and by Mrs. Hamilton, which lady states that "the larva of this butterfly feeds on the Lime and Citron; the eggs are laid singly on the tips of the leaves, and generally on the upper-side. The larva is at first of a greenish-brown."

113. No. 227. The larva and pupa of *Papilio Eurypylus* are figured on Plate XII., fig. 10, 10a, from A. Grote, Esq., of Calcutta.

126. No. 249. The transformations of *Danais Chrysippus* were also observed by Lady Gilbert, and figured in Vol. II., p. 33, of her ladyship's drawings. "Feeds on the Madar, *Asclepias gigantea*. The larva taken July 14th, changing on the 15th, and the imago coming forth on the 23rd."

No. 256. Add specimens b. c. ♂. Borneo. Presented by Edwin Shepherd, Esq.

131. No. 262. The transformations of *Euploea Core* were also observed and figured by Captain Mortimer Slater.

142. No. 288. The larva and pupa of *Junonia Almana* are figured on Plate XII., fig. 12, 12a, from A. Grote, Esq., of Calcutta.

163. Genus NEPTIS. Vide monograph in Proc. Zool. Soc.

167. No. 345. Dele the letter h in Jumbah.

169. Genus ATHYMA. Vide monograph in Proc. Zool. Soc.

185. Genus ADOLIAS. Vide monograph in Proc. Zool. Soc., where the new species are figured.

188. No. 379. The larva and pupa of *Adolias Lubentina* are figured on Plate XII., fig. 13, 13a, from A. Grote, Esq., of Calcutta.

207. No. 424. The larva and pupa of *Nymphalis Baya* are figured on Plate XII., fig. 14, 14a, copied from Mrs. Hamilton's original drawings.

211. No. 431. The larva and pupa of *Discophora Tullia* are figured on Plate XII., fig. 15, 15a, from A. Grote, Esq., of Calcutta.

222. No. 461. The transformations of this species also observed

- and figured by Lady Gilbert, who states: "Feeds upon grass. Larva taken August 1st, changing to a pupa on the 5th, and the imago emerging on the 14th."—(Vol. I. p. 41.)
237. No. 511. "Not uncommon at Dacca, 1845. Flies in a weak, hovering manner, generally over the deep damp pits formerly used by the muslin manufacturers, and many years deserted."—(Captain Mortimer Slater's Notes, p. 248.)

INDEX

OF

GENERA AND SPECIES.

VOL. I.

ABAEIS , <i>Hübner</i>	<i>p.</i> 63	Alcathoë , <i>Godart</i> , <i>Danaïs</i>	<i>p.</i> 131
Ablata , <i>Zincken-Sommer</i> , <i>Euplecta</i>	124	Alcibiades , <i>Fabr.</i> , <i>Papilio</i>	116
Acalis , <i>Godart</i> , <i>Pieris</i>	80	Alcidice , <i>Godart</i> , <i>Danaïs</i>	126
Acanthe , <i>Hübner</i> , <i>Tisiphone</i>	228	Alcinous , <i>Klug</i> , <i>Papilio</i>	95
Acca , <i>Hübner</i>	163, 170, 178	Alciope , <i>Godart</i> , <i>Argynnus</i>	151
Aceris , <i>Esper</i> , <i>Papilio</i>	168	Alcippe , <i>Cramer</i> , <i>Papilio</i>	151
Achanta , <i>Donovan</i> , <i>Papilio</i>	228	Alcithoë , <i>Cramer</i> , <i>Papilio</i>	159
Achates , <i>Cramer</i> , <i>Papilio</i>	100	Alcemene , <i>Cramer</i> , <i>Papilio</i>	159
Achatiades , <i>Hübner</i> , <i>Papilio</i>	101	Alcemeone , <i>Cramer</i> , <i>Papilio</i>	56
ACHERONTIA , <i>Hübner</i>	266	Alcyone , <i>Cramer</i> , <i>Papilio</i>	56
ACHILLIDES , <i>Hübner</i>	89	Alea , <i>Hübner</i> , <i>Terpsichrois</i>	131
ACHLYODES , <i>Hübner</i>	251	Alecto , <i>Linn.</i> , <i>Sphinx</i>	275
ACIDALIA , <i>Hübner</i>	155	Alexis , <i>Stoll</i> , <i>Papilio</i>	25
ACONTHEA , <i>Horsf.</i>	185	Allica , <i>Fabr.</i> , <i>Papilio</i>	242
Aconthea , <i>Cramer</i> , <i>Papilio</i>	185	Almana , <i>Linn.</i> , <i>Papilio</i>	142
Acte , <i>Doubleday</i> , <i>Myrina</i>	47	Alopia , <i>Godart</i> , <i>Danaïs</i>	131
Acteus , <i>Cramer</i> , <i>Sphinx</i>	272	Alpheda , <i>Godart</i> , <i>Nymphalis</i>	188
Adima , <i>Moore</i> , <i>Adolias</i>	194	Alphenor , <i>Fabr.</i> , <i>Papilio</i>	102
ADOLIAS , <i>Boisd.</i>	185	Alysos , <i>Boisd.</i> , <i>Hesperia</i>	253
Adonia , <i>Cramer</i> , <i>Papilio</i>	188	Amara , <i>Moore</i> , <i>Ismene</i>	247
Ægistus , <i>Cramer</i> , <i>Papilio</i>	114	AMARYSSUS , <i>Dalman</i>	89
Ælianuſ , <i>Fabr.</i> , <i>Hesperia</i>	25	Amata , <i>Fabr.</i> , <i>Papilio</i>	67
Ænipe , <i>Cramer</i> , <i>Papilio</i>	66	AMATHUSIA , <i>Fabr.</i>	208, 209
Æole , <i>De Haan</i> , <i>Cethosia</i>	154	AMAURIS , <i>Hübner</i>	89, 121
Æsopuſ , <i>Fabr.</i> , <i>Papilio</i>	53	Ambalika , <i>Moore</i> , <i>Adolias</i>	192
Agama , <i>Moore</i> , <i>Pyrgus</i>	249	Ambasa , <i>Moore</i> , <i>Ismene</i>	248
Agamemnon , <i>Linn.</i> , <i>Papilio</i>	114	Ambasara , <i>Moore</i> , <i>Melanitis</i>	223
Agathon , <i>G. R. Gray</i> , <i>Pieris</i>	83	Ambica , <i>Kollar</i> , <i>Apatura</i>	201
Agenor , <i>Linn.</i> , <i>Papilio</i>	101	AMBLYPODIA , <i>Horsf.</i>	39
Agestor , <i>G. R. Gray</i> , <i>Papilio</i>	91	AMBLYPTERUS , <i>Hübner</i>	265
Agetes , <i>Westw.</i> , <i>Papilio</i>	116	AMBULYX , <i>Boisd.</i>	265
AGLAURA , <i>Boisd.</i>	210	Amerta , <i>Moore</i> , <i>Nisoniades</i>	250
Aglea , <i>Cramer</i> , <i>Papilio</i>	122	AMNOSIA , <i>Boisd.</i>	146
Agna , <i>Moore</i> , <i>Hesperia</i>	255	Amor , <i>Fabr.</i> , <i>Papilio</i>	49
Agostina , <i>Hewitson</i> , <i>Pieris</i>	81	AMPHION , <i>Hübner</i>	274
AGRIUS , <i>Hübner</i>	267	AMPHRISIUS , <i>Swainson</i>	86
Akasa , <i>Horsf.</i> , <i>Polyommatus</i>	20	Amphrisius , <i>Cramer</i> , <i>Papilio</i>	88
Alankara , <i>Horsf.</i> , <i>Aconthea</i>	179	Amyntas , <i>Fabr.</i> , <i>Papilio</i>	22
ALAZONIA , <i>Hübner</i>	153	Amyntor , <i>Herbst</i> , <i>Papilio</i>	49
Alecanor , <i>Hübner</i> , <i>Symphaædra</i>	204	Amythaon , <i>Doubleday</i> , <i>Amathusia</i>	210
Alcanor , <i>Cramer</i> , <i>Papilio</i>	100	Anais , <i>Bougainville</i> , <i>Danaïs</i>	61

- Ananta, *Moore*, Neptis p. 166
 ANAPHLEA, *Hübner* 70
 Anapita, *Moore*, Mycalesis 232
 ANARTIA, *Hübner* 145
 Anceus, *Cramer*, Papilio 101
 — *Cramer*, Sphinx 270
 Androcles, *Boisd.*, Thecla 29
 Androgeos, *Cramer*, Papilio 99
 Anexibia, *Hübner*, Ixias 67
 Angelica, *Cramer*, Papilio 138
 Annada, *Moore*, Erebia 226
 Anomala, *Kollar*, Acræa 136
 ANOFS, *Boisd.* 52
 ANOSIA, *Hübner* 121
 Anosia (*Boisd.*), *Moore*, Adolias 189
 ANTEOS, *Hübner* 59
 ANTHOCHARIS, *Doubleday* 69
 Anthrax, *Boisd.*, Euploea 131
 Antierates, *Doubleday*, Papilio 115
 Antigone, *Cramer*, Papilio 159
 Antilope, *Cramer*, Papilio 160
 Antiope, *Linn.*, Papilio 137
 Antiphates, *Cramer*, Papilio 116
 Antiphus, *Fabr.*, Papilio 94
 Antonöe, *Cramer*, Papilio 81
 Aonis, *Cramer*, Papilio 139
 Aoris, *Doubleday*, Cirrochroa 149
 APATURA, *Fabr.* 158, 201
 Apaturina, *Horsf.*, Aconthea 197
 Aphnaeus, *Hübner* 30, 37
 APHRODITE, *Hübner* 68
 Apiades, *Menetries*, Adolias 195
 Apidanus, *Cramer*, Papilio 39
 APORIA, *Hübner* 83
 APPIAS, *Hübner* 70
 Apriate, *Hübner*, Delias 79
 Aratus, *Cramer*, Papilio 22
 Arbates, *Zinken-Sommer*, Papilio 99
 Areadia, *Cramer*, Papilio 219
 Arcensia, *Cramer*, Papilio 223
 Arcesilaus, *Fabr.*, Papilio 213
 Arcturus, *Westwood*, Papilio 108
 Arcuatum, *Horsf.*, Macroglossa 262
 Arete, *Cramer*, Papilio 216
 ARGEUS, *Hübner* 270
 ARGYNNIS, *Fabr.* 155
 Argynnis, *Drury*, Papilio 156
 Argyrus, *Sparm.*, Papilio 156
 ARGYRONOME, *Hübner* 155
 ARIOPALA, *Boisd.* 39
 Aria, *Moore*, Hesperia 254
 ARIADNE, *Horsf.* 144
 Ariadne, *Linn.*, Papilio 144
 ARISBEE, *Hübner* 89
 Arjuna, *Horsf.*, Papilio 107
 Arne, *Klug*, Pontia 68
 Arsinoë, *Cramer*, Papilio 145
 Aruna, *Moore*, Argynnus 156
 ASCANIDES, *Hübner* 89
 Asmara, *Moore*, Hesperia 253
 Assimilis, *Linn.*, Papilio 160
- Astenous, *Fabr.*, Papilio p. 87
 Asterie, *Cramer*, Papilio 142
 Astina, *Horsf.*, Papilio 89
 Astorion, *Westwood*, Papilio 98
 ASTRAPTES, *Hübner* 244
 Astyanax, *Fabr.*, Papilio 103
 Asura, *Moore*, Athyma 171
 Atalanta, *Cramer*, Papilio 138
 Ataxus, *Boisd.*, Thecla 31
 ATELLA, *Doubleday* 151
 Athamas, *Drury*, Papilio 205
 Athamis, *Hübner*, Eriboea 205
 ATHYMA, *Westwood* 169
 Atratus, *Horsf.*, Lycena 22
 Atymnus, *Cramer*, Papilio 51
 Auge, *Cramer*, Papilio 158
 AUGIADES, *Hübner* 250
 Augias, *Linn.*, Papilio 251
 Aurora, *Cramer*, Papilio 68
 Aurota, *Fabr.*, Papilio 72
 Automedon, *Boisd.*, Deilephila 269
 AUTONEMA, *Boisd.* 148
 Autothisbe, *Hübner*, Delias 82
 Avatar, *Moore*, Eronia 61
 Avatara, *Moore*, Satyrus 229
 Aventina, *Cramer*, Papilio 122
 Avia, *Fabr.*, Papilio 159
 Badra, *Moore*, Goniloba 245
 Bajadate, *Moore*, Cirrochroa 150
 Baldus, *Fabr.*, Papilio 235
 Balidia, *Boisd.*, Thesrias 67
 Banksia, *Fabr.*, Papilio 222
 BASIANA, *Walker* 265
 Basilissa, *Cramer*, Papilio 133
 Batara, *Moore*, Ismene 249
 Bathycles, *Zink.-Som.*, Papilio 114
 Baya, *Moore*, Nymphalis 207
 Bebryce, *Hübner*, Ixias 66
 Bela, *Moore*, Melanitis 223
 BELENOIS, *Hübner* 70
 Belenus, *Hübner*, Heliophorus 29
 Belia, *Westwood*, Hestia 135
 Belis, *Cramer*, Sphinx 263
 Belisama, *Fabr.*, Papilio 82
 Belladonna, *Cramer*, Papilio 78
 Benjaminii, *Guérin*, Thymele 248
 Bernardus, *Fabr.*, Papilio 206
 Beroë, *Cramer*, Papilio 216
 Bhadra, *Moore*, Enope 227
 Bhagava, *Moore*, Goniloba 246
 Bhairava, *Moore*, Debis 217
 Bianor, *Cramer*, Papilio 110
 BIBLIA, *Hübner* 239
 Biblina, *Godart*, Cethosia 154
 BIBLIS, *Fabr.* 144, 170, 236
 Biblis, *Drury*, Papilio 153
 Bisaltide, *Cramer*, Papilio 209
 Bisecta, *Horsf.*, Deilephila 278
 Blanda, *Boisd.*, Terias 64
 Boerhaviae, *Fabr.*, Sphinx 276
 Boeticus, *Linn.*, Papilio 23

Boisduvalii, <i>Moore</i> , Miletus	p. 19	Chiron, <i>Cramer</i> , Sphinx	p. 269
— <i>G. R. Gray</i> , Adolias	198	Chirona, <i>G. R. Gray</i> , Danais	125
Bolina, <i>Linn.</i> , Papilio	158	Chitra, <i>Horsf.</i> , Thecla	51
BRACHYGLOSSA, <i>Boisd.</i>	266	CHLORIPPE, <i>Boisd.</i>	201
Brahma, <i>Moore</i> , Ilerda	29	CHLORISSES, <i>Swainson</i>	89
Brahmina, <i>Boisd.</i> , Zeritis	37	Chromus, <i>Cramer</i> , Papilio	244
Brahminus, <i>Blanch.</i> , Satyrus	229	Chryseis, <i>Drury</i> , Papilio	56
BRENTHIS, <i>Hübner</i>	155	Chrysippus, <i>Linn.</i> , Papilio	126
Buddha, <i>Moore</i> , Calinaga	163	Chrysomallus, <i>Hübner</i> , Zesiuss ..	33
Bulis, <i>Boisd.</i> , Anops	53	CHRYSOPLANUS, <i>Hübner</i>	27
Busiris, <i>Boisd.</i> , Deilephila	270	Cicero, <i>Fabr.</i> , Hesperia	252
Cadma, <i>Boisd.</i> , Thecla	28	Cinnara, <i>Moore</i> , Hesperia	255
CELENORRHINUS, <i>Hübner</i>	250	Cinyla, <i>Cramer</i> , Papilio	53
Caius, <i>Fabr.</i> , Hesperia	21	Ciprea, <i>Fabr.</i> , Papilio....	67
Calais, <i>Cramer</i> , Papilio	67	Circe, <i>De Cerisy</i> , Acherontia	267
CALAIDES, <i>Hübner</i>	89	CIRROCHROA, <i>Doubleday</i>	149
CALINAGA, <i>Moore</i>	162	Clagia, <i>Godart</i> , Argynnis	149
CALLIDRYAS, <i>Boisd.</i>	85	Clarissa, <i>Boisd.</i> , Terinos	149
Callirhoe, <i>Hübner</i> , Hamadryas ..	138	Clemanthe, <i>Doubleday</i> , Pieris....	77
CALLOSUNE, <i>Doubleday</i>	68	Cleona, <i>Cramer</i> , Papilio	123
Calybe, <i>Godart</i> , Vanessa	141	CLEROME, <i>Boisd.</i>	213
CALYMNIA, <i>Boisd.</i>	266	Cloanthus, <i>Westwood</i> , Papilio ..	112
Cama, <i>Moore</i> , Athyma.....	174	Clotho, <i>Drury</i> , Sphinx....	277
Camadeva, <i>Westw.</i> , Thaumantis..	216	CLYTIA, <i>Swainson</i>	89
Camdeo, <i>Doubleday</i> , Amblypodia	41	Clytia, <i>Linn.</i> , Papilio	92
Canens, <i>Hübner</i> , Faunis	213	Cneus, <i>Fabr.</i> , Hesperia	22
Carduelis, <i>Cramer</i> , Papilio	139	Cocytta, <i>Fabr.</i> , Papilio	194
Cardui, <i>Linn.</i> , Papilio	138	Cocytina, <i>Horsf.</i> , Aconthea	193
Cashmirensis, <i>Kollar</i> , Vanessa ..	137	COCYTIUS, <i>Hübner</i>	268
CASTALIA, <i>Boisd.</i>	199	Cocytus, <i>Fabr.</i> , Papilio	194
CASTNIUS, <i>Hübner</i>	247	CŒLITES, <i>Boisd.</i>	225
Castor, <i>Westwood</i> , Papilio	93	Coenæos, <i>Hübner</i> , Euchloë	69
— <i>Boisd.</i> , Deilephila	273	COLIAS, <i>Fabr.</i>	55, 62
CATARGYRIA, <i>Hübner</i>	201	COLOTIS, <i>Hübner</i>	55, 62
CATHÆMIA, <i>Hübner</i>	70	Columbina, <i>Cramer</i> , Papilio ...	152
Catilla, <i>Cramer</i> , Papilio	57	Columella, <i>Cramer</i> , Papilio....	166
CATOPHAGA, <i>Hübner</i>	70	Columena, <i>Hübner</i> , Acca....	167
CATOPSILIA, <i>Hübner</i>	55	Coluteæ, <i>Rossi</i> , Papilio	24
Caumas, <i>Godart</i> , Satyrus	219	Consimilis, <i>Westwood</i> , Hestina ..	161
C. aureum, <i>Linn.</i> , Papilio	138	Constantia, <i>Cramer</i> , Papilio ...	221
Celeno, <i>Cramer</i> , Papilio	24	Convolvul, <i>Linn.</i> , Sphinx	267
Celerio, <i>Fabr.</i> , Papilio	24	Coon, <i>Fabr.</i> , Papilio	95
— <i>Linn.</i> , Sphinx	274	Core, <i>Cramer</i> , Papilio	131
Celinde, <i>Stoll</i> , Papilio	211	Coreisia, <i>Hübner</i> , Hypolimnas ..	197
Centaurus, <i>Fabr.</i> , Papilio	40	Coreta, <i>Godart</i> , Danais	132
Cephea, <i>Cramer</i> , Papilio	135	Coriolanus, <i>Fabr.</i> , Hesperia	243
CEPHONODES, <i>Hübner</i>	260, 261	Cornelia, <i>Fabr.</i> , Papilio	58
Cervina, <i>Walker</i> , Basiana ..	265	Coronea, <i>Cramer</i> , Papilio	82
CETHOSIA, <i>Fabr.</i>	153	Coronis, <i>Cramer</i> , Papilio....	76
CHÆROCAMPÀ, <i>Dupont.</i>	274	Corydon, <i>Cramer</i> , Papilio ...	26
Chandica, <i>Moore</i> , Debis	219	Coryta, <i>Cramer</i> , Papilio	144
Chandra, <i>Moore</i> , Castalia ..	200	Corythus, <i>Boisd.</i> , Macroglossa ..	262
Chandrasa, <i>Moore</i> , Achlyodes..	251	Crameri, <i>Moore</i> , Euplora ..	129
Chaon, <i>Westwood</i> , Papilio	102	CRASTIA, <i>Hübner</i>	126
— <i>Boisd.</i> , Ismene....	249	Cresphontes, <i>Fabr.</i> , Papilio ...	105
Chara, <i>Westwood</i> , Papilio	98	CRETEUS, <i>Hübner</i>	244
CHARAXES, <i>Ochsenheimer</i>	204	Cretica, <i>Boisd.</i> , Deilephila....	275
Charonia, <i>Drury</i> , Papilio....	137	Crino, <i>Fabr.</i> , Papilio....	109
Chaya, <i>Moore</i> , Hesperia	255	Crishna, <i>Westwood</i> , Cyollo ..	221
Chenu, <i>Guérin</i> , Satyrus	235	Crisia, <i>Hübner</i> , Eptychia	226
Childreni, <i>G. R. Gray</i> , Argynnis	157	Crithoë, <i>Boisd.</i> , Pieris....	79

- Crocale, *Cramer*, Papilio p. 57
 Curius, *Fabr.*, Papilio 85
 Cyane, *Fabr.*, Papilio 155
CYCLOPIDES, *Hübner* 250
 Cycenus, *Westwood*, Emispe 212
CYLLO, *Boisd.* 221
 CYMOTHOE, *Hübner* 185
CYNTHIA, *Fabr.* 136, 145
 Cyrene, *Westwood*, Deilephila 277
CYRESTIS, *Boisd.* 146
 Cyrus, *Fabr.*, Papilio 104
 Daha, *Moore*, Nisoniades 250
 Damaris, *G. R. Gray*, Pieris 225
 Damoëtes, *Fabr.*, Papilio 23
 Dan, *Fabr.*, Hesperia 253
 Danaë, *Fabr.*, Papilio 69
DANAIDA, *Latr.* 121
DANAI, *Latr.* 121, 126
DANAUS, *Latr.* 121, 126
 Danava, *Moore*, Limenitis 180
 Danna, *Moore*, Pyrgus 249
 Daos, *Boisd.*, Idea 124
DAPHNIS, *Hübner* 270, 271, 272
 Daplidice, *Linn.*, Papilio 75
DARAPSA, *Walker* 271
 Daraxa, *Doubleday*, Limenitis 180
 Darsius, *G. R. Gray*, Ornithoptera 87
 Dasa, *Moore*, Ismene 248
 Dasarada, *Moore*, Papilio 96
DEBIS, *Boisd.* 216
 Decora, *Boisd.*, Amnossia 146
DEILEPHILA, *Ochs.* 272, 273
 Deione, *Westwood*, Euplcea 132
 — *Erichson*, Cynthia 145
 Deiopeia, *Donovan*, Papilio 83
 Delcessertii, *Guérin*, Papilio 91
DELIAS, *Hübner* 70
 Delphis, *Doubleday*, Charaxes 206
 Demeter, *Doubleday*, Nemeobius 242
 Demolion, *Cramer*, Papilio 105
 Dentatus, *Cramer*, Sphinx 264
DERCAS, *Boisd.* 59
 Derma, *Kollar*, Adolias 193
 — *Moore*, Gonilobia 245
 Descombesi, *Boisd.*, Pieris 81
 Deva, *Moore*, Amblypodia 46
DIADEMA, *Boisd.* 158
 Dichroa, *Kollar*, Limenitis 199
DIDONIS, *Hübner* 126
DILIPA, *Moore* 201
 Diocippus, *Cramer*, Papilio 158
 Diocles, *Boisd.*, Thanaos 250
 Diocletianus, *Fabr.*, Papilio 126
 Diore, *Doubleday*, Thaumantis 215
 Diphius, *Esper*, Papilio 94
DIPSAS, *Doubleday* 30, 39
DIRA, *Hübner* 227
 Dirtea, *Fabr.*, Papilio 198
 Discistriga, *Walker*, Macrosila 268
 Disconthea, *Godart*, Nymphalis 185
DISCOPHORA, *Boisd.* 211
 Dissimilis, *Linn.*, Papilio p. 91
 Divergens, *Walker*, Macroglossa 263
 Divodasa, *Moore*, Hesperia 255
 Dodonea, *Boisd.*, Amblypodia 43
 Dolichus, *Westwood*, Chærocampa 273
 Dolon, *Westwood*, Charaxes 206
 Dorimond, *Stoll*, Papilio 39
 Dorippa, *Boisd.*, Danais 123
 Doris, *Cramer*, Papilio 234
DORITIS, *Fabr.* 84
 Doubledayii, *Boisd.*, Adolias 196
 — *Westwood*, Aconthea 198
DOXOCOPA, *Hübner* 201
 Drancus, *Cramer*, Sphinx 277
 Drona, *Horsf.*, Terias 65
 Druna, *Moore*, Hesperia 254
 Drupadi, *Horsf.*, Emesis 244
 Drusia, *Cramer*, Papilio 231
DRUSILLA, *Swainson* 214
 Dryas, *Boisd.*, Smerinthus 264
 Dudu, *Doubleday*, Limenitis 180
DUPO, *Hübner* 270
 Durga, *Moore*, Adolias 196
 — *Kollar*, Melitæa 242
 Durvaza, *Moore*, Pieris 73
 Dusara, *Horsf.*, Melanitis 238
 Dynamene, *Klug*, Pontia 68
 Eacus, *Latr.*, Hesperia 253
EANTIS, *Boisd.* 251
 Eborea, *Cramer*, Papilio 69
 Echerius, *Stoll*, Papilio 243
 Echidna, *De Haan*, Papilio 92
 Edmondi, *Bougainville*, Danais 125
 Edusa, *Fabr.*, Papilio 62
 Ega, *Boisd.*, Pieris 72
 Egeon, *Boisd.*, Emesis 242
 Egialea, *Cramer*, Papilio 79
 Egista, *Cramer*, Papilio 152
 Eleonor, *Boisd.*, Pieris 71
ELIBIA, *Walker* 273
 Elpis, *Godart*, Polyommatus 24
 Elusine, *Cramer*, Papilio 127
ELYMNIAS, *Hübner* 144, 236
 Emarginata, *Horsf.*, Sphinx 268
EMESIS, *Horsf.* 242
ENISPE, *Doubleday* 212
ENODIA, *Hübner* 228
ENOPE, *Moore* 228
ENYO, *Hübner* 263, 270
 Enyo, *Boisd.*, Pieris 71
EPARGYREUS, *Hübner* 244
 Epicharis, *Godart*, Pieris 80
 Epicles, *Godart*, Polyommatus 29
EPIGEA, *Hübner* 226
 Epitarbas, *Boisd.*, Thecla 32
 Epilais, *Hübner*, Doxocopa 207
 Epiminthia, *Boisd.*, Cœlites 225
 Epimuta, *Boisd.*, Amblypodia 42
EPINEPHILE, *Hübner* 227, 228
 Epiona, *G. R. Gray*, Adolias 196
 Epius, *Fabr.*, Papilio 106

Erato, Boisd., Emesia	p. 243	Ganga, Moore, Abrota	p. 178
EREBIA, Dalman	226	Garuda, Moore, Adolias	186
ERGOLIS, Boisd.	144	Gaura, Horsf., Idea	134
ERIBOEA, Hübner	204	Genutia, Cramer, Papilio	124
Erigone, Cramer, Papilio	139	Geometricum, Horsf., Macroglossum	
Eriphile, Cramer, Papilio	159	sum	263
ERITES, Boisd.	229	GERYDUS, Boisd.	18
Erithonius, Cramer, Papilio	105	Gilia, Boisd., Macroglossa	262
ERONIA, Hübner	60	Glypha, Godart, Pieris	75
Erosine, Cramer, Papilio	170	Glaucippe, Linn., Papilio	60
Erylus, Godart, Polyommatus	50	Glicinia, Cramer, Papilio	75
Erymanthis, Drury, Papilio	150	Glycerion, G. R. Gray, Papilio	116
Esa, Moore, Hesperia	254	Gnoma, Fabr., Papilio	56
Eson, Cramer, Sphinx	276	Godartii, G. R. Gray, Adolias	194
ESOPTRIA, Hübner	158	Gokala, Moore, Melanitis	224
Etolus, Cramer, Papilio	38	GONEPTERYX, Leach	59
— Fabr., Papilio	49	GONIAPTERYX, Westwood	59
Etrida, Boisd., Anthocharis	69	GONILOBA, Westwood	244
Eucharis, Fabr., Papilio	68	Gopala, Moore, Goniloba	246
— Drury, Papilio	80	Gortys, Hübner, Xylophanes	278
EUCHLOE, Hübner	68	Gotama, Moore, Mycalesis	232
Eudamippus, Doubl., Charaxes	206	Grammica, Boisd., Danais	122
Eudora, G. R. Gray, Hestia	134	GRAPTA, Kirby	137
EUGONIA, Hübner	136, 137	Gyas, Westwood, Papilio	110
EUMENIS, Hübner	228	Gyrans, Boisd., Macroglossa	262
Eumolpus, Cramer, Papilio	41	Halitherses, Doubleday, Euripus	199
Eunice, Godart, Danaus	129	Hallirothius, Westwood, Euripus	199
EUPHLEADES, Hübner	89	HAMANUMIDA, Hübner	242
Euphanes, Esper, Papilio	204	Hara, Moore, Precis	134
Euphranor, Doubl., Amblypodia	31	Hardwickii, G. R. Gray, Parnassius	84
Euphrone, Westwood, Nymphalis	160	Harina, Horsf., Terias	64
EUPLEA, Fabr.	121, 126	Harisa, Moore, Ismene	247
EUREMA, Hübner	63	HEBOMOIA, Hübner	60
EURIFUS, Doubleday	198	Hecabe, Linn., Papilio	63
Europa, Fabr., Papilio	216	HECAERGE, Hübner	239
EURYMUS, Swainson	62	Hector, Linn., Papilio	93
Eurynome, Westwood, Limenitis	169	HECTORIDES, Hübner	89
Euryphilus, Linn., Papilio	113	Hegesippus, Cramer, Papilio	125
EUTHALIA, Hübner	185	Helenus, Linn., Papilio	101
Euthymius, Doubleday, Enispe	212	Heliacon, Fabr., Papilio	87
Evagete, Cramer, Papilio	77	Heliodore, Fabr., Papilio	164
Evan, Doubleday, Papilio	110	Helus, Cramer, Papilio	40
Evelina, Stoll, Papilio	193	HEMARIS, Dalman	260
Fabius, Fabr., Papilio	204	HERACLIDES, Hübner	89
Faro, Cramer, Sphinx	263	Heri, Moore, Mycalesis	233
Fatih, Kollar, Hesperia	252	HERONA, Doubleday	119
Fatna, Boisd., Emesia	243	Hesione, Cramer, Papilio	234
FAUNIS, Hübner	213	Hespera, Fabr., Sphinx	269
Fausta, Olivier, Pieris	68	HESPERIA, Fabr.	252
Flacilla, Doubleday, Vanessa	137	HESTIA, Hübner	121, 134
Flegyas, Cramer, Papilio	242	HESTINA, Westwood	160
Folus, Cramer, Papilio	252	Hierta, Fabr., Papilio	140
Francie, G. R. Gray, Adolias	197	Hierte, Hübner, Pontia	81
Francisca, Cramer, Papilio	230	Hilaria, Cramer, Papilio	57
Francki, Godart, Nymphalis	148	Hilda, Westwood, Neorina	226
Fylla, Boisd., Emesia	244	Himachala, Moore, Theope	234
Gambrius, Fabr., Papilio	148	HIPPARCHIA, Fabr.	228
Gana, Moore, Goniloba	245	Hippia, Fabr., Pieris	61
Ganesa, Doubleday, Papilio	107	HIPPO, Hübner	221, 222
— Moore, Amblypodia	44	Hippo, Cramer, Papilio	71
Ganescha, Kollar, Amathusia	146	HIPPOTION, Hübner	274

Hordonia, <i>Stoll</i> , <i>Papilio</i>	p. 164	Issaea, <i>G. R. Gray</i> , <i>Argynnus</i>	p. 156
Horsfieldii, <i>G. R. Gray</i> , <i>Pieris</i> ..	78	ISSORIA, <i>Hübner</i>	155
— <i>Swainson</i> , <i>Drusilla</i>	214	Issoria, <i>Hübner</i> , <i>Telechinia</i>	136
— <i>De Cerisy</i> , <i>Smerinthus</i>	264	Iswara, <i>White</i> , <i>Papilio</i>	101
— <i>Moore</i> , <i>Miletus</i>	19	ITANUS, <i>Doubleday</i>	185
Hübneri, <i>Moore</i> , <i>Euploea</i>	128	ITHOBALUS, <i>Hübner</i>	89
Hügeli, <i>Kollar</i> , <i>Paphia</i>	208	Iva, <i>Moore</i> , <i>Adolias</i>	195
HYADES, <i>Boisd.</i>	214	IXIAS, <i>Hübner</i>	65
Hyagriva, <i>Moore</i> , <i>Yphthima</i>	236	Jacintha, <i>Fabr.</i> , <i>Papilio</i>	159
Hyale, <i>Linn.</i> , <i>Papilio</i>	62	Jacquemontii, <i>Boisd.</i> , <i>Satyrus</i>	229
Hyas, <i>Boisd.</i> , <i>Lophura</i>	263	Jafra, <i>Godart</i> , <i>Myrina</i>	47
Hylas, <i>Hübner</i> , <i>Marpesia</i>	146	Jahnu, <i>Moore</i> , <i>Adolias</i>	192
— <i>Linn.</i> , <i>Sphinx</i>	260	Jaina, <i>Moore</i> , <i>Ismene</i>	247
Hylax, <i>Fabr.</i> , <i>Hesperia</i>	19	Jalindra, <i>Horsf.</i> , <i>Amblypodia</i>	50
HYLES, <i>Hübner</i>	273	Janaka, <i>Moore</i> , <i>Papilio</i>	97
HYLOCUS, <i>Hübner</i>	268	Janardana, <i>Moore</i> , <i>Mycalesis</i>	234
HYPANARTIA, <i>Hübner</i>	152	Jangala, <i>Horsf.</i> , <i>Amblypodia</i>	46
HYPANIS, <i>Boisd.</i>	239	Japetus, <i>Cramer</i> , <i>Papilio</i>	245
Hyparete, <i>Linn.</i> , <i>Papilio</i>	80	Jarbas, <i>Fabr.</i> , <i>Hesperia</i>	32
Hypatada, <i>Boisd.</i> , <i>Thecla</i>	45	JASIA, <i>Swainson</i>	204
Hypenor, <i>Godart</i> , <i>Papilio</i>	95	Jasonia, <i>Westwood</i> , <i>Hestia</i>	135
Hyperbius, <i>Linn.</i> , <i>Papilio</i>	155	JASONIDES, <i>Hübner</i>	89
HYPOLIMNAS, <i>Hübner</i>	158	Jayadeva, <i>Moore</i> , <i>Ismene</i>	248
Hypomenes, <i>Kaden</i> , <i>Argynnus</i>	198	Jina, <i>Moore</i> , <i>Athyma</i>	172
Hypotheüs, <i>Cramer</i> , <i>Sphinx</i>	271	Judith, <i>Fabr.</i> , <i>Papilio</i>	77
Hypoocla, <i>Cramer</i> , <i>Papilio</i>	153	Jugurtha, <i>Cramer</i> , <i>Papilio</i>	57
Hypselis, <i>Godart</i> , <i>Vanessa</i>	153	Juliana, <i>Cramer</i> , <i>Papilio</i>	145
Iapis, <i>Godart</i> , <i>Nymphalis</i>	192	Jumba, <i>Moore</i> , <i>Neptis</i>	167
Ida, <i>Cramer</i> , <i>Papilio</i>	142	JUNONIA, <i>Hübner</i>	139
IDAIDES, <i>Hübner</i>	89	Justinia, <i>Cramer</i> , <i>Papilio</i>	231
Idamene, <i>Godart</i> , <i>Vanessa</i>	143	Juventa, <i>Cramer</i> , <i>Papilio</i>	122
IDEA, <i>Fabr.</i>	134	Jynx, <i>Hübner</i> , <i>Elmynias</i>	237
Idea, <i>Stoll</i> , <i>Papilio</i>	135	KALLIMA, <i>Doubleday</i>	208
IDEOPSIS, <i>Horsf.</i>	133	Kamala, <i>Moore</i> , <i>Argynnus</i>	156
IDMAIS, <i>Boisd.</i>	67	Kamara, <i>Moore</i> , <i>Elymnias</i>	239
ILERDA, <i>Doubleday</i>	28	Kandarpa, <i>Horsf.</i> , <i>Lycæna</i>	23
ILIADES, <i>Hübner</i>	89	Kansa, <i>Moore</i> , <i>Debis</i>	220
Ilithya, <i>Drury</i> , <i>Papilio</i>	239	Kessuma, <i>Horsf.</i> , <i>Thecla</i>	37
Imperialis, <i>Hope</i> , <i>Teinopalpus</i> ..	86	Klugii, <i>Moore</i> , <i>Euploea</i>	130
INACHIS, <i>Hübner</i>	136	Klugius, <i>Zinken-Sommer</i> , <i>Morpho</i> ..	215
Inachis, <i>Boisd.</i> , <i>Amathusia</i>	208	Kotzebæus, <i>Esch.</i> , <i>Papilio</i>	95
Inara, <i>Doubleday</i> , <i>Limenitis</i>	173	Krishna, <i>Moore</i> , <i>Papilio</i>	108
Inaria, <i>Cramer</i> , <i>Papilio</i>	158	Kumara, <i>Moore</i> , <i>Hesperia</i>	255
Indica, <i>Doubleday</i> , <i>Pieris</i>	61	Ladon, <i>Cramer</i> , <i>Papilio</i>	248
Indra, <i>Moore</i> , <i>Pieris</i>	74	LAERTIADES, <i>Hübner</i>	89
Indrani, <i>Moore</i> , <i>Hesperia</i>	253	Læta, <i>Boisd.</i> , <i>Terias</i>	65
Infernalis, <i>Westwood</i> , <i>Sesia</i>	261	Lais, <i>Linn.</i> , <i>Papilio</i>	237
Inquilina, <i>Harris</i> , <i>Deilephila</i>	274	Laius, <i>Cramer</i> , <i>Papilio</i>	21
Insularis, <i>Horsf.</i> , <i>Phædra</i>	53	LAMPROSURA, <i>Boisd.</i>	85
IPHIAS, <i>Boisd.</i>	60	Laodocus, <i>De Haan</i> , <i>Papilio</i>	91
IPHICLIDES, <i>Hübner</i>	85, 89	LAOGONA, <i>Boisd.</i>	152
Iphigenia, <i>Cramer</i> , <i>Papilio</i>	159	Laomedia, <i>Linn.</i> , <i>Papilio</i>	140
Iphita, <i>Cramer</i> , <i>Papilio</i>	143	Laomedon, <i>Cramer</i> , <i>Papilio</i>	100
Irava, <i>Moore</i> , <i>Hesperia</i>	254	Lapithis, <i>Boisd.</i> , <i>Thecla</i>	48
Isa, <i>Moore</i> , <i>Hestina</i>	161	Larymna, <i>Doubleday</i> , <i>Limenitis</i> ..	172
Isama, <i>Kollar</i> , <i>Satyrus</i>	217	Lasinassa, <i>Cramer</i> , <i>Papilio</i>	159
ISMENE, <i>Swainson</i>	247	LASIOMMATA, <i>Westwood</i>	227
Ismene, <i>Cramer</i> , <i>Papilio</i>	223	Lathonia, <i>Kollar</i> , <i>Argynnus</i>	156
— <i>Doubleday</i> , <i>Limenitis</i>	180	Lathyrus, <i>Boisd.</i> , <i>Deilephila</i>	274
Isocrates, <i>Fabr.</i> , <i>Hesperia</i>	33	Latreillei, <i>Donovan</i> , <i>Papilio</i>	97
ISOPLES, <i>Hübner</i>	274	Lea, <i>Doubleday</i> , <i>Pieris</i>	77

Leda, <i>Linn.</i> , <i>Papilio</i>	p. 222	Malaya, <i>Horsf.</i> , <i>Lycæna</i>	p. 26
Ledeckbouria, <i>Esch.</i> , <i>Papilio</i>	104	Malika, <i>Horsf.</i> , <i>Thecla</i>	37
Leis, <i>Hübner</i> , <i>Catopthaga</i>	37	Malsara, <i>Moore</i> , <i>Mycalesis</i>	231
Lemonias, <i>Linn.</i> , <i>Papilio</i>	139	Mamerta, <i>Cramer</i> , <i>Papilio</i>	231
Leonteus, <i>Zinken-Sommer</i> , <i>Morpho</i>	213	Manasa, <i>Moore</i> , <i>Neptis</i>	165
Leos, <i>Gérvin</i> , <i>Symmachus</i>	18	MANCIPUM, <i>Horsf.</i>	67, 68
Lepida, <i>Moore</i> , <i>Eunispe</i>	213	Mandata, <i>Moore</i> , <i>Mycalesis</i>	233
Lepita, <i>Moore</i> , <i>Libythea</i>	240	Mangala, <i>Moore</i> , <i>Hesperia</i>	255
LEPTOCIRCUS, <i>Swainson</i>	85	Manilia, <i>Cramer</i> , <i>Papilio</i>	159
LEPTOSIA, <i>Hübner</i>	63	Marathus, <i>Doubleday</i> , <i>Herona</i>	199
LETHE, <i>Hübner</i>	216	Marianne, <i>Cramer</i> , <i>Papilio</i>	66
Lethe, <i>Westwood</i> , <i>Acherontia</i>	266	Marmax, <i>Westwood</i> , <i>Charaxes</i>	206
LETHIA, <i>Hübner</i>	267	MARMESSUS, <i>Hübner</i>	51
Leucocerca, <i>Kollar</i> , <i>Hesperia</i>	252	Matuta, <i>Hübner</i> , <i>Acca</i>	169
Leucocyma, <i>Godart</i> , <i>Biblis</i>	238	Mazares, <i>Doubleday</i> , <i>Euploea</i>	127
LEUCONIA, <i>Donzel</i>	70	Medusa, <i>De Cerisy</i> , <i>Acherontia</i>	267
Leuconoë, <i>Erichson</i> , <i>Hestia</i>	135	Megarus, <i>Westwood</i> , <i>Papilio</i>	90
LEUCOPHLEBIA, <i>Westwood</i>	265	Meges, <i>Zinken-Sommer</i> , <i>Papilio</i>	85
Leucothoë, <i>Linn.</i> , <i>Papilio</i>	169	Mekara, <i>Moore</i> , genus	219
LEXIAS, <i>Boisd.</i>	203	MELAMPIAS, <i>Hübner</i>	226
Libitina, <i>Godart</i> , <i>Pieris</i>	73	Melampus, <i>Cramer</i> , <i>Papilio</i>	32
LIBYTHEA, <i>Fabr.</i>	239	Melaneus, <i>Cramer</i> , <i>Papilio</i>	123
Libythea, <i>Fabr.</i> , <i>Papilio</i>	73	Melanides, <i>De Haan</i> , <i>Papilio</i>	91
Lignaria, <i>Walker</i> , <i>Panaera</i>	270	Melanippus, <i>Cramer</i> , <i>Papilio</i>	125
LIMENITIS, <i>Fabr.</i>	163, 170, 178	MELANITIS, <i>Fabr.</i>	222, 236
Limniace, <i>Cramer</i> , <i>Papilio</i>	121	Melina, <i>Godart</i> , <i>Danais</i>	128
Lineata, <i>Fabr.</i> , <i>Sphinx</i>	274	Melissa, <i>Cramer</i> , <i>Papilio</i>	121
—, <i>Westwood</i> , <i>Leucophlebia</i>	265	Melitta, <i>Cramer</i> , <i>Papilio</i>	159
Linoesa, <i>Walker</i> , <i>Chærocampa</i>	276	Memercus, <i>Godart</i> , <i>Papilio</i>	89
Liria, <i>Fabr.</i> , <i>Papilio</i>	159	Memnon, <i>Linn.</i> , <i>Papilio</i>	99
Lisarda, <i>Doubleday</i> , <i>Diadema</i>	160	Menaka, <i>Moore</i> , <i>Goniloba</i>	246
Lisias, <i>Fabr.</i> , <i>Papilio</i>	48	MENELAIDES, <i>Hübner</i>	89
Livornica, <i>Esper</i> , <i>Sphinx</i>	274	Menetho, <i>Fabr.</i> , <i>Papilio</i>	211
Lohita, <i>Horsf.</i> , <i>Amblypodia</i>	38	MERINTHUS, <i>Meigen</i>	264
Longinus, <i>Fabr.</i> , <i>Hesperia</i>	45	Merione, <i>Cramer</i> , <i>Papilio</i>	144
LOPHURA, <i>Boisd.</i>	263	Mesentina, <i>Cramer</i> , <i>Papilio</i>	72
Lotis, <i>Cramer</i> , <i>Papilio</i>	125	MESSARAS, <i>Doubleday</i>	150
LOXURA, <i>Horsf.</i>	51	Mestor, <i>Hübner</i> , <i>Iliades</i>	99
LUBENTINA, <i>Cramer</i> , <i>Papilio</i>	188	METOPSILUS, <i>Hübner</i>	274
Lucasi, <i>Boisd.</i> , <i>Deilephila</i>	277	Miah, <i>Moore</i> , <i>Neptis</i>	164
Lucina, <i>Cramer</i> , <i>Papilio</i>	153	Midamis, <i>Hübner</i> , <i>Limnas</i>	131
Lucipor, <i>Westwood</i> , <i>Thaumantis</i>	215	Midamus, <i>Linn.</i> , <i>Papilio</i>	133
Lutca, <i>Zinken-Sommer</i> , <i>Amathusia</i>	146	MLETUS, <i>Hübner</i>	18
Luxerii, <i>Hübner</i> , <i>Zeuxidia</i>	210	MIMAS, <i>Hübner</i>	264
LYCÆNA, <i>Fabr.</i>	19, 20, 21, 27	Minereus, <i>G. R. Gray</i> , <i>Papilio</i>	97
Lycetus, <i>Cramer</i> , <i>Sphinx</i>	277	MINETRA, <i>Boisd.</i>	147
Lycô, <i>Doubleday</i> , <i>Amblypodia</i>	44	Mineus, <i>Linn.</i> , <i>Papilio</i>	231
Lycorias, <i>Doubleday</i> , <i>Rhodocera</i>	59	MINOIS, <i>Hübner</i>	228
Lynceus, <i>Drury</i> , <i>Papilio</i>	134	Minos, <i>Cramer</i> , <i>Papilio</i>	87
Lyneida, <i>Cramer</i> , <i>Papilio</i>	71	Misippus, <i>Linn.</i> , <i>Papilio</i>	158
Lysandra, <i>Cramer</i> , <i>Papilio</i>	235	Mitra, <i>Moore</i> , <i>Euploea</i>	127
Macareus, <i>Godart</i> , <i>Papilio</i>	90	MŒRA, <i>Hübner</i>	209, 211
Machaon, <i>Linn.</i> , <i>Papilio</i>	111	Moorei, <i>Boisd.</i> , <i>Ambulyx</i>	266
MACROGLOSSA, <i>Ochsenheimer</i>	261	Morgiana, <i>Westwood</i> , <i>Apatura</i>	201
MACROGLOSSUM, <i>Scopoli</i>	261	Morpheus, <i>Cramer</i> , <i>Sphinx</i>	269
MACROSILA, <i>Boisd.</i>	268	Morta, <i>Hübner</i> , <i>Acherontia</i>	267
Maculatum, <i>Horsf.</i> , <i>Macroglossum</i>	262	Mulciber, <i>Cramer</i> , <i>Papilio</i>	133
Madhava, <i>Moore</i> , <i>Goniloba</i>	246	MURTIA, <i>Hübner</i>	55
Madura, <i>Horsf.</i> , <i>Hipparchia</i>	229	Mutius, <i>Fabr.</i> , <i>Papilio</i>	102
Mahesa, <i>Moore</i> , <i>Athyma</i>	176	MYCALESIAS, <i>Hübner</i>	230
Makuta, <i>Horsf.</i> , <i>Hipparchia</i>	226	Mycena, <i>Cramer</i> , <i>Papilio</i>	223

- MYLOTHRIS, *Hübner* p. 60, 70
 MYRINA, *Godart* 47, 51
 Myrrha, *Godart*, *Libythea* 240
 Nada, *Moore*, *Debis* 218
 Naga, *Moore*, *Philampelus* 271
 Nakula, *Moore*, *Apatura* 203
 Nama, *Doubleday*, *Pieris* 76
 —, *Boisd.*, *Diadema* 162
 Namouna, *Doubleday*, *Pieris* 71
 —, *Doubleday*, *Apatura* 201
 Nandina, *Moore*, *Neptis* 168
 Narada, *Horsf.*, *Amblypodia* 39
 Naranata, *Moore*, *Pamphila* 251
 Narasingha, *Moore*, *Yphthima* 236
 Nasaka, *Horsf.*, *Thecla* 37
 Nata, *Moore*, *Neptis* 168
 Neduna, *De Haan*, *Cyrestis* 147
 Nedymond, *Cramer*, *Papilio* 50
 Nesiuss, *Zinken-Sommer*, *Papilio* 89
 Nefte, *Cramer*, *Papilio* 173
 Neilgheriensis, *Guérin*, *Satyrus* 217
 Nemertes, *Hübner*, *Salpinx* 129
 Neombo, *Boisd.*, *Pieris* 72
 Neophron, *Boisd.*, *Emesis* 243
 NEORINA, *Westwood* 225
 NEPIELE, *Hübner* 268
 Nephtie, *Fabr.*, *Papilio* 56
 Nepos, *Fabr.*, *Hesperia* 245
 NEPTIS, *Fabr.* 163
 Neriene, *Fischer*, *Colias* 62
 Nerii, *Linn.*, *Sphinx* 272
 Nerina, *Fabr.*, *Papilio* 159
 Nerissa, *Fabr.*, *Papilio* 72
 Nero, *Fabr.*, *Papilio* 70
 Nessimachus, *Boisd.*, *Adolias* 197
 Nessus, *Drury*, *Sphinx* 276
 NESTORIDES, *Hübner* 89
 Niamus, *Godart*, *Papilio* 155
 Nicea, *G. R. Gray*, *Adolias* 197
 Nila, *Horsf.*, *Lycæna* 21
 —, *Kollar*, *Thecla* 44
 NINA, *Horsf.* 69
 Nina, *Fabr.*, *Papilio* 69
 Nipalensis, *G. R. Gray*, *Pieris* 75
 —, *Doubleday*, *Gonepteryx* 59
 Niphe, *Linn.*, *Papilio* 155
 NISONIADES, *Hübner* 250
 Nissa, *Kollar*, *Thecla* 46
 Nivca, *Zinken-Sommer*, *Amathusia* 146
 Nomius, *Esper*, *Papilio* 115
 Nourmahal, *Westwood*, *Thaumantis* 215
 Noureddin, *Westwood*, *Thaumantis* 215
 Nox, *Swainson*, *Papilio* 89
 Numa, *Weber*, *Papilio* 103
 NYCTALEMON, *Dalman* 256
 Nycteris, *Kollar*, *Macroglossa* 263
 Nyctiphanes, *Boisd.*, *Mecrosila* 268
 Nymphalides, *Swains.*, *Amphrisius* 87
 NYMPHALIS, *Latr.* 158, 178, 185, 204
 Nyseus, *Guérin*, *Polyommatus* 27
 Ochsenheimeri, *Moore*, *Euploea* 132
- Ocyale, *Hübner*, *Junonia* p. 141
 Ocys, *Hübner*, *Hippotion* 274
 Odana, *Godart*, *Morpho* 214
 Edipodea, *Swainson*, *Ismene* 247
 Oeneis, *Hübner* 228
 Enome, *Linn.*, *Papilio* 140
 Oldenlandiæ, *Fabr.*, *Sphinx* 278
 Onyx, *Boisd.*, *Thecla* 30
 Opalina, *Kollar*, *Limenitis* 171
 OREAS, *Hübner* 216, 221
 OREINA, *Westwood* 226
 Orestes, *Fabr.*, *Papilio* 115
 OREUS, *Hübner* 274
 ORINOMA, *Doubleday* 225
 Orithyia, *Linn.*, *Papilio* 141
 ORNEUS, *Hübner* 270
 ORNITHOPTERA, *Boisd.* 86
 ORONTES, *Swainson* 256
 ORPHEIDES, *Hübner* 89
 Otreia, *Cramer*, *Papilio* 230
 Padma, *Kollar*, *Satyrus* 228
 Paduka, *Moore*, *Limenitis* 179
 Palephates, *Westwood*, *Papilio* 92
 Palguna, *Moore*, *Adolias* 190
 Palicosta, *Boisd.*, *Deilephila* 276
 Pammom, *Linn.*, *Papilio* 104
 PAMPHILA, *Fabr.* 250
 PANACRA, *Walker* 269
 Panava, *Westwood*, *Thecla* 28
 Panda, *Godart*, *Pieris* 71
 Pandava, *Horsf.*, *Lycæna* 22
 Pandia, *Moore*, *Hesperia* 254
 Pandione, *Hübner*, *Pieris* 73
 PANDITA, *Moore* 181
 Pandoces (*Boisd.*), *Moore*, *Yphthima* 235
 —, *ma* 235
 Pandu, *Horsf.*, *Symetha* 18
 Pann, *Fabr.*, *Hesperia* 34
 Panope, *Linn.*, *Papilio* 92
 PANOPEA, *Hübner* 158
 Panopus, *Cramer*, *Sphinx* 266
 PAONIAS, *Hübner* 264
 PAPHIA, *Fabr.* 145, 204
 PAPILIO, *Linn.* 89
 Paralekta, *Horsf.*, *Paphia* 208
 Parallelis, *Horsf.*, *Smerinthus* 264
 PARARGE, *Hübner* 227
 PAREBA, *Doubleday* 136
 PARIDES, *Hübner* 89
 Paris, *Linn.*, *Papilio* 107
 Parisatis, *Westwood*, *Apatura* 202
 PARNASSIUS, *Latr.* 84
 Parrhasius, *Fabr.*, *Hesperia* 22
 Parryæ, *Hope*, *Teinopalpus* 86
 Parta, *Moore*, *Adolias* 185
 PARTHENOS, *Hübner* 147
 Parvata, *Moore*, *Apatura* 202
 Pasithœ, *Linn.*, *Papilio* 79
 Passalus, *Drury*, *Sphinx* 262
 Patala, *Kollar*, *Adolias* 196
 Patalena, *Westwood*, *Amathusia* 210

Patnia, <i>Westwood</i> , Melanitis	...	p. 238	Polyuectes, <i>Doubleday</i> , Papilio	...	p. 96
Patnia, <i>Moore</i> , Mycalesis	...	232	POLYGONIA, <i>Hübner</i>	...	137
Patroclus, <i>Linn.</i> , Papilio	...	256	Polymnestor, <i>Cramer</i> , Papilio	...	101
Paulina, <i>Cramer</i> , Papilio	...	73	Polynice, <i>Cramer</i> , Papilio	...	114
Pavana, <i>Horsf.</i> , Lycæna	...	24	Polynissa, <i>Godart</i> , Vanessa	...	144
— <i>Kollar</i> , Polyommatus	...	28	POLYOMMATUS, <i>Latr.</i>	...	20, 21, 27, 30
Pavetia, <i>Zinken-Sommer</i> , Euploea	...	131	POLYPTYCHUS, <i>Hübner</i>	...	264
Payeni, <i>Boisd.</i> , Papilio	...	110	Polytes, <i>Linn.</i> , Papilio	...	103
PENTHEMA, <i>Doubleday</i>	...	160	Polyzena, <i>Donovan</i> , Papilio	...	170
Penthesilea, <i>Cramer</i> , Papilio	...	154	— <i>Cramer</i> , Papilio	...	207
Peranthus, <i>Fabr.</i> , Papilio	...	109	Polyxo, <i>Godart</i> , Nymphalis	...	207
PERGESA, <i>Walker</i>	...	272	Pomona, <i>Fabr.</i> , Papilio	...	57
Periander, <i>Fabr.</i> , Papilio	...	147	Pompeus, <i>Cramer</i> , Papilio	...	87
Perimele, <i>Cramer</i> , Papilio	...	159	Pompilius, <i>Fabr.</i> , Papilio	...	116
Perimuta, <i>Boisd.</i> , Amblypodia	...	42	PONTIA, <i>Fabr.</i>	...	64, 69, 70
PERRYHYBRIS, <i>Hübner</i>	...	70	Porphyria, <i>Cramer</i> , Papilio	...	159
Persimilis, <i>Westwood</i> , Hestina	...	161	Porsema, <i>Cramer</i> , Papilio	...	80
PHÆDRA, <i>Horsf.</i>	...	52	Praba, <i>Moore</i> , Hesperia	...	253
PHÆDRUS, <i>Fabr.</i> , Papilio	...	53	Pralaya, <i>Moore</i> , Goniloba	...	246
PHALANTA, <i>Horsf.</i>	...	151	Pravara, <i>Moore</i> , Athyma	...	173
Phalanta, <i>Drury</i> , Papilio	...	151	PRECIS, <i>Hübner</i>	...	142
Phaon, <i>Erichson</i> , Drusilla	...	214	PRIAMIDES, <i>Hübner</i>	...	89
Pharis, <i>Doubleday</i> , Myrina	...	48	Primaria, <i>Horsf.</i> , Aconthea	...	185
Phedima, <i>Cramer</i> , Papilio	...	223	Procris, <i>Cramer</i> , Papilio	...	179
Phemius, <i>Doubleday</i> , Itanus	...	187	Proserpina, <i>Cramer</i> , Papilio	...	159
Pheres, <i>Boisd.</i> , Pamphila	...	251	PROTEIDES, <i>Hübner</i>	...	244
Phidippus, <i>Linn.</i> , Papilio	...	209	Protenor, <i>Cramer</i> , Papilio	...	98
PHILAMPELUS, <i>Harris</i>	...	270	PROTESILAUS, <i>Swainson</i>	...	89
Philene, <i>Cramer</i> , Papilio	...	124	PROTHOE, <i>Hübner</i>	...	118
Philippinus, <i>Cramer</i> , Papilio	...	56	Prothoë, <i>Godart</i> , Danais	...	130
Philomela, <i>Fabr.</i> , Papilio	...	61	Protogenia, <i>Cramer</i> , Papilio	...	237
— <i>Hübner</i> , Ypthima	...	235	Pseudocentaurus, <i>Doubleday</i> , Ani-	...	
— <i>Zinken-Sommer</i> , Euploea	...	123	blypoda	...	40
Philoxenus, <i>G. R. Gray</i> , Papilio	...	96	Pseudolonginus, <i>Doubleday</i> , Am-	...	
Phisadia, <i>Godart</i> , Pieris	...	68	blypoda	...	45
Phleas, <i>Linn.</i> , Papilio	...	27	Pseudoroxus, <i>Doubleday</i> , Lycæna	...	27
PHLEGETRONTIUS, <i>Hübner</i>	...	268	PSITHYROS, <i>Hübner</i>	...	261
Phœbis, <i>Hübner</i>	...	55	Pulaha, <i>Moore</i> , Enope	...	227
PHOLUS, <i>Hübner</i>	...	265, 270	Pulasara, <i>Moore</i> , Adolias	...	190
Pholus, <i>Boisd.</i> , Thecla	...	30	Pulomaya, <i>Moore</i> , Hesperia	...	252
Phryne, <i>Fabr.</i> , Papilio	...	71	Purendra, <i>Moore</i> , Pyrgus	...	250
Phryxe, <i>Boisd.</i> , Pieris	...	83	Puseda, <i>Moore</i> , Adolias	...	190
PIRYXUS, <i>Hübner</i>	...	273	Puspa, <i>Horsf.</i> , Polyommatus	...	20
Picus, <i>Cramer</i> , Sphinx	...	260	Putra, <i>Moore</i> , Hesperia	...	253
PIERIS, <i>Fabr.</i>	...	65, 70, 83, 84	PYRAMEIS, <i>Hübner</i>	...	138
Pindarus, <i>Fabr.</i> , Hesperia	...	37	Pyranthe, <i>Linn.</i> , Papilio	...	56
Pita, <i>Horsf.</i> , Luxura	...	52	Pyrene, <i>Linn.</i> , Papilio	...	66
PITHECOPS, <i>Horsf.</i>	...	19	PYRGUS, <i>Hübner</i>	...	249
Plautilla, <i>Hübner</i> , Papilio	...	169	PYRONIA, <i>Hübner</i>	...	228
Plexippus, <i>Linn.</i> , Papilio	...	124	Pyrrhus, <i>Donovan</i> , Papilio	...	205
Plinius, <i>Fabr.</i> , Hesperia	...	25	Quercetum, <i>Boisd.</i> , Amblypodia	...	42
Pluto, <i>Fabr.</i> , Hesperia	...	21	Quercetorum, <i>Boisd.</i> , Amblypodia	...	42
— <i>Fabr.</i> , Sphinx	...	276	Radha, <i>Moore</i> , Neptis	...	166
PODALIRIUS, <i>Swainson</i>	...	89	Rafflesii, <i>Horsf.</i> , Deilephila	...	276
Polibetes, <i>Cramer</i> , Papilio	...	209	RAGADIA, <i>Westwood</i>	...	226
Polinice, <i>Boisd.</i> , Hypanis	...	239	Rahria, <i>Westwood</i> , Cyrestis	...	147
Pollux, <i>Westwood</i> , Papilio	...	93	Rama, <i>Kollar</i> , Thecla	...	42
Polyctor, <i>Boisd.</i> , Papilio	...	109	Ramdeo, <i>Moore</i> , Thaumantis	...	215
Polydecta, <i>Fabr.</i> , Papilio	...	230	Ranga, <i>Moore</i> , Athyma	...	175
POLYDORUS, <i>Swainson</i>	...	89	Ravana, <i>Moore</i> , Papilio	...	96
Polydorus, <i>Cramer</i> , Papilio	...	94	— <i>Moore</i> , Cirrochroa	...	150

- Ravi, *Moore*, Goniloba p. 246
 Ravindra, *Horsf.*, Myrina 51
 Recaranus, *Westwood*, Cyrestis .. 146
 Remba, *Moore*, Pieris 75
 Rhadamanthus, *Fabr.*, Papilio .. 126
 — *Boisd.*, Ornithoptera 88
 Rhadamina, *Godart*, Danais 126
 Rhebus, *De Cerasy*, Macroglossa .. 263
 Rhetenor, *Westwood*, Papilio 98
 Rhexia, *Fabr.*, Papilio 67
 RHODOCERA, *Boisd.* 59
 Richmondia, *G. R. Gray*, Papilio .. 86
 Risa, *Doubleday*, Cyrestis 147
 Rochana, *Horsf.*, Amblypodia .. 44
 Rohria, *Fabr.*, Papilio 217
 Romulus, *Cramer*, Papilio 102
 Rouxii, *Boisd.*, Pieris 71
 Roxus, *Godart*, Polyommatus 26
 Rudra, *Moore*, Argynnis 156
 Runeka, *Moore*, Mycalesis 234
 Safitza, *Hewitson*, Mycalesis .. 233
 Sakontala, *Kollar*, Argynnis 157
 Sakra, *Moore*, Yphthima 236
 SALAMIS, *Boisd.* 142
 Salia, *Moore*, Adolias 189
 SALPINX, *Hübner* 126
 Salsala, *Moore*, Nisoniades 250
 Samba, *Moore*, Mycalesis 233
 Sambara, *Moore*, Goniloba 246
 Sanaca, *Moore*, Pieris 79
 Sanatana, *Moore*, Mycalesis .. 231
 Sancara, *Moore*, Adolias 195
 Santana, *Moore*, Anops 54
 Saraswati, *Kollar*, Satyrus 229
 Sari, *Horsf.*, Terias 64
 Sarpedon, *Linn.*, Papilio 113
 Sataas, *Boisd.*, Acherontia 267
 SATASPE, *Moore* 261
 SATYRUS, *Latr.* 228
 Scanda, *Kollar*, Erebia 226
 — *Moore*, Debis 218
 Scapularis, *Horsf.*, Deilephila .. 269
 Schakra, *Kollar*, Satyrus 227
 Schreiberi, *Godart*, Nymphalis .. 205
 Scylla, *Linn.*, Papilio 58
 Selenophora, *Kollar*, Limenitis .. 175
 Sema, *Moore*, Hesperia 253
 Sena, *Kollar*, Polyommatus 28
 — *Moore*, Goniloba 245
 Sericeus, *Walker*, Philampelus .. 271
 SESIA, *Fabr.* 260
 Sesia, *Fabr.*, Papilio 66
 Seta, *Moore*, Pieris 78
 Setho, *Doubleday*, Myrina 33
 Sidera, *Boisd.*, Adolias 194
 Sikandi, *Moore*, Adolias 193
 Silhetensis, *Boisd.*, Amblypodia .. 40
 — *Boisd.*, Deilephila 278
 Similis, *Linn.*, Papilio 122
 Sinha, *Kollar*, Termos 152
 Sinope, *Boisd.*, Adolias 182
 Sinta, *Moore*, Ismene p. 248
 SITHON, *Hübner* 30
 Sitiene, *Boisd.*, Macroglossa .. 262
 Siva, *Westwood*, Adolias 198
 SMERINTHUS, *Latr.* 264
 Soracta, *Boisd.*, Pieris 83
 Solandra, *Fabr.*, Papilio 222
 Sorya, *Kollar*, Thecla 32
 SPHINX, *Linn.* 267
 Sphyrus, *Hübner*, Papilio 111
 Spilota, *Horsf.*, Deilephila .. 277
 Stellarum, *Linn.*, Sphinx 261
 STEROPES, *Hübner* 250
 Stichius, *Hübner*, Papilio 103
 Striatus, *Zinken-Sommer*, Papilio .. 90
 Styx, *Westwood*, Acherontia 266
 Substrigilis, *Westwood*, Ambulyx .. 266
 Sufusa, *Walker*, Deilephila .. 275
 Sugriva, *Horsf.*, Amblypodia .. 51
 Sumbadra, *Moore*, Pamphila 251
 Superba, *Herbst*, Papilio 131
 Superna, *Moore*, Pyrgus 249
 Sura, *Doubleday*, Zophoessa .. 220
 — *Moore*, Achlyodes 251
 Suradeva, *Moore*, Melanitis 225
 Surya, *Moore*, Loxura 52
 Sutapa, *Moore*, Hesperia 255
 Suyudana, *Moore*, Melanitis 224
 Swaha, *Kollar*, Satyrus 229
 Syama, *Horsf.*, Amblypodia .. 38
 Syla, *Kollar*, Thecla 30
 Sylvia, *Cramer*, Papilio 147
 Symbialis, *Hübner*, Alazonia 154
 SYMBRENTHA, *Hübner* 152
 SYMETHA, *Horsf.* 18
 Symethus, *Cramer*, Papilio 18
 SYMPHÆDRA, *Hübner* 185, 203
 SYNCHLOE, *Hübner* 70
 SYRICHTHUS, *Boisd.* 249
 Talazala, *Boisd.*, Thecla 39
 TALIDES, *Hübner* 244
 Tamu, *Kollar*, Polyommatus 28
 Tapania, *Moore*, Goniloba 246
 TAXILA, *Doubleday* 242
 TEINOPALpus, *Hope* 86
 TELCHINIA, *Hübner* 135
 TELEGONUS, *Hübner* 244
 TENARIS, *Hübner* 214
 Tephnia, *Godart*, Argynnis .. 156
 TERIAS, *Swainson* 63
 TERINOS, *Boisd.* 149
 Terpsichore, *Cramer*, Papilio .. 136
 TERPSICHOIROS, *Hübner* 126
 Terricola, *Horsf.*, Phædra 53
 Teuta, *Doubleday*, Adolias 196
 Thais, *Fabr.*, Papilio 149
 THANAOs, *Boisd.* 250
 Tharis, *Hübner*, Oxylydes 47
 THAUMANTIS, *Hübner* 214
 THAUMAS, *Hübner* 273, 274
 Thea, *Godart*, Argynnis 149

THECLA, Boisd.....	p. 28	Vasudeva, Moore, Mycalesis .. p. 230	
Theophrastus, Fabr., Hesperia ..	25	Vasutana, Moore, Ismene .. 247	
THERETRA, Hübner	272, 274	Vedanga, Moore, Pamphila..... 250	
Thescus, Cramer, Papilio	94	Velleda, Cramer, Papilio..... 159	
Thessalus, Boisd., Thoræssa ..	210	Vellida, Fabr., Papilio..... 141	
THESTIAS, Boisd.....	65	Venata, Moore, Terias..... 65	
Thestylis, Doubleday, Pieris ..	78	Venilia, Godart, Pieris	67
Thetys, Drury, Papilio.....	52	Verhuellii, Van der Hoeven, Colias .. 59	
Thisbe, Cramer, Papilio	80	Verma, Kollar, Satyrus	217
THOAS, Swainson	89	Vesta, Fabr., Papilio	136
Thoas, Swainson, Polydorus ..	94	Vidura, Horsf., Amblypodia .. 45	
Thoë, Hübner, Hestia	125	Vigil, Guérin, Sphinx	270
Thoosa, Hübner, Terpsichrois ..	126	Vikasi, Horsf., Neptis	165
THRÆSSA, Boisd.....	209	Violeæ, Fabr., Papilio	135
THRACIDES, Hübner.....	244	Virgauriæ, Scop., Papilio	27
Thrax, Linn., Papilio	254	Visala, Moore, Mycalesis .. 230	
Thyelia, Linn., Sphinx	276	Vishnu, Moore, Pieris	83
—, Fabr., Papilio	204	Vivarna, Horsf., Amblypodia .. 39	
THYMELE, Boisd.....	247	Volucris, Boisd., Macroglossa .. 263	
THYMELICUS, Hübner	250	Vulcania, Godart, Vanessa .. 138	
Thyodamas, Boisd., Cyrestis ..	146	Vulcanus, Fabr., Papilio	38
Thyria, Godart, Pieris	70	XANTHIDIA, Boisd.....	63
TIGRIDIA, Hübner	204	Xanthomelas, Denis et Schiff., Papilio	137
Tilaha, Horsf., Terias	64	Xanthopogon, Kollar, Hesperia .. 248	
Timæus, Cramer, Papilio	27	XENICA, Westwood	228
Timesius, Stoll, Sphinx	264	Xenocles, Doubleday, Papilio	90
Timoleon, Stoll, Papilio	44	Xenodice, Hübner, Lycaena	243
Timora, Doubleday, Discophora ..	211	Xiphia, Fabr., Papilio	70
TINOPE, Moore	234	Xuthus, Linn., Papilio	111
TISIPHONE, Hübner	228	Yama, Moore, Zophoessa	220
Titania, Fabr., Papilio	57	YPHTHIMA, Hübner	235
Titea, Godart, Pieris	68	ZARETIS, Hübner	208
Trigerta, Moore, Adolias	191	Zayla, Doubleday, Limenitis .. 181	
Triopas, Cramer, Papilio	49	ZELIDES, Hübner	89
TROIDES, Hübner	86	ZELIMA, Fabr.....	89
Tullia, Cramer, Papilio	211	Zelima, Donovan, Papilio	143
Turcicus, Gerhard, Papilio	27	Zelmira, Cramer, Papilio	73
Tytia, G. R. Gray, Euplœa	123	ZEMEROS, Boisd.....	242
Undularis, Fabr., Papilio	237	ZERENE, Hübner	62
URANIA, Latr.	256	ZERITIS, Boisd.....	37
Urania, Zinken-Sommer, Morpho ..	214	ZERYNTHIA, Hübner	211
VALERIA, Horsf.	60	ZETIDES, Hübner	89
Valeria, Cramer, Papilio	61	ZEUXIDIA, Hübner	210
Vamana, Moore, Melanitis	223	Zeuzippe, Cramer, Papilio	77
VANESSA, Fabr.	136-9	ZONILIA, Boisd.....	268
Varaha, Moore, Melanitis	224	ZOPHOESSA, Doubleday	220
Varuna, Horsf., Thecla	37	Zulema, Doubleday, Limenitis ..	197
—, White, Papilio	98		
Vasava, Moore, Achlyodes	252		

CONTENTS OF PLATES.

PLATE I.

FIG.		PAGE.
1.	Lycaena Alexis, <i>Stoll</i> , larva, 1a. chrysalis	25
2.	Dipsas Melampus, <i>Cramer</i> , larva, 2a. chrysalis.....	32
3.	— Xenophon, <i>Fabricius</i> , larva, 3a. chrysalis.....	31
4.	Amblypodia Narada, <i>Horsfield</i> , larva, 4a. chrysalis ...	39
5.	— Apidanus, <i>Cramer</i> , larva, 5a. chrysalis ...	39
6.	— Pseudolonginus, <i>Doubleday</i> , larva, 6a. chrysalis	45
7.	Callidryas Alcemeone, <i>Cramer</i> , larva, 7a. chrysalis.....	56
8.	— Pyranthe, <i>Linnæus</i> , larva, 8a. chrysalis ...	56
9.	— Scylla, <i>Linnæus</i> , larva, 9a. chrysalis	58
10.	Hebomoia Glaucippe, <i>Linnæus</i> , larva, 10a. chrysalis	60
11.	Terias Hecabe, <i>Linnæus</i> , larva, 11a. chrysalis	63
12.	Pieris Coronis, <i>Cramer</i> , larva, 12a. chrysalis.....	76
13.	— Eucharis, <i>Drury</i> , larva, 13a. chrysalis	80
14.	— Belisama, <i>Cramer</i> , larva, 14a. chrysalis	82

PLATE II.

1.	Ornithoptera Pompeus, <i>Cramer</i> , larva, 1a. chrysalis...	87
2.	— Darsius, <i>G. R. Gray</i> , larva, 2a. chrysalis	87
3.	Papilio dissimilis, <i>Linnæus</i> , larva, 3a. chrysalis.....	91
4.	— Hector, <i>Linnæus</i> , larva, 4a. 4b. chrysalis	93
5.	— Diphilus, <i>Esper</i> , larva, 5a. chrysalis	94
6.	— Alcinous, <i>Klug</i> , larva.....	95
7.	— Memnon, <i>Linnæus</i> , larva, 7a. chrysalis.....	99

CONTENTS OF PLATES.

PLATE III.

FIG.		PAGE.
1.	Papilio Polymnestor, <i>Cramer</i> , larva, 1 α . chrysalis.....	101
2.	——— Helenus, <i>Linnæus</i> , larva, 2 α . chrysalis	101
3.	——— Polytes, <i>Linnæus</i> , larva, 3 α . chrysalis	103
4.	——— Pammon, <i>Linnæus</i> , larva, 4 α . chrysalis	104
5.	——— Demolion, <i>Cramer</i> , larva, 5 α . chrysalis	105
6.	——— Erithonius, <i>Cramer</i> , larva, 6 α . chrysalis	105
7.	——— Arjuna, <i>Horsfield</i> , larva, 7 α . chrysalis	107
8.	——— Sarpedon, <i>Linnæus</i> , larva, 8 α . chrysalis	113
9.	——— Agamemnon, <i>Linnæus</i> , larva, 9 α . chrysalis ...	114
10.	——— Antiphates, <i>Cramer</i> , larva, 10 α . chrysalis.....	116

PLATE IV.

1.	Papilio Xuthus, <i>Linnæus</i> , larva	111
2.	——— Machaon, <i>Linnæus</i> , larva, 2 α . chrysalis	111
3.	Danais Limniacæ, <i>Cramer</i> , larva, 3 α . chrysalis	121
4.	——— Juventa, <i>Cramer</i> , larva, 4 α . chrysalis	122
5.	——— Philene, <i>Cramer</i> , larva, 5 α . chrysalis.....	124
6.	——— Plexippus, <i>Linnæus</i> , larva, 6 α . chrysalis	124
7.	——— Chrysippus, <i>Linnæus</i> , larva, 7 α . chrysalis.....	126
8.	Euplœa Prothoö, <i>Godart</i> , larva, 8 α . chrysalis	130
9.	——— Core, <i>Cramer</i> , larva, 9 α . chrysalis	131
10.	——— Midamus, <i>Linnæus</i> , larva, 10 α . chrysalis	133
11.	Ideopsis Daos, <i>Boisduval</i> , larva, 11 α . chrysalis.....	134
12.	Hestia Belia, <i>Westwood</i> , chrysalis	135

CONTENTS OF PLATES.

PLATE V.

FIG.		PAGE.
1.	Telchinia Violæ, <i>Fabricius</i> , larva, 1 α . chrysalis	135
2.	Pareba Vesta, <i>Fabricius</i> , larva, 2 α . chrysalis.....	136
3.	Pyrameis Cardui, <i>Linnæus</i> , larva, 3 α . chrysalis.....	138
4.	Junonia Laomedea, <i>Linnæus</i> , larva, 4 α . chrysalis	140
5.	— Orithyia, <i>Linnæus</i> , larva, 5 α . chrysalis	141
6.	— Asterie, <i>Linnæus</i> , larva, 6 α . chrysalis	142
7.	Atella Phalanta, <i>Drury</i> , larva, 7 α . chrysalis	151
8.	Cethosia Cyane, <i>Fabricius</i> , larva, 8 α . chrysalis	155
9.	Diadema Auge, <i>Cramer</i> , larva, 9 α . chrysalis	158
10.	Neptis Aceris, <i>Esper</i> , larva, 10 α . chrysalis	168
11.	Athyma Leucothoë, <i>Linnæus</i> , larva, 11 α . chrysalis ...	170
12.	Limenitis Procris, <i>Cramer</i> , larva, 12 α . chrysalis	179

PLATE VI.

1.	Adolias Aconthea, <i>Cramer</i> , larva, 1 α . chrysalis	185
2.	— Garuda, <i>Moore</i> , larva, 2 α . chrysalis	186
3.	Nymphalis Athamas, <i>Drury</i> , larva, 3 α . chrysalis	205
4.	Amathusia Phidippus, <i>Linnæus</i> , larva, 4 α . chrysalis...	209
5.	Discophora Celinde, <i>Stoll</i> , larva, 5 α . chrysalis	211
6.	Ergolis Coryta, <i>Cramer</i> , larva, 6 α . chrysalis	244
7.	Elymnias undularis, <i>Fabricius</i> , larva, 7 α . chrysalis ...	237
8.	Melanitis Leda, <i>Linnæus</i> , larva, 8 α . chrysalis	222

CONTENTS OF PLATES.

PLATE VII.

FIG.		PAGE.
1.	Pyrgus Agama, <i>Moore</i> , larva, 1 α . chrysalis	249
2.	Goniloba Japetus, <i>Cramer</i> , larva, 2 α . chrysalis.....	245
3.	——— Badra, <i>Moore</i> , larva, 3 α . chrysalis	245
4.	Ismene Oedipodea, <i>Swainson</i> , larva, 4 α . chrysalis	247
5.	Hesperia Thrax, <i>Linnæus</i> , larva, 5 α . chrysalis	253
6.	Ismene Ambasa, <i>Moore</i> , larva, 6 α . chrysalis	248
7.	? Hesperia Folus, <i>Cramer</i> , larva, 7 α . chrysalis.....	252
8.	——— <i>Sp. undetermined</i> , larva, 8 α . chrysalis	
9.	? Ismene Ladon, <i>Cramer</i> , larva, 9 α . chrysalis	248
10.	Hesperia Pandia, <i>Moore</i> , larva, 10 α . chrysalis	254

PLATE VIII.

1.	Sesia Hylas, <i>Linnæus</i> , larva, 1 α . chrysalis	260
2.	Macroglossa corythus, <i>Boisduval</i> , larva, 2 α . chrysalis	262
3.	——— Gilia, <i>Boisduval</i> , larva, 3 α . chrysalis.....	262
4.	Lophura Hyas, <i>Walker</i> , larva, 4 α . chrysalis	263
5.	Leucophlaea lineata, <i>Westwood</i> , larva	265
6.	Calymnia Panopus, <i>Cramer</i> , larva, 6 α . chrysalis	266

PLATE IX.

1.	Acherontia Styx, <i>Westwood</i> , larva, 1 α . chrysalis	266
2.	——— Satanas, <i>Boisduval</i> , larva, 2 α . chrysalis ...	267
3.	Sphinx Convolvuli, <i>Linnæus</i> , larva, 3 α . chrysalis	267
4.	Philampelus Anceus, <i>Cramer</i> , larva, 4 α . chrysalis.....	270
5.	Panaeca scapularis, <i>Horsfield</i> , larva, 5 α . chrysalis ...	269

CONTENTS OF PLATES.

PLATE X.

FIG.		PAGE.
1.	Pergesa Acteus, <i>Cramer</i> , larva, 1a. chrysalis.....	272
2.	Darapsa Hypothoüs, <i>Cramer</i> , larva, 2a. chrysalis	271
3.	Daphnis Nerii, <i>Linnæus</i> , larva, 3a. chrysalis	272
4.	Chærocampa Alecto, <i>Linnæus</i> , larva, 4a. chrysalis ...	275

PLATE XI.

1.	Chærocampa Celerio, <i>Linnæus</i> , larva, 1a. chrysalis ...	274
2.	————— Nessus, <i>Drury</i> , larva, 2a. chrysalis	276
3.	————— Lucasi, <i>Boisduval</i> , larva, 3a. chrysalis ...	277
4.	————— Oldenlandiæ, <i>Fabr.</i> , larva, 4a. chrysalis	278
5.	————— bisecta, <i>Horsfield</i> , larva, 5a. chrysalis ...	277

PLATE XII.

1.	Polyommatus Laius, <i>Cramer</i> , larva, 1a. chrysalis	21
2.	Dipsas (biocellatus, <i>Grote</i> , MS.), larva, 2a. chrysalis	
3.	Amblypodia Timoleon, <i>Stoll</i> , larva, 3a. chrysalis	44
4.	————— <i>Spec. undetermined</i> , larva, 4a. chrysalis...	
5.	Myrina Triopas, <i>Cramer</i> , larva, 5a. chrysalis	49
6.	————— Etolus, <i>Fabricius</i> , larva, 6a. chrysalis	49
7.	Anops Thetys, <i>Drury</i> , larva, 7a. chrysalis.....	52
8.	Callidryas Philippina, <i>Cramer</i> , larva, 8a. chrysalis ...	56
9.	Pieris Mesentina, <i>Cramer</i> , larva, 9a. chrysalis	72
10.	Papilio Eurypylus, <i>Linnæus</i> , larva, 10a. chrysalis.....	113
12.	Junonia Almana, <i>Linnæus</i> , larva, 12a. chrysalis	142
13.	Adolias Lubentina, <i>Cramer</i> , larva, 13a. chrysalis	188
14.	Nymphalis Baya, <i>Moore</i> , larva, 14a. chrysalis	207
15.	Discophora Tullia, <i>Cramer</i> , larva, 15a. chrysalis	211

CONTENTS OF PLATES.

PLATE Ia.

FIG.		PAGE.
1.	<i>Miletus Boisduvali, Moore</i>	19
2.	——— <i>Horsfieldi, Moore, ♂</i>	19
3.	<i>Lycæna Parrhasius, Fabricius</i>	22
4.	<i>Ilerda Brahma, Moore, ♂</i>	29
5.	<i>Dipsas Malika, Horsfield, ♂</i>	37
6.	<i>Amblypodia Camdeo, Doubleday, ♀</i>	41
7.	——— <i>Quercetorum, Boisduval</i>	42
8.	——— <i>Dodonea, Boisduval</i>	43
9.	——— <i>Ganesa, Moore</i>	44
10.	——— <i>Rochana, Horsfield</i>	44
11.	——— <i>Jangala, Horsfield, ♂</i>	46
12.	<i>Myrina Sugriva, Horsfield</i>	51
13.	<i>Loxura Surya, Moore</i>	52
14.	<i>Anops insularis, Horsfield</i>	53

PLATE IIa.

1.	<i>Eronia Avatar, Moore</i>	61
2.	<i>Terias Venata, Moore</i>	65
3.	<i>Pieris Neombo, Boisduval</i>	72
4.	——— <i>Vishnu, Moore</i>	83
5.	——— <i>Autothisbe, Hübner</i>	82
6.	<i>Papilio Krishna, Moore</i>	108

CONTENTS OF PLATES.

PLATE IIIa.

FIG.		PAGE.
1.	Precis Hara, <i>Moore</i>	143
2.	Cyrestis Rabria, <i>Westwood</i>	147
3.	Cirrochroa Bajadeta, <i>Moore</i>	150
4.	Argynnис Aruna, <i>Moore</i>	156
5.	Calinaga Buddha, <i>Moore</i>	163

PLATE IVa.


1.	Neptis Miah, <i>Moore</i>	164
2.	— Manasa, <i>Moore</i>	165
3.	— Ananta, <i>Moore</i>	166
4.	— Radha, <i>Moore</i>	166
5.	— Jumba, <i>Moore</i>	167
6.	— Nata, <i>Moore</i>	168
7.	— Nandina, <i>Moore</i>	168


PLATE Va.

1.	Athyma Asura, <i>Moore</i>	171
2.	— Opalina, <i>Kollar</i>	171
3.	— Jina, <i>Moore</i>	172
4.	— Pravara, <i>Moore</i>	173
5.	— Cama, <i>Moore</i> , ♂ ♀	174
6.	— Ranga, <i>Moore</i>	175
7.	— Mahesa, <i>Moore</i>	176

PLATE VIa.


1.	Abrota Ganga, <i>Moore</i> , ♂ ♀	178
2.	Limenitis Danava, <i>Moore</i> , ♂ ♀	180
3.	Pandita Sinope, <i>Moore</i>	182
4.	Castalia Chandra, <i>Moore</i>	200
5.	Dilipa Morgiana, <i>Moore</i> , ♂ ♀	201
6.	Apatura Parvata, <i>Moore</i> , ♂	202
























VIII.






X.






XII









IVa



VII









SMITHSONIAN INSTITUTION LIBRARIES



3 9088 00316971 1

nhent QL545.E13

v. 1 A catalogue of the lepidopterous